

On $\mathbb{Z}/2$ -e-invariants

メタデータ	<p>言語: English</p> <p>出版者: Osaka University and Osaka City University, Departments of Mathematics</p> <p>公開日: 2024-09-09</p> <p>キーワード (Ja):</p> <p>キーワード (En):</p> <p>作成者: 南, 春男</p> <p>メールアドレス:</p> <p>所属: Osaka City University</p>
URL	https://ocu-omu.repo.nii.ac.jp/records/2008812

ON $\mathbb{Z}/2$ -e-INVARIANTS

HARUO MINAMI

(Received November 25, 1981)

Let G be the group $\mathbb{Z}/2$. Denote by $\pi_{p,q}^S$ the equivariant stable homotopy group of Landweber [12]. In a similar way to the usual e -invariants we define equivariant e -invariants e_G and $e_{G,R}$ on $\pi_{p,2q-1}^S$ by using the Adams operations in the K_G - and KO_G -theories and the equivariant Chern character. And we compute these invariants, in particular $e_{G,R}$, on the image of the equivariant J -homomorphism, making use of the Adams' result for e'_R . Here we study the case when $\widetilde{KO}_G^{-1}(\Sigma^{p,2q-1})$ is torsion-free. The torsion case is discussed by Löffler [14].

1. Definitions

Let $R^{p,q}$ denote the R^{p+q} with non trivial G -action on the first p coordinates. By $B^{p,q}$ and $S^{p,q}$ we denote the unit ball and unit sphere in $R^{p,q}$ and by $\Sigma^{p,q}$ the $B^{p,q}/S^{p,q}$. If p and q are even then $R^{p,q}$ is a complex G -module. In particular, we write 1 and L for $R^{0,2}$ and $R^{2,0}$. Then $\{1, L\}$ are basis of the complex representation ring $R(G)$ of G .

For the Thom class of $R^{2p,2q}$ as a complex G -vector bundle over a point we write $\lambda_{2p,2q}$, so that $\tilde{K}_G(\Sigma^{2p,2q}) = R(G) \cdot \lambda_{2p,2q}$ [16]. Here let $A \cdot x$ denote the module generated by x over a ring A . Then we have the formula

$$\psi^t(\lambda_{2p,2q}) = \rho^t(2p, 2q) \lambda_{2p,2q}, \quad \rho^t(2p, 2q) \in R(G)$$

for the t -th Adams operation ψ^t , and $\rho^t(2p, 2q)$ is computed briefly, using the result for ψ^t in $\tilde{K}(S^{2n})$, as follows.

Lemma 1.1. $\rho^t(0, 2q) = t^q$, and if $p > 0$ then

$$\rho^t(2p, 2q) = \begin{cases} \frac{1}{2} t^{p+q}(L+1) & (t \text{ even}) \\ t^{p+q} + \frac{1}{2} t^q(t^p-1)(L-1) & (t \text{ odd}). \end{cases}$$

As is easily seen, $\tilde{K}_G(\Sigma^{1,0})$ is isomorphic to the augmentation-ideal of $R(G)$. Identifying $\tilde{K}_G(\Sigma^{1,0})$ with $Z \cdot (1-L)$ it is clear that $\tilde{K}_G(\Sigma^{2p+1,2q}) = Z \cdot$

$(1-L)\lambda_{2p,2q}$. Hence we have the following

Corollary 1.2. ψ^t operates on $\tilde{K}_G(\Sigma^{2p+1,2q})$ as multiplication by 0 if t is even and by t^q if t is odd.

For $p, q-1 \geq 0$ suppose given a base point preserving G -map $f: \Sigma^{p+2k, 2q-1+2l} \rightarrow \Sigma^{2k, 2l}$ for k, l large, which is fixed in this section. f yields a cofiber sequence

$$\Sigma^{p+2k, 2q-1+2l} \xrightarrow{f} \Sigma^{2k, 2l} \xrightarrow{i} C_f \xrightarrow{j} \Sigma^{p+2k, 2q+2l} \xrightarrow{-\Sigma^{0,1}f} \Sigma^{2k, 2l+1}$$

where i, j are the inclusion and projection maps and C_f is the mapping cone of f . Applying \tilde{K}_G we obtain the following exact sequence.

$$\begin{aligned} 0 \leftarrow \tilde{K}_G(\Sigma^{2k, 2l}) \xleftarrow{i^*} \tilde{K}_G(C_f) \xleftarrow{j^*} \tilde{K}_G(\Sigma^{p+2k, 2q+2l}) \leftarrow 0 \\ \approx R(G) \qquad \qquad \approx \begin{cases} R(G) & (p \text{ even}) \\ Z & (p \text{ odd}) \end{cases} \end{aligned}$$

Choose generators ξ, η of $\tilde{K}_G(C_f)$ so that

$$i^*(\xi) = \lambda_{2k, 2l} \text{ and } \eta = \begin{cases} j^*(\lambda_{p+2k, 2q+2l}) & (p \text{ even}) \\ j^*((1-L)\lambda_{p-1+2k, 2q+2l}) & (p \text{ odd}). \end{cases}$$

For any odd integer $t (\neq \pm 1)$, $\psi^t(\xi)$ must be given by the formula

$$\psi^t(\xi) = \rho^t(2k, 2l)\xi + \begin{cases} (c(t)+d(t)(L-1))\eta & (p \text{ even}) \\ c(t)\eta & (p \text{ odd}), \end{cases}$$

$c(t), d(t) \in Z$. So we set

$$\begin{aligned} \lambda(f) &= \frac{c(t)}{t^{p/2+k+q+l} - t^{k+l}} & (p \text{ even}) \\ \mu(f) &= \begin{cases} \frac{1}{2} \left(\frac{c(t)}{t^{p/2+k+q+l} - t^{k+l}} + \frac{2d(t)-c(t)}{t^{q+l} - t^l} \right) & (p \text{ even}) \\ \frac{c(t)}{t^{q+l} - t^l} & (p \text{ odd}). \end{cases} \end{aligned}$$

Using Lemma 1.1, Corollary 1.2 and the relation $\psi^s \psi^t = \psi^{st}$ we can check that the values $\{\lambda(f)\}, \{\mu(f)\}$ do not depend on the choice of an integer t where $\{ \}$ denotes the coset in Q/Z . As in [1, IV], §7 we see that the assignment

$$f \mapsto \begin{cases} (\{\lambda(f)\}, \{\mu(f)\}) & (p \text{ even}) \\ \{\mu(f)\} & (p \text{ odd}) \end{cases}$$

induces a group homomorphism

$$e_G: \pi_{p, 2p-1}^S \rightarrow \begin{cases} Q/Z \oplus Q/Z & (p \text{ even}) \\ Q/Z & (p \text{ odd}) \end{cases} \text{ for } p, q-1 \geq 0.$$

Regard e_G as taking values in $\tilde{K}_G(\Sigma^{p+2k, 2q+2l}) \otimes Q/Z$, namely let $e_G[f]$ be $(\{\lambda(f)\} + \{\mu(f)\}(L-1))\lambda_{p+2k, 2q+2l}$ or $\{\mu(f)\}(1-L)\lambda_{p-1+2k, 2l}$ according as p is even or odd where $[f]$ is the stable homotopy class of f . Then we have easily the following

Proposition 1.3. e_G is natural for stable maps from $\Sigma^{p, 2q-1}$ to $\Sigma^{r, 2q-1}$.

To evaluate $\psi^t(\xi)$ we shall next describe e_G in terms of the equivariant Chern character. Let ch_G be as in [18] and ch_G^n denote the $2n$ -dimensional component of ch_G which is a homomorphism of K_G to $H_G^{2n}(_, R_G)$ in the notation of [18]. By the definition of equivariant Bredon cohomology [7] we have the following canonical isomorphisms

$$\begin{aligned} H_G^{p+2k+2q+2l}(C_f, R_G) &\approx H^{p+2k+2q+2l}(C_{\psi f}, Q) \\ &\approx H^{p+2k+2q+2l}(S^{p+2k+2q+2l}, Q), \\ H_G^{2q+2l}(C_f, R_G) &\approx H^{2q+2l}(C_{\phi f}, Q) \cdot (1-L) \\ &\approx H^{2q+2l}(S^{2q+2l}, Q) \cdot (1-L). \end{aligned}$$

Here ψ and ϕ are the forgetful and fixed point functors [3]. Under the identification of the above isomorphisms we may set

$$ch_G^{p/2+k+q+l}(\xi) = a(f)h^{p+2k+2q+2l}$$

and

$$ch_G^{q+l}(\xi) = b(f)h^{2q+2l}(1-L),$$

$a(f), b(f) \in Q$ (p even) where $h^{2i} \in H^{2i}(S^{2i}, Z)$ is a canonical generator such that $ch^i(\psi^* \lambda_{0, 2i}) = h^{2i}$. Then we obtain

Proposition 1.4. If p even then

$$\lambda(f) = a(f), \mu(f) = \frac{1}{2} \left(a(f) - \frac{b(f)}{2^{p/2+k-1}} \right)$$

and if p is odd then

$$\mu(f) = \frac{b(f)}{2^{(p-1)/2+k}}.$$

Proof. Consider the following commutative diagram with the exact sequence which ϕf yields as f does.

$$(*) \quad \begin{array}{ccccccc} 0 & \leftarrow & \tilde{K}_G(\Sigma^{2k, 2l}) & \xleftarrow{i^*} & \tilde{K}_G(C_f) & \xleftarrow{j^*} & \tilde{K}_G(\Sigma^{p+2k, 2q+2l}) \leftarrow 0 \\ & & \downarrow h^* & & \downarrow h^* & & \downarrow h^* \\ 0 & \leftarrow & \tilde{K}_G(\Sigma^{0, 2l}) & \xleftarrow{i_1^*} & \tilde{K}_G(C_{\phi f}) & \xleftarrow{j_1^*} & \tilde{K}_G(\Sigma^{0, 2q+2l}) \leftarrow 0 \end{array}$$

(Here h 's are the inclusions.) Choose $\xi_1 \in \tilde{K}_G(C_{\phi_f})$ so that $i_1^*(\xi_1) = \lambda_{0,2l}$ and put $\eta_1 = j_1^*(\lambda_{0,2q+2l})$. Then we may write

$$h^*(\xi) = 2^{k-1}(1-L)\xi_1 + x(1-L)\eta_1, \quad x \in Z$$

for a cohomological reason and the fact that $h^*(\lambda_{2k,2l}) = 2^{k-1}(1-L)\lambda_{0,2l}$. Applying ψ^t we have

$$(1) \quad \psi^t(h^*\xi) = 2^{k-1}(1-L)\psi^t(\xi_1) + xt^{q+l}(1-L)\eta_1.$$

On the other hand, apply h^* to the defining formula of $c(t)$, $d(t)$ we have

$$(2) \quad \begin{aligned} \psi^t(h^*\xi) &= 2^{k-1}t^l(1-L)\xi_1 + xt^l(1-L)\eta_1 \\ &+ \begin{cases} 2^{p/2+k-1}(c(t)-2d(t))(1-L)\eta_1 & (p \text{ even}) \\ 2^{(p-1)/2+k}c(t)(1-L)\eta_1 & (p \text{ odd}). \end{cases} \end{aligned}$$

Combining (1) and (2) shows

$$\psi^t(\xi_1) = t^l\xi_1 + \frac{x(t^l - t^{q+l})}{2^{k-1}}\eta_1 + \begin{cases} 2^{p/2}(c(t)-2d(t))\eta_1 & (p \text{ even}) \\ 2^{(p+1)/2}c(t)\eta_1 & (p \text{ odd}). \end{cases}$$

Case p even. From the definition of ch_G it follows easily that

$$ch_G^{p/2+k+q+l}(\xi) = ch^{p/2+k+q+l}(\psi\xi)$$

and

$$ch_G^{q+l}(\xi) = 2^{k-1}ch^{q+l}(\psi\xi_1)(1-L) + xh^{2q+2l}(1-L).$$

Hence we get

$$ch^{p/2+k+q+l}(\psi\xi) = a(f)h^{p+2k+2q+2l} \quad \text{and} \quad ch^{q+l}(\psi\xi_1) = \frac{b(f)-x}{2^{k-1}}h^{2q+2l}.$$

Therefore [1, IV], Proposition 7.5 for ψf and ϕf leads to the equalities

$$a(f) = \frac{c(f)}{t^{p/2+k+q+l} - t^{k+l}} \quad \text{and} \quad \frac{b(f)}{2^{p/2+k-1}} = \frac{c(t)-2d(t)}{t^{q+l} - t^l}.$$

Case p odd. Similar to the proof of the above case.

q.e.d.

2. $(0, 2q-1)$ -stem

Let $\pi: \Sigma^{2k, 2q-1+2l} \rightarrow \Sigma^{2k, 2q-1+2l} / \Sigma^{0, 2q-1+2l}$ be the canonical projection map for k, l large. Let $\lambda_{p,q}^S$ denote the equivariant stable homotopy group introduced in [12]. Then we have by [12] a split short exact sequence

$$0 \rightarrow \lambda_{0, 2q-1}^S \xrightarrow{\pi^*} \pi_{0, 2q-1}^S \xrightleftharpoons[\theta]{\phi} \pi_{2q-1}^S \rightarrow 0$$

where π^* is the homomorphism induced by π and θ denotes a left inverse of ϕ as in [4], §5.

By the definition we can easily describe the values of e_G on $\text{Im } \theta$ in terms of the complex e -invariant e_C in [1, IV]. So we consider e_G on $\text{Im } \pi^*$ in this section.

Suppose given a base point preserving G -map $\tilde{f}: \Sigma^{2k, 2q-1+2l}/\Sigma^{0, 2q-1+2l} \rightarrow \Sigma^{2k, 2l}$, so that \tilde{f} and $\tilde{f}\pi$ define elements $[\tilde{f}]$ and $[\tilde{f}\pi]$ of $\lambda_{0, 2q-1}^S$ and $\pi_{0, 2q-1}^S$ respectively. We consider $\tilde{f}\pi$ as f in §1.

Since $\Sigma^{i, j}/\Sigma^{0, j}$ is equivariantly homeomorphic to $\Sigma^{0, j+1}S_+^{i, 0}$ ([12], Lemma 4.1), we have $\tilde{K}_G(\Sigma^{i, j}/\Sigma^{0, j}) \approx K^{-j-1}(RP^{i-1})$ [16] where RP^n is the real n -dimensional projective space. Let η_n be the complexification of a canonical real line bundle over RP^n and put $\tilde{\eta}_n = 1 - \eta_n$. We now recall [6] that

$$\begin{aligned}\tilde{K}^0(RP^{2n}) &= Z/2^n \cdot \tilde{\eta}_{2n}, \quad K^1(RP^{2n}) = 0 \\ \tilde{K}^0(RP^{2n+1}) &= Z/2^n \cdot \tilde{\eta}_{2n+1}, \quad K^1(RP^{2n+1}) \approx Z.\end{aligned}$$

Then we can identify

$$\tilde{K}_G^0(\Sigma^{2k, 2q-1+2l}/\Sigma^{0, 2q-1+2l}) = Z \oplus Z/2^{k-1} \cdot (\psi \lambda_{0, 2q+2l}) \tilde{\eta}_{2k-1}.$$

Consider $\tilde{f}^*: \tilde{K}_G(\Sigma^{2k, 2l}) \rightarrow \tilde{K}_G(\Sigma^{2k, 2q-1+2l}/\Sigma^{0, 2q-1+2l})$. Because $[\tilde{f}] \in \lambda_{0, 2q-1}^S$ for $q \geq 1$ is of finite order ([12], Theorem 2.4 and Corollary 6.3) we may put

$$\tilde{f}^*(\lambda_{2k, 2l}) = [\tilde{b}(\tilde{f})] (\psi \lambda_{0, 2q+2l}) \tilde{\eta}_{2k-1}, \quad \tilde{b}(\tilde{f}) \in Z$$

where $[\]$ denotes the coset in $Z/2^{k-1}$.

Lemma 2.1. $\tilde{b}(\tilde{f}) = -b(\tilde{f}\pi) \bmod 2^{k-1}$
where $b(\tilde{f}\pi)$ is as in §1.

Proof. Observe the following commutative diagram involving (*) in §1.

$$\begin{array}{ccccccc} \tilde{K}_G(\Sigma^{2k, 2q-1+2l}/\Sigma^{0, 2q-1+2l}) & \xleftarrow{\tilde{f}^*} & \tilde{K}_G(\Sigma^{2k, 2l}) & \xleftarrow{\quad} & \tilde{K}_G(C_{\tilde{f}}) & \xleftarrow{\quad} & \tilde{K}_G(\Sigma^{2k, 2q+2l}/\Sigma^{0, 2q+2l}) \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \leftarrow & \tilde{K}_G(\Sigma^{2k, 2l}) & \xleftarrow{i^*} & \tilde{K}_G(C_{\tilde{f}\pi}) & \xleftarrow{j^*} & \tilde{K}_G(\Sigma^{2k, 2q+2l}) & \leftarrow 0 \\ & & \downarrow h^* & & \downarrow h^* & & \downarrow h^* & \\ 0 & \leftarrow & \tilde{K}_G(\Sigma^{0, 2l}) & \xleftarrow{i_1^*} & \tilde{K}_G(C_{\phi(\tilde{f}\pi)}) & \xleftarrow{j_1^*} & \tilde{K}_G(\Sigma^{0, 2q+2l}) & \leftarrow 0 \\ & & & & & & \downarrow \delta & \\ & & & & & & K_G^1(\Sigma^{2k, 2q+2l}/\Sigma^{0, 2q+2l}) & \end{array}$$

where the right-hand sequence is the exact sequence for a pair $(\Sigma^{2k, 2q+2l}, \Sigma^{0, 2q+2l})$. Clearly $C_{\phi(\tilde{f}\pi)} \approx \Sigma^{0, 2q+2l} \vee \Sigma^{0, 2l}$, hence we can verify that $\tilde{f}^*(\lambda_{2k, 2l}) = -\delta j_1^{*-1} h^*(\xi)$ where ξ is as in §1. Hence the canonical identification such that $\tilde{K}_G(\Sigma^{0, 2q+2l}) = \tilde{K}(S^{2q+2l}) \otimes R(G) = H^{2q+2l}(S^{2q+2l}, Z) \otimes R(G)$ leads to the desired assertion. q.e.d.

Let BG denote the real infinite dimensional projective space. There is an integer $c(n)$ such that $c(n)\eta_{2n-1}$ becomes trivial (see, e.g. [9], p. 219). So we have an equivariant homeomorphism $\Sigma^{c(n),0}S_+^{n,0} \approx \Sigma^{0,c(n)}S_+^{n,0}$. This homeomorphism, the equivariant suspension theorem and the Spanier-Whitehead duality theorem yield an isomorphism

$$\lambda_{0,n}^S \xrightarrow[\approx]{} \pi_n^S(BG_+),$$

denoted by I , as follows. Let τ be the tangent bundle of RP^{2k-1} and ν be a normal bundle of RP^{2k-1} for an embedding of RP^{2k-1} in R^{2m-1} for m suitably large. Note that the Thom complex $T(\nu)$ of ν is a $(2m-1)$ -dual of RP_+^{2k-1} [5], and $\tau \oplus 1 \approx 2k\eta'_{2k-1}$ so that $S^{2m}T((sc-k)\eta_{2k-1}) \approx S^{2sc}T(\nu)$ for $sc > k$ where η'_{2k-1} denotes the underlying real vector bundle of η_{2k-1} and $c=c(k)$ is as above. Then we have the following isomorphisms.

$$\begin{aligned} \lambda_{0,n}^S &= \lim_{k,l} [\Sigma^{2k,n+2l}/\Sigma^{0,n+2l}, \Sigma^{2k,2l}]^G && \text{by definition [12]} \\ &\approx \lim_{k,l} [\Sigma^{0,n+2l+1}S_+^{2k,0}, \Sigma^{2k,2l}]^G \\ &\approx \lim_{k,l} [\Sigma^{2sc,n+2l-2sc+1}S_+^{2k,0}, \Sigma^{2k,2l}]^G && \text{for some } c \\ &\approx \lim_{k,l} [\Sigma^{2sc-2k,n+2l-2sc+1}S_+^{2k,0}, \Sigma^{0,2l}]^G && \text{by [3], Theo. 11.9} \\ &\approx \lim_{k,l} [S^{n+2l-2sc+1}T((sc-k)\eta_{2k-1}), S^{2l}] \\ &\approx \lim_{k,l} [S^{n+2l-2m+1}T(\nu), S^{2l}] \\ &\approx \lim_k \{S^n, RP_+^{2k-1}\} && \text{by [19], Cor. (7.10)} \\ &= \pi_n^S(BG_+) \end{aligned}$$

On the other hand, the geometrical interpretation of I by Landweber [12] shows that the composite $\psi\pi^*I^{-1}: \pi_n^S(BG_+) \rightarrow \pi_n^S$ agrees with the $Z/2$ -transfer. So we write $t=\psi\pi^*I^{-1}$ as usual.

Following the homotopical construction of I we see that $I[\tilde{f}]$ is represented by a stable map $g: S^{2q-1} \rightarrow RP_+^{2k-1}$. Let $\tilde{g}: S^{2q-1} \rightarrow RP^{2k-1}$ be the composite g and the canonical projection from RP_+^{2k-1} to RP^{2k-1} and let

$$\alpha_1 \in \pi_{2q-1}^S(BG)$$

denote the stable homotopy class induced by \tilde{g} . Then we have

Proposition 2.2. $\left\{ \frac{\tilde{b}(\tilde{f})}{2^{k-1}} \right\} = e_c t(\alpha_1)$

where e_c is as in [1, IV].

We prepare a lemma for a proof of Proposition 2.2. We recall the following universal coefficient sequence for a finite CW -complex X [2]

$$0 \rightarrow \text{Ext}(\tilde{K}^0(X), \mathbb{Z}) \rightarrow K_1(X) \xrightarrow{k} \text{Hom}(K^1(X), \mathbb{Z}) \rightarrow 0$$

where k is a map induced by the Kronecker product. Here we denote by ι the injection map. Furthermore we have a natural homomorphism

$$\text{Hom}(\tilde{K}^0(X), \mathbb{Q}/\mathbb{Z}) \rightarrow \text{Ext}(\tilde{K}^0(X), \mathbb{Z}),$$

which we denote by Δ . In particular, for $X=RP^{2k}$, ι and Δ are isomorphisms.

Denote by p the collapsing map $RP^{2k-1} \rightarrow RP^{2k-1}/RP^{2k-2}$ and identify RP^{2k-1}/RP^{2k-2} with S^{2k-1} . Then, clearly $p^*: \tilde{K}^0(S^{2k}) = K^1(S^{2k-1}) \rightarrow K^1(RP^{2k-1})$ is an isomorphism and hence by using the universal coefficient sequence we see that $p_*: K_1(RP^{2k-1}) \rightarrow K_1(S^{2k-1}) = \tilde{K}_0(S^{2k})$ is an epimorphism. Therefore, if we put $z' = p^*(\psi\lambda_{0,2k}) \in K^1(RP^{2k-1})$ then we have an element $z \in K_1(RP^{2k-1})$ such that p_*z is a dual element of $\psi\lambda_{0,2k}$, i.e. $\langle z', z \rangle = 1$, which is a fundamental class of RP^{2k-1} ([19], p. 217). By [19], Corollary (7.8) we have an isomorphism

$$P = z \cap : \tilde{K}^0(RP^{2k-1}) \rightarrow K_1(RP^{2k-1}).$$

Consider the composite

$$\tilde{K}^0(RP^{2k-1}) \xrightarrow{P} K_1(RP^{2k-1}) \xrightarrow{i'_*} K_1(RP^{2k}) \xrightarrow{(\iota\Delta)^{-1}} \text{Hom}(\tilde{K}^0(RP^{2k}), \mathbb{Q}/\mathbb{Z})$$

where $i': RP^{2k-1} \hookrightarrow RP^{2k}$ is the inclusion map. Then

$$\textbf{Lemma 2.3.} \quad ((\iota\Delta)^{-1}i'_*P\tilde{\eta}_{2k-1})\tilde{\eta}_{2k} = -\left\{\frac{1}{2^{k-1}}\right\}.$$

Proof. Let γ^* be the co-Hopf bundle on the complex $(k-1)$ -dimensional projective CP^{k-1} and γ be its dual. By D and S we denote the total spaces of the unit disk and unit sphere bundles of $\gamma^* \otimes \gamma^*$ with respect to some metric. Then $D \simeq CP^{k-1}$ clearly and $S \approx RP^{2k-1}$ (see [10], IV.1.14. Example). We identify S with RP^{2k-1} . Because, if we put $\tilde{\gamma} = 1 - \gamma$ then $K^*(D) \approx Z[\tilde{\gamma}]/(\tilde{\gamma}^k)$ and $i^*\tilde{\gamma} = \tilde{\eta}_{2k-1}$, we have a short exact sequence

$$0 \rightarrow K^1(S) \xrightarrow{\delta} K^0(D, S) \xrightarrow{j^*} K^0(D) \xrightarrow{i^*} K^0(S) \rightarrow 0$$

where δ is a coboundary homomorphism and i, j are the inclusion maps. As is well known, $j^*\lambda = -\tilde{\gamma}^{*2} + 2\tilde{\gamma}^*$ where $\tilde{\gamma}^* = 1 - \gamma^*$ and λ is the Thom class of $\gamma^* \otimes \gamma^*$. Hence $K^*(D, S) \approx \bigoplus_{i=0}^{k-1} \mathbb{Z} \cdot \lambda \tilde{\gamma}^i$. Moreover, by an observation for $\tilde{\gamma}^{k-1}$ in [6], p. 100 we have

$$\delta^{-1}\lambda\tilde{\gamma}^{k-1} = z'.$$

Put $z'_1 = \delta z'$ and denote by z_1 a dual element of z'_1 so that we may suppose that $\partial z_1 = z$ where ∂ is the boundary homomorphism. Similarly $P_1 = z_1 \cap : K^0(D) \rightarrow K_0(D, S)$ is then an isomorphism and the diagram

$$\begin{array}{ccc} K^0(D) & \xrightarrow{i^*} & K^0(S) \\ P_1 \downarrow & & P \downarrow \\ K_0(D, S) & \xrightarrow{\partial} & K_1(S) \end{array}$$

commutes.

A routine computation shows that $\lambda\tilde{\gamma}^{k-2} \in K^0(D, S)$ is a dual element of $P_1\tilde{\gamma}$, i.e.,

$$\langle \lambda\tilde{\gamma}^{k-2}, P_1\tilde{\gamma} \rangle = 1.$$

Let put $M = D \times S^{2k-1}$ and $i_1: S \subset M$ be an embedding given by $i_1(x) = (i(x), p(x))$ $x \in S$. Then we get a short exact sequence

$$0 \rightarrow K^*(M, S) \xrightarrow{j_1^*} \tilde{K}^*(M) \xrightarrow{i_1^*} \tilde{K}^*(S) \rightarrow 0,$$

which is a free resolution of $\tilde{K}^*(S)$, where j_1 is the inclusion map. Hence we see that

$$\tilde{K}^0(M) = \bigoplus_{i=1}^{k-1} Z \cdot q^* \tilde{\gamma}^i \quad \text{and} \quad K^0(M, S) = \bigoplus_{i=0}^{k-2} Z \cdot q^* \lambda \tilde{\gamma}^i,$$

where q is the projection map of M to D .

Here we adopt the above resolution as a free resolution in the proof of [2], Theorem 3.1 for $K_1(S)$. Define $f \in \text{Hom}(K^0(M, S), Z)$ by

$$f(q^* \lambda \tilde{\gamma}^i) = \begin{cases} 1 & \text{if } i = k-2 \\ 0 & \text{otherwise.} \end{cases}$$

Then

$$\text{Hom}(q^*, 1)f = \langle \quad, P_1\tilde{\gamma} \rangle.$$

This implies that because $\text{Coker Hom}(j_1^*, 1) = \text{Ext}(\tilde{K}^0(S), Z)$,

$$\iota[f] = P\tilde{\eta}_{2k-1},$$

where $[f]$ denotes the equivalence class of f in $\text{Coker Hom}(j_1^*, 1)$.

By the definition of Δ it is verified that

$$(\Delta^{-1}[f])\tilde{\eta}_{2k-1} = -\left\{ \frac{1}{2^{k-1}} \right\}.$$

Hence,

$$(\iota\Delta)^{-1}(P\tilde{\eta}_{2k-1})\tilde{\eta}_{2k-1} = -\left\{\frac{1}{2^{k-1}}\right\}.$$

This proves the lemma because $i'_*\iota\Delta = \iota\Delta\text{Hom}(i'^*, 1)$.

Proof of Proposition 2.2. We may suppose that ν is a complex vector bundle, since the stable tangent bundle of RP^{2k-1} has a complex structure.

Observing the construction of I we have the following commutative diagram.

$$\begin{array}{ccc} \tilde{K}_G^0(\Sigma^{0,2q+2l}S_+^{2k,0}) & \xleftarrow{\tilde{f}^*} & \tilde{K}_G^0(\Sigma^{2k,2l}) \\ I_0 \downarrow & & \uparrow I_1 \\ \tilde{K}^0(S^{2l+2q-2m}T(\nu)) & & \tilde{K}_G^0(\Sigma^{0,2l}) = \tilde{K}^0(S^{2l}) \otimes R(G) \\ D_2 \downarrow & & \uparrow D_3 \\ K_1(RP^{2k}) \xleftarrow{i'_*} K_1(RP^{2k-1}) & & \xleftarrow{\tilde{g}^*} K_1(S^{2q-1}) \end{array}$$

Here D_2, D_3 are the duality isomorphisms as in [19], Corollary (7.10), and I_0, I_1 are isomorphisms given by $I_0((\psi\lambda_{0,2q+2l})\tilde{\eta}_{2k-1}) = (\psi\lambda_{0,2l+2q-2m})\lambda_\nu\tilde{\eta}_{2k-1}$, $I_1(\lambda_{0,2l}) = \lambda_{2k,2l}$ where λ_ν denotes the Thom class of ν .

By [19], Corollaries (7.8) and (7.10) we have

$$D_2I_0((\psi\lambda_{0,2q+2l})\tilde{\eta}_{2k-1}) = P\tilde{\eta}_{2k-1},$$

which is pointed out by Dyer in [8]. By Lemma 2.3 we therefore have

$$((\iota\Delta)^{-1}(i'\tilde{g})_*\beta)\tilde{\eta}_{2k} = -\left\{\frac{\tilde{b}(\tilde{f})}{2^{k-1}}\right\}$$

where $\beta = D_3(\psi\lambda_{0,2l})$.

Identifying $K_1(RP^{2k})$ with $\text{Hom}(\tilde{K}^0(RP^{2k}), \mathbb{Q}/\mathbb{Z})$ through the isomorphism $\iota\Delta$, we may write

$$(h\alpha_1)\tilde{\eta}_{2k} = -\left\{\frac{\tilde{b}(\tilde{f})}{2^{k-1}}\right\}$$

in terms of the Hurewicz homomorphism $h: \pi_{2q-1}^S(BG) \rightarrow K_1(BG)$. Hence by [11], Theorem 2.1 we obtain

$$(CH^q(\alpha_1))\tilde{\eta}_{2k} = -\left\{\frac{\tilde{b}(\tilde{f})}{2^{k-1}}\right\}$$

where CH^q is the functional Chern character. By the naturality of CH^q we get

$$e_c t(\alpha_1) = -(CH^i(\alpha_1))\tilde{\eta}_{2k}.$$

(For the sign, see Remark 4 of [11], p. 128.)

Therefore

$$\left\{ \frac{\tilde{b}(\tilde{f})}{2^{k-1}} \right\} = e_c t(\alpha_1).$$

q.e.d.

Consequently we get the following

Theorem 2.4. For $\alpha \in \pi_{0,2q-1}^S$ ($q \geq 1$),

$$e_c(\alpha) = \begin{cases} (e_c(\psi\alpha), 0) & \text{for } \alpha \in \text{Im } \theta \\ (e_c(\psi\alpha), \frac{1}{2}(e_c(\psi\alpha) + e_c t(\alpha_1) + \varepsilon)) & \text{for } \alpha \in \text{Im } \pi^* \end{cases}$$

($\varepsilon=0, 1$) where α_1 denotes the first factor of $I\pi^{*-1}(\alpha)$ under the identification $\pi_{2q-1}^S(BG_+) = \pi_{2q-1}^S(BG) \oplus \pi_{2q-1}^S$.

Proof. As to the first factors this is clear from the definitions of e_c and e_c . As to the second this follows in addition from Proposition 1.4, Lemma 2.1 and Proposition 2.2. q.e.d.

3. Images of the S^1 -transfer

Let $\tilde{t}: \pi_n^S(BS_+^1) \rightarrow \pi_{n+1}^S(BG_+)$ denote the S^1 -transfer, where BS^1 is the complex infinite dimensional projective space.

Proposition 3.1. Let $\alpha \in \text{Im } \{\pi^*: \lambda_{0,4q-1}^S \rightarrow \pi_{0,4q-1}^S\}$ ($q \geq 1$) and $I\pi^{*-1}(\alpha) \in \text{Im } \tilde{t}$. Then

$$e_c t(\alpha_1) = (1 - 2^{2q})e_c(\psi\alpha)$$

where α_1 is as in Theorem 2.4.

Proof. Consider the isomorphisms

$$\lambda_{0,4q-1} \xrightarrow{I} \pi_{4q-1}^S(BG_+) = \pi_{4q-1}^S(BG) \oplus \pi_{4q-1}^S.$$

We may write $I\pi^{*-1}(\alpha) = (\alpha_1, \alpha_2)$. Applying t we have

$$\psi\alpha = t\alpha_1 + 2\alpha_2,$$

Since $t = \psi\pi^*I^{-1}$ and t operates on π_{4q-1}^S as multiplication by 2. From [13], Theorem 3.4 it follows that

$$e_c(\alpha_2) = 2^{2q-1}e_c(\psi\alpha).$$

Therefore we get the proposition.

The following theorem follows immediately from Theorem 2.4 and Proposition 3.1.

Theorem 3.2. *For $\alpha \in \pi_{0,4q-1}^S$ as in Proposition 3.1 we have*

$$e_G(\alpha) = (e_C(\psi\alpha), (1-2^{2q-1})e_C(\psi\alpha) + \frac{\varepsilon}{2}), \quad (\varepsilon = 0, 1).$$

Let $J_G: \widetilde{KO}_G^{-1}(\Sigma^{0,4q-1}) \rightarrow \pi_{0,4q-1}^S$ ($q \geq 1$) be the equivariant J -homomorphism [14, 17]. Set $\alpha = J_G(H\nu) \in \pi_{0,4q-1}^S$ where ν is a canonical generator of $\widetilde{KO}^{-1}(S^{4q-1})$ and $H = R^{1,0}$. Then $\alpha \in \text{Im } \pi^*$ because $\phi(\alpha) = 0$.

Lemma 3.3. *Let α be as above. Then $I\pi^{*-1}(\alpha)$ or $2I\pi^{*-1}(\alpha) \in \text{Im } \tilde{t}$ according as q is odd or even.*

Proof. We consider the S^1 -homotopy theory. Replace $R^{1,0}$ by the standard complex 1-dimensional non trivial representation V of S^1 in the $Z/2$ -homotopy theory. Then by the same argument as in [12] we have the S^1 -homotopy groups $\pi_n^{V,S}$, $\lambda_n^{V,S}$ and an exact sequence $\lambda_n^{V,S} \xrightarrow{\pi^*} \pi_n^{V,S} \xrightarrow{\phi} \pi_n^S$. Moreover, we have an isomorphism $\lambda_n^{V,S} \approx \pi_{n-1}^S(BS_+^1)$. Clearly the diagram

$$\begin{array}{ccccc} \lambda_n^{V,S} & \xrightarrow{\pi^*} & \pi_n^{V,S} & \xrightarrow{\phi} & \pi_n^S \\ r \downarrow & & r \downarrow & & \parallel \\ 0 \rightarrow \lambda_{0,n}^S & \xrightarrow{\pi^*} & \pi_{0,n}^S & \xrightarrow{\phi} & \pi_n^S \end{array}$$

commutes where r denotes the restriction of S^1 -actions. Identifying the left-hand groups with the cobordism groups canonically, r agrees with the S^1 -transfer \tilde{t} .

Analogously for S^1 -actions we can define the equivariant J -map J_V as follows. Denote by $U(kV+l)$ the unitary group of $kV \oplus C^l$ with the induced action and by U_V the infinite unitary group obtained by taking a limit with respect to canonical inclusions of $U(kV+l)$'s. Then we have a map J_V from the equivariant homotopy group $[S^n, U_V]^{S^1}$ to $\pi_n^{V,S}$ as usual.

Now a generator μ of $\tilde{K}^{-1}(S^{4q-1})$, viewed as a map from S^{4q-1} to a unitary group, comes from $[S^{4q-1}, U_V]^{S^1}$ and so $V\mu$ does. Generally an equivariant map from S^{4q-1} to U_V defines an element of $\tilde{K}_{S^1}^{-1}(S^{4q-1})$. So we have a map $[S^{4q-1}, U_V]^{S^1} \rightarrow \tilde{K}_{S^1}^{-1}(S^{4q-1})$.

Because $J_V(V\mu) = 0$, using the same notation for $V\mu$ in $[S^{4q-1}, U_V]^{S^1}$, there exists $x \in \lambda_{4q-1}^{V,S}$ such that $\pi^*x = J_V(V\mu)$. From the above discussion it follows that $r(J_V(V\mu)) = \alpha$ or 2α , so that $r(x) = \pi^{*-1}(\alpha)$ or $2\pi^{*-1}(\alpha)$, according as q is odd or even. q.e.d.

Let J_0 be the real J -homomorphism. By [1, IV], Theorem 7.16 we may write

$$e'_R J_o(\nu) = \frac{a_q}{m(2q)} \in Q/Z, \quad (a_q, m(2q)) = 1$$

where $m(2q)$, e'_R are as in [1, II]. Then we have

Theorem 3.4. For $\alpha = J_G(H\nu) \in \pi_{0,4q-1}^S$ ($q \geq 1$),

$$e_G(\alpha) = \begin{cases} \left(\frac{2a_q}{m(2q)}, 2(1-2^{2q-1})\frac{a_q}{m(2q)} + \frac{\varepsilon}{2} \right) & (q \text{ odd}) \\ \left(\frac{a_q}{m(2q)}, (1-2^{2q-1})\frac{a_q}{m(2q)} + \frac{\varepsilon}{4} + \frac{\varepsilon'}{2} \right) & (q \text{ even}) \end{cases}$$

($\varepsilon, \varepsilon' = 0, 1$) as rational numbers mod 1 and the order of each factor of $e_G(\alpha)$ is $\frac{m(2q)}{2}$ or $m(2q)$ according as q is odd or even.

Proof. The first claim follows from Theorem 3.2, Lemma 3.3 and [1, IV], Proposition 7.14. The second follows from [1, II], Lemma (2.12) and the equality $\nu_2(m(2q)) = 3 + \nu_2(q)$ ([1, II], p. 139) immediately. e.d.q.

4. Real $Z/2$ - e -invariants

We take a base point preserving G -map $f: \Sigma^{p+8k, 2q-1+8l} \rightarrow \Sigma^{8k, 8l}$ as a representative of elements of $\pi_{p, 2q-1}^S$ for $p, q-1 \geq 0$. Then the parallel argument to e_G , using the Adams operation in the KO_G -theory [12] and Table of [14], yields the following equivariant e -invariants.

$$(1) \quad e_{G,R}: \pi_{8p+4\zeta+i, 8q+4\delta-1}^S \rightarrow \begin{cases} (Q/Z)^2 & (i=0) \\ Q/Z & (i=1, 2, 3) \end{cases}$$

$$(2) \quad e_{G,R}: \pi_{8p+4\zeta+2, 8q+4\delta+1}^S \rightarrow Q/Z$$

for $\zeta, \delta = 0, 1$.

Theorem 4.1. For $\bar{\alpha} = J_G(\nu)$, $\alpha = J_G(H\nu) \in \pi_{0,4q-1}^S$ ($q \geq 1$),

$$e_{G,R}(\bar{\alpha}) = \left(\frac{a_q}{m(2q)}, 0 \right),$$

$$e_{G,R}(\alpha) = \left(\frac{a_q}{m(2q)}, (1-2^{2q-1})\frac{a_q}{m(2q)} + \frac{\varepsilon}{4} + \frac{\varepsilon'}{2} \right)$$

($\varepsilon, \varepsilon' = 0, 1$) as rational numbers mod 1 and the order of the second factor of $e_{G,R}(\alpha)$ is $m(2q)$.

Proof. As to the first factors of the equalities this follows immediately from the definitions of $e_{G,R}$ and e'_R . As to the second this follows in addition from Theorem 3.4 and the fact that $e_G = e_{G,R}$ or $2e_{G,R}$ according as q is even or odd. The proof of the last claim is similar to that of Theorem 3.4. q.e.d.

Finally we shall consider $e_{G,R}$ on $\text{Im } J_G$ for $\pi_{p,4q-1}^S$ ($p \geq 1$). Let χ, ρ be as in [3] and $\hat{\eta}$ be the homomorphism induced by the element of [4], (8.1). Observe χ, ρ and $\hat{\eta}$ on the groups $\widetilde{KO}_G^{-1}(\Sigma^{p,2q-1})$ (see [15], §2), then since $e_{G,R} J_G$ commutes with χ, ρ and $\hat{\eta}$ (by an analogue of Proposition 1.3), we can compute $e_{G,R}$ of (1) on $\text{Im } J_G$ inductively by using Theorem 4.1. For $e_{G,R}$ of (2), considering $\psi e_{G,R}$ we get readily $e_{G,R}$ on $\text{Im } J_G$. Specifically we have

Theorem 4.2. Let $v_1 \in \widetilde{KO}_G^{-1}(\Sigma^{8p+4\zeta, 8q+4\delta-1})$ ($8p+4 > 0$), $v_2 \in \widetilde{KO}_G^{-1}(\Sigma^{8p+4\zeta+i, 8q+4\delta-1})$ ($1 \leq i \leq 3$) and $v_3 \in \widetilde{KO}_G^{-1}(\Sigma^{8p+4\zeta+2, 8q+4\delta+1})$ be generators as modules over the real representation ring of G respectively and set $\alpha_k = J_G(v_k)$ ($1 \leq k \leq 3$). Then as rational numbers mod 1

$$\begin{aligned} e_{G,R}(\alpha_1) &= \left(\frac{a_{2p+2q+\zeta+\delta}}{m(4p+4q+2\zeta+2\delta)}, \frac{1}{2} \left\{ \frac{a_{2p+2q+\zeta+\delta}}{m(4p+4q+2\zeta+2\delta)} \right. \right. \\ &\quad \left. \left. - (1-2^{4q+2\delta-1}) \frac{a_{2q+\delta}}{m(4q+2\delta)} - \frac{\varepsilon}{4} - \frac{\varepsilon'}{2} + \varepsilon'' \right\} \right), \\ e_{G,R}(\alpha_2) &= (1-2^{4q+2\delta-1}) \frac{a_{2q+\delta}}{m(4q+2\delta)} + \frac{\varepsilon}{4} + \frac{\varepsilon'}{2}, \\ e_{G,R}(\alpha_3) &= \frac{a_{2p+2q+\zeta+\delta+1}}{m(4p+4q+2\zeta+2\delta+2)} + \frac{\varepsilon}{2} \end{aligned}$$

($\varepsilon, \varepsilon', \varepsilon'' = 0, 1$) up to sign and

$$\begin{aligned} \text{order } e_{G,R}(\alpha_1) &= \frac{m(4p+4q+2\zeta+2\delta)m(4q+2\delta)}{2^\kappa d}, \\ \text{order } e_{G,R}(\alpha_2) &= m(4q+2\delta), \\ \text{order } e_{G,R}(\alpha_3) &= m(4p+4q+2\zeta+2\delta+2) \end{aligned}$$

where

$$d = \left(\frac{m(4p+4q+2\zeta+2\delta)}{2^{\nu_2(2p+2q+\zeta+\delta)+3}}, \frac{m(4q+2\delta)}{2^{\nu_2(2q+\delta)+3}} \right)$$

and κ is the following integer:

$$\begin{array}{ll} \nu_2(2q+\zeta)+2 & \text{if } \zeta = \delta \text{ and } \nu_2(2q+\zeta) \leq \nu_2(p+q+\zeta), \\ \nu_2(2q+\zeta)+3 & \text{if } \zeta = \delta \text{ and } \nu_2(2q+\zeta) = \nu_2(p+q+\zeta)+1, \\ \nu_2(p+q+\zeta)+3 & \text{if } \zeta = \delta \text{ and } \nu_2(2q+\zeta) \geq \nu_2(p+q+\zeta)+2, \\ 3 & \text{if } \zeta = 0 \text{ and } \delta = 1, \\ 2 & \text{if } \zeta = 1 \text{ and } \delta = 0. \end{array}$$

Here let $\nu_2(s)$ denote the exponent to which 2 occurs in s .

By Theorems 4.1, 4.2 and the results of [15] we have

Corollary 4.3. For $\pi_{p,q}^S$ in [15], Theorems 3.1, 3.2 and 3.3,

$$\operatorname{Im} J_G \xhookrightarrow{i} \pi_{p,q}^S \xrightarrow{e_{G,R}} \operatorname{Im} e_{G,R}$$

provides a direct sum splitting.

References

- [1] J.F. Adams: *On the groups $J(X)$* —II, IV, *Topology* **3** (1965), 137–171, **5** (1966), 21–71.
- [2] D.W. Anderson: *Universal coefficient theorems for K -theory*, Preprint.
- [3] S. Araki and M. Murayama: *τ -Cohomology theories*, *Japan. J. Math.* **4** (1978), 363–416.
- [4] ——— and K. Iriye: *Equivariant stable homotopy groups of spheres with involutions*, I, *Osaka J. Math.* **19** (1982), 1–55.
- [5] M.F. Atiyah: *Thom complexes*, *Proc. London Math. Soc.* (3) **11** (1961), 291–310.
- [6] ———: *K -theory*, W.A. Benjamin, Inc., 1967.
- [7] G.E. Bredon: *Equivariant cohomology theories*, *Lecture Notes in Math.* **34**, Springer-Verlag, 1967.
- [8] E. Dyer: *Relations between cohomology theories*, *Coll. on Algebraic Topology*, Aarhus 1962, 89–93.
- [9] D. Husemoller: *Fiber bundles*, McGraw-Hill, Inc., 1966.
- [10] M. Karoubi: *K -theory. An introduction*, Berlin-Heidelberg-New York, Springer, 1978.
- [11] K. Knapp: *Das bild des Hurewicz-homomorphismus $h: \pi_*^S(BZ_p) \rightarrow K_1(BZ_p)$* , *Math. Ann.* **223** (1976), 119–138.
- [12] P.S. Landweber: *On equivariant maps between spheres with involutions*, *Ann. of Math.* **89** (1969), 125–137.
- [13] P. Löffler and L. Smith: *Line bundles over framed manifolds*, *Math. Z.* **138** (1974), 35–52.
- [14] P. Löffler: *Equivariant framability of involutions on homotopy spheres*, *Manuscripta Math.* **23** (1978), 161–171.
- [15] H. Minami: *On equivariant J -homomorphism for involutions*, *Osaka J. Math.* **20** (1983), 109–122.
- [16] G.B. Segal: *Equivariant K -theory*, *Inst. Hautes Études Sci. Publ. Math.* **34** (1968), 129–151.
- [17] ———: *Equivariant stable homotopy theory*, *Actes Congrès intern. Math.*, 1970, t. **2**, 59–63.
- [18] J. Slomińska: *On the equivariant Chern homomorphism*, *Bull. Acad. Polon. Sci.* **XXIV** (1976), 909–913.
- [19] G.W. Whitehead: *Generalized homology theories*, *Trans. Amer. Math. Soc.* **102** (1962), 227–283.

Department of Mathematics
Osaka City University
Sugimoto, Sumiyoshi-ku
Osaka 558, Japan