

Global solutions to a chemotaxis system with non-diffusive memory

メタデータ	言語: English 出版者: OCAMI 公開日: 2019-10-07 キーワード (Ja): キーワード (En): chemotaxis system, global solutions 作成者: 杉山, 由恵, 筒井, 容平, Velazquez, J. J. L. メールアドレス: 所属: Kyushu University, Waseda University, Universitat Bonn
URL	https://ocu-omu.repo.nii.ac.jp/records/2016785

Global solutions to a chemotaxis system with non-diffusive memory

Y. Sugiyama, Y. Tsutsui and J.J.L. Velazquez

Citation	OCAMI Preprint Series
Issue Date	2013
Type	Preprint
Textversion	Author
Relation	The following article has been submitted to Journal of Mathematical Analysis and Applications. This is not the published version. Please cite only the published version. The article has been published in final form at https://doi.org/10.1016/j.jmaa.2013.08.065 .
Is version of	https://doi.org/10.1016/j.jmaa.2013.08.065 .

From: Osaka City University Advanced Mathematical Institute

<http://www.sci.osaka-cu.ac.jp/OCAMI/publication/preprint/preprint.html>

Global solutions to a chemotaxis system with non-diffusive memory

Y. Sugiyama*, Y. Tsutsui† and J.J.L. Velázquez‡

Abstract

In this article, a existence theorem of global solutions with small initial data belonging to $L^1 \cap L^p$, ($n < p \leq \infty$) for a chemotaxis system are given on the whole space \mathbb{R}^n , $n \geq 3$. In the case $p = \infty$, our global solution is integrable with respect to the space variable on some time interval, and then conserves the mass for a short time, at least. The system consists of a chemotaxis equation with a logarithmic term and an ordinary equation without diffusion term.

Keywords chemotaxis system, global solutions.

2000 Mathematics Subject Classification 35A01, 92C17

1 Introduction

This paper concerns global solutions to the system

$$(E_\lambda) \begin{cases} \partial_t u = \Delta u - \nabla \cdot \left(u \frac{\nabla v}{v} \right), & x \in \mathbb{R}^n, \quad t > 0 \\ \partial_t v = uv^\lambda, & x \in \mathbb{R}^n, \quad t > 0 \quad \text{and} \quad \lambda \in \mathbb{R} \\ u(0, x) = a(x) \geq 0 \quad v(0, x) = b(x) \geq 0, & x \in \mathbb{R}^n. \end{cases}$$

Here, u is the unknown cell density of the chemotactic species and v is the unknown density of non-diffusive chemical substance, which is produced by the species. This system is a particular case of Keller-Segel system [8] and related to the dynamics of self-reinforced random walks [14] [16], and also used as haptotaxis and angiogenesis models. One of interesting features of the system is the absence of diffusion term in the second equation. There are many papers which studied the classical Keller-Segel system with diffusion term in the second equation. Levine and Sleeman [10] investigated finite time blow-up phenomena for the system (E_1) in one dimensional case. Additional properties for the solutions of (E_1) have been obtained in [12]. In smooth bounded domains in \mathbb{R}^n with $\lambda \leq 1$, Rascle [15] and Yang, Chen and Liu [20] showed the existence of global solutions for the system. Corrias, Perthame and Zaag [5], [6] studied the same topics in a general system, which do not cover the case $\lambda = 1$. In [17], asymptotic behavior of radial symmetric solutions to (E_λ) is studied. When $\lambda \in [0, 1)$ and $n = 1$, Kang, Stevens and Velázquez proved that for some initial data the corresponding solutions u tends to Dirac mass as $t \rightarrow \infty$ in [7].

Through the transformation

$$z = \frac{v^{1-\lambda}}{1-\lambda} \quad \text{with} \quad \theta = \frac{1}{1-\lambda} \in \mathbb{R},$$

*Faculty of Mathematics, Kyushu University, Fukuoka 819-0395, Japan. e-mail: sugiyama@math.kyushu-u.ac.jp

†Department of Mathematics, Faculty of Science and Engineering, Waseda University, Tokyo, Japan. e-mail: y.tsutsui@kurenai.waseda.jp

‡Universität Bonn, Institute für Angewandte Mathematik, Endencher Allee 60, D-53115 Bonn, Germany. e-mail: velazquez@iam.uni-bonn.de

the system (E_λ) becomes

$$(\tilde{E}_\theta) \begin{cases} \partial_t u = \Delta u - \theta \nabla \cdot \left(u \frac{\nabla z}{z} \right), & x \in \mathbb{R}^n, \quad t > 0 \\ \partial_t z = u, & x \in \mathbb{R}^n, \quad t > 0 \\ u(0, x) = a(x) \quad z(0, x) = c(x) = \theta b(x)^{1-\lambda}, & x \in \mathbb{R}^n. \end{cases}$$

This system is invariant with respect to the scaling

$$(u(t, x), z(t, x)) \mapsto (\mu^{\alpha+2} u(\mu^2 t, \mu x), \mu^\alpha z(\mu^2 t, \mu x))$$

for all $\alpha \in \mathbb{R}$. In this article, we give a global existence theorem of solutions to the system (\tilde{E}_θ) with the special initial data $c \equiv 1$, which corresponds to the case $\lambda < 1$ in (E_λ) , in the sense of mild solutions, more precisely, we construct solutions to the integral equations;

$$(I.E.) \begin{cases} u(t) = e^{t\Delta} a - \theta B[u](t), \text{ where } B[u](t) = \int_0^t e^{(t-\tau)\Delta} \nabla \cdot \left(u(\tau) \frac{\nabla z(\tau)}{z(\tau)} \right) d\tau \\ z(t) = 1 + \int_0^t u(\tau) d\tau. \end{cases}$$

As a consequence of the absence of diffusion term in the second equation, the regularity of z and ∇z with respect to space variable is not better than that of 1 and ∇u respectively. For the initial data $c \equiv -1$ that corresponds to another case $\lambda > 1$ in (E_λ) , the same results as in below hold. D. Li, K. Li and Zhao [11] treated with the case $\lambda = 1$ in which the system is changed to a hyperbolic-parabolic system through the transform $V = -\frac{\nabla v}{v}$, and constructed local and global solutions in Sobolev spaces with positive smoothness. Very recently, Ahn and Kang [1] proved the local existence of solutions to (E_λ) with $\lambda > 0$, and the non-existence of self-similar solutions.

Our main result reads as follows

Theorem 1.1 (Small data global existence). *Let $n \geq 3$, $n/(n-1) < q < n < r < p \leq \infty$ and $\theta \in \mathbb{R}$. There exists $\delta = \delta(n, p, q, r, |\theta|) > 0$ such that if $\|a\|_{L^1 \cap L^p} \leq \delta$, then there exists a global solution $u \in L^1(0, \infty; L^\infty(\mathbb{R}^n))$ of (I.E.) satisfying $\|u\|_{X_t^1} + \|u\|_{X_\infty^2} + \|u\|_{X_\infty^3} \lesssim \|a\|_{L^1 \cap L^p}$ where*

$$\|u\|_{X_t^1} = \int_0^t \|u(\tau)\|_{L^\infty} d\tau, \\ \|u\|_{X_t^2} = \sup_{\tau < t} \left\| \int_0^\tau \nabla u(\sigma) d\sigma \right\|_{L^r} \quad \text{and} \quad \|u\|_{X_t^3} = \sup_{\tau < t} \left\| \int_0^\tau \nabla u(\sigma) d\sigma \right\|_{L^q}.$$

Moreover, with $U_\infty(t) = \int_0^t \|u(\tau)\|_{L^\infty} d\tau$ and $U(t, x) = \int_0^t u(\tau, x) d\tau$, one has

$$U_\infty \in C([0, \infty)) \quad \text{and} \quad \nabla U \in C([0, \infty); L^r \cap L^q). \quad (1)$$

Remark 1.1. *The smallness assumption on the initial data a is used to guarantee*

$$\sup_{t>0} \left\| \frac{1}{z_0(t)} \right\|_{L^\infty} \lesssim 1, \quad \text{where} \quad z_0(t) = 1 + \int_0^t e^{\tau\Delta} a d\tau.$$

Indeed, $\int_0^\infty \|e^{\tau\Delta} a\|_{L^\infty} d\tau \approx \|a\|_{\dot{B}_{\infty,1}^{-2}}$ and $L^1 \cap L^p \hookrightarrow \dot{B}_{\infty,1}^{-2}$. We then have

$$|z_0(t, x)| \geq 1 - \int_0^\infty |e^{\tau\Delta} a(x)| d\tau \geq 1/2.$$

And, this property is preserved by the evolution of the system and $\sup_{t>0} \left\| \frac{1}{z(t)} \right\|_{L^\infty} \lesssim 1$.

Because the first equation of (\tilde{E}_θ) is a divergence form, there formally holds the conservation of mass;

$$\int_{\mathbb{R}^n} u(t) dx = \int_{\mathbb{R}^n} a dx \quad (t \geq 0). \quad (2)$$

However, we do not know whether or not the global solution, constructed in Theorem 1.1, is in $L^1(\mathbb{R}^n)$. The following theorem ensures that the solution belongs to $L^1(\mathbb{R}^n)$ in some time interval, when $p = \infty$ in Theorem 1.1. As a consequence, we can make sure that our global solution constructed in Theorem 1.1 fulfills the conservation of mass (2) for a short time, at least.

Theorem 1.2 (Large data local existence). *Let $n \geq 1$, $0 < s < 1$ and $\theta \in \mathbb{R}$.*

(i): *There exists a small constant $C = C(n, s, |\theta|) \in (0, 1)$ such that for any $a \in L^\infty$, we can find a local solution $u \in L^\infty((0, T) \times \mathbb{R}^n)$ with $T = \frac{C}{\|a\|_{L^\infty}}$ satisfying*

$$\|u\|_{Y_T^1} + \|u\|_{Y_T^2} + \|u\|_{Y_T^3} + \|u\|_{Y_T^4} \lesssim \|a\|_{L^\infty}$$

where for $t > 0$

$$\begin{aligned} \|u\|_{Y_t^1} &= \sup_{\tau < t} \|u(\tau)\|_{L^\infty}, \quad \|u\|_{Y_t^2} = \sup_{\tau < t} t^{s/2} \|u(\tau)\|_{\dot{B}_{\infty, \infty}^s}, \\ \|u\|_{Y_t^3} &= \sup_{\tau < t} \tau^{1/2} \|\nabla u(\tau)\|_{L^\infty} \quad \text{and} \quad \|u\|_{Y_t^4} = \sup_{\tau < t} \tau^{(1+s)/2} \|\nabla u(\tau)\|_{\dot{B}_{\infty, \infty}^s}. \end{aligned}$$

(ii): *In addition, if the initial data a also belongs to $L^1(\mathbb{R}^n)$, then the solution u , constructed in (i), is in $C([0, T]; L^1)$ where $T = \frac{C}{\|a\|_{L^1 \cap L^\infty}}$ and C depending on n , s and $|\theta|$, and fulfills*

$$\|u\|_{Z_t} = \sup_{\tau < t} \|u(\tau)\|_{L^1} \lesssim \|a\|_{L^1}.$$

Remark 1.2. 1. *The condition on T guarantees the lower bound of $1/z$. Indeed, for all t and x*

$$|z_0(t, x)| \geq 1 - \int_0^t \|e^{\tau \Delta} a\|_{L^\infty} d\tau \geq 1 - C > 0.$$

2. *Ahn and Kang [1] proved the local existence under the assumption $a, c (= z(0)) \in L^\infty \cap W^{1,p}$ with $p > n$ and certain lower bound of $c = z(0)$.*

Uniqueness of solutions in Theorem 1.2 is valid as follows;

Theorem 1.3 (Uniqueness). *Let $n \geq 1$, $0 < s < 1$ and $T \in (0, \infty)$.*

(i): *If u and v are solutions in the class $C((0, T); L^\infty) \cap \cap_{j=2}^4 Y_T^j$ with the same initial data $a \in L^\infty$, then $u(t, x) = v(t, x)$ for all $t \in (0, T)$ and a.e. $x \in \mathbb{R}^n$.*

(ii): *If u and v are solutions in the class $C([0, T]; L^1) \cap \cap_{j=1}^4 Y_T^j$ with the same initial data $a \in L^1 \cap L^\infty$, then $u(t, x) = v(t, x)$ for all $t \in (0, T)$ and a.e. $x \in \mathbb{R}^n$.*

This paper is organized as follows. In next section, we recall the definition and equivalence norms of (homogeneous) Besov space and establish our basic estimates. Theorems 1.1 and 1.2 are proved in Sections 3 and 4, respectively. In Section 5, the proof of Theorem 1.3 is provided.

2 Preliminaries

Throughout this paper we use the following notations. \mathcal{S} and \mathcal{S}' denote the Schwartz spaces of rapidly decreasing smooth functions and tempered distributions, respectively. $A \lesssim B$ means $A \leq cB$ with a positive constant c . $A \approx B$ means $A \lesssim B$ and $B \lesssim A$. $C = C(a, b, c)$ means C depends on a, b and c .

Let us recall the definition of Besov spaces. We fix $\varphi \in \mathcal{S}(\mathbb{R}^n)$ satisfying $\text{supp } \varphi \subset \{1/2 \leq |\xi| \leq 2\}$ and $\sum_{j \in \mathbb{Z}} \varphi\left(\frac{\xi}{2^j}\right) = 1$ for $\xi \in \mathbb{R}^n \setminus \{0\}$, and then $\varphi_j(D)f = \mathcal{F}^{-1} \left[\varphi\left(\frac{\cdot}{2^j}\right) \hat{f}(\cdot) \right]$.

Definition 2.1. Let $0 < p, q \leq \infty$ and $s \in \mathbb{R}$. Besov space $\dot{B}_{p,q}^s$ is defined to be the space of $f \in \mathcal{S}'$ modulo polynomials such that

$$\|f\|_{\dot{B}_{p,q}^s} = \left(\sum_{j \in \mathbb{Z}} 2^{jsq} \|\varphi_j(D)f\|_{L^p}^q \right)^{1/q}, \quad q < \infty,$$

$$\|f\|_{\dot{B}_{p,\infty}^s} = \sup_{j \in \mathbb{Z}} 2^{js} \|\varphi_j(D)f\|_{L^p}.$$

It is well-known that $\dot{B}_{\infty,\infty}^s$ with $0 < s \notin \mathbb{N}$ coincides with Hölder space \dot{C}^s , i.e.,

$$\|f\|_{\dot{B}_{\infty,\infty}^s} \approx \|f\|_{\dot{C}^s} = \sum_{|\alpha|=[s]} \sup_{x \neq y} \frac{|\partial^\alpha f(x) - \partial^\alpha f(y)|}{|x - y|^{s-[s]}},$$

where $[s]$ means the integer part of s , see for example [4]. Triebel [19] showed that Besov spaces are also characterized by means of heat semigroups $e^{t\Delta}$; with a non-negative integer $m > s/2$

$$\|f\|_{\dot{B}_{p,q}^s} \approx \left(\int_0^\infty \left(t^{m-s/2} \|(-\Delta)^m e^{t\Delta} f\|_{L^p} \right)^q \frac{dt}{t} \right)^{1/q}, \quad q < \infty,$$

$$\|f\|_{\dot{B}_{p,\infty}^s} \approx \sup_{t > 0} t^{m-s/2} \|(-\Delta)^m e^{t\Delta} f\|_{L^p}. \quad (3)$$

The following lemma is our basic estimate for the proof of Theorem 1.1.

Lemma 2.1 (Estimates for the non-linear term). Let $n \geq 1$. For vector functions $F = (f_1, \dots, f_n)$, we have the following.

(i): For $1 \leq p < n < q \leq \infty$,

$$\int_0^\infty \left\| \int_0^t e^{(t-\tau)\Delta} \nabla \cdot F(\tau) d\tau \right\|_{L^\infty} dt \lesssim \int_0^\infty \|F(\tau)\|_{\dot{B}_{\infty,1}^{-1}} d\tau \lesssim \int_0^\infty \|F(\tau)\|_{L^p \cap L^q} d\tau.$$

(ii): For all $r \in (1, \infty)$

$$\left\| \int_0^T \nabla \int_0^t e^{(t-\tau)\Delta} \nabla \cdot F(\tau) d\tau dt \right\|_{L^r} \lesssim \int_0^T \|F(\tau)\|_{L^r} d\tau.$$

Proof. (i): Changing order of integrals, we see that the right hand side is bounded by

$$\int_0^\infty \int_0^\infty \|e^{t\Delta} \nabla \cdot F(\tau)\|_{L^\infty} dt d\tau \approx \int_0^\infty \|F(\tau)\|_{\dot{B}_{\infty,1}^{-1}} d\tau$$

and then from the embedding $L^p \cap L^q \hookrightarrow \dot{B}_{\infty,1}^{-1}$, the desired inequality is verified.

(ii): We also change order of integrals and have that the right hand side is bounded by

$$\int_0^T \left\| \nabla \int_0^{T-\tau} e^{t\Delta} \nabla \cdot F(\tau) dt \right\|_{L^r} d\tau.$$

Though the boundedness of Riesz transform on $L^r(\mathbb{R}^n)$ and the duality, the L^r norm above is dominated by

$$\left| \int_{\mathbb{R}^n} \left(\int_0^{T-\tau} e^{t\Delta} F(\tau) dt \right) \Delta G dx \right|$$

with some $G = (g_1, \dots, g_n)$ and each $g_j \in \mathcal{S}$ satisfying $\|g_j\|_{L^{r'}} \leq 1$. The proof is completed as follows;

$$\begin{aligned} \int_0^T \left| \int_{\mathbb{R}^n} F(\tau) \int_0^{T-\tau} \partial_t e^{t\Delta} G dt dx \right| d\tau &= \int_0^T \left| \int_{\mathbb{R}^n} F(\tau) \left(e^{(T-\tau)\Delta} G - G \right) dx \right| d\tau \\ &\lesssim \int_0^T \|F(\tau)\|_{L^r} d\tau \|G\|_{L^{r'}}. \end{aligned}$$

□

The following estimates for products are verified by paraproduct formula due to Bony [2] and is applied in the proof of Theorem 1.2.

Lemma 2.2. For $s > 0$,

$$\|fg\|_{\dot{B}_{\infty,\infty}^s} \lesssim \|f\|_{\dot{B}_{\infty,\infty}^s} \|g\|_{L^\infty} + \|f\|_{L^\infty} \|g\|_{\dot{B}_{\infty,\infty}^s}.$$

Since it is not hard to prove this, we omit the proof.

Decay estimates of the heat semigroup $e^{t\Delta}$ on Besov spaces are used in the proof.

Lemma 2.3 ([9]). If $\beta < \alpha$, it follows

$$\|e^{t\Delta}f\|_{\dot{B}_{\infty,1}^\beta} \lesssim t^{-(\alpha-\beta)/2} \|f\|_{\dot{B}_{\infty,\infty}^\alpha}.$$

See [9] for the proof.

The following estimate is useful to control the norm Y_t^4 . For the We give a proof based on a method due to Meyer [13], which involves the real interpolation theory.

Lemma 2.4. For any $\alpha \in \mathbb{R}$, there holds

$$\left\| \int_0^t e^{(t-\tau)\Delta} \nabla f(\tau) d\tau \right\|_{\dot{B}_{\infty,\infty}^{\alpha+1}} \lesssim \sup_{\tau < t} \|f(\tau)\|_{\dot{B}_{\infty,\infty}^\alpha}.$$

Proof. We define $\tilde{f}(\tau) = f(\tau)\chi_{(0,t)}(\tau)$ where χ_I is the characteristic function of $I \subset \mathbb{R}$, and decompose

$$\int_0^t e^{(t-\tau)\Delta} \nabla f(\tau) d\tau = \int_0^\infty e^{\tau\Delta} \nabla \tilde{f}(t-\tau) d\tau = I_A + II_A,$$

where $A > 0$ and

$$I_A = \int_0^A \nabla^\theta e^{\tau\Delta} \nabla^{1-\theta} \tilde{f}(t-\tau) d\tau, \quad II_A = \int_A^\infty \nabla^{\tilde{\theta}} e^{\tau\Delta} \nabla^{1-\tilde{\theta}} \tilde{f}(t-\tau) d\tau.$$

Here, $0 < \theta, \tilde{\theta} < 1$. From Lemma 2.3, one has that

$$\begin{aligned} \|I_A\|_{\dot{B}_{\infty,1}^{\alpha-1/2}} &\lesssim \int_0^A \tau^{-3/4} \|\tilde{f}(t-\tau)\|_{\dot{B}_{\infty,\infty}^{\alpha-1}} d\tau \lesssim A^{1/4} \sup_{\tau < t} \|f(\tau)\|_{\dot{B}_{\infty,\infty}^{\alpha-1}} \\ \|II_A\|_{\dot{B}_{\infty,1}^{\alpha+1/2}} &\lesssim \int_A^\infty \tau^{-5/4} \|\tilde{f}(t-\tau)\|_{\dot{B}_{\infty,\infty}^{\alpha-1}} d\tau \lesssim A^{-1/4} \sup_{\tau < t} \|f(\tau)\|_{\dot{B}_{\infty,\infty}^{\alpha-1}}. \end{aligned}$$

Applying the well-known property of Besov spaces, [18] or [3];

$$\dot{B}_{\infty,\infty}^\alpha = \left(\dot{B}_{\infty,\infty}^{\alpha-1/2}, \dot{B}_{\infty,\infty}^{\alpha+1/2} \right)_{1/2,\infty},$$

we end the proof as follows;

$$\begin{aligned} \left\| \int_0^t e^{(t-\tau)\Delta} \nabla f(\tau) d\tau \right\|_{\dot{B}_{\infty,\infty}^{\alpha+1}} &\lesssim \sup_{\lambda > 0} \lambda^{-1/2} \left(\|I_{A_\lambda}\|_{\dot{B}_{\infty,1}^{\alpha-1/2}} + \lambda \|II_{A_\lambda}\|_{\dot{B}_{\infty,1}^{\alpha+1/2}} \right) \\ &\lesssim \sup_{\tau < t} \|f(\tau)\|_{\dot{B}_{\infty,\infty}^{\alpha-1}}. \end{aligned}$$

□

3 Proof of Theorem 1.1

Let

$$\|u\|_{X_t} = \|u\|_{X_t^1} + \|u\|_{X_t^2} + \|u\|_{X_t^3}.$$

The assumption on exponents ensures the following linear estimates.

Lemma 3.1 (Linear estimates).

$$\|e^{-\Delta}a\|_{X_\infty} \lesssim \|a\|_{\dot{B}_{\infty,1}^{-2}} + \|a\|_{\dot{B}_{r,1}^{-1}} + \|a\|_{\dot{B}_{q,1}^{-1}} \lesssim \|a\|_{L^1 \cap L^p}.$$

Proof. The first inequality follows from (3). From

$$\begin{aligned} \|a\|_{\dot{B}_{\infty,1}^{-2}} &\lesssim \sum_{j \leq 0} 2^{j(n-2)} \|a\|_{L^1} + \sum_{j \geq 1} 2^{j(n/p-2)} \|a\|_{L^p} \\ \|a\|_{\dot{B}_{l,1}^{-1}} &\lesssim \sum_{j \leq 0} 2^{j(n/l'-1)} \|a\|_{L^1} + \sum_{j \geq 1} 2^{-j} \|a\|_{L^l}, \quad (l = r, q) \quad \text{and} \end{aligned}$$

the interpolation inequality $\|a\|_{L^r \cap L^q} \leq \|a\|_{L^1 \cap L^p}$, the second inequality is showed. \square

With lemmas 2.1 and 3.1, we construct solutions by successive approximation;

$$u_0(t) = e^{t\Delta}a \quad \text{and} \quad u_{m+1}(t) = u_0(t) - B[u_m](t)$$

where

$$z_m(t) = 1 + \int_0^t u_m(\tau) d\tau.$$

We begin the proof of Theorem 1.1 with showing that $\{u_m\}_m$ is a bounded sequence in X_∞ .

From Lemma 3.1, $\|u_0\|_{X_\infty} \leq C_X \|a\|_{L^1 \cap L^p}$ where the constant C_X depends on n, p, q and r . Taking δ so that

$$\delta \leq \frac{1}{4C_X}, \tag{4}$$

we see that $\sup_{t>0} \left\| \frac{1}{z_0(t)} \right\|_{L^\infty} \leq 4/3$. Assuming $\|u_m\|_{X_\infty} \leq 2C_X \|a\|_{L^1 \cap L^p}$, we shall check $\|u_{m+1}\|_{X_\infty} \leq 2C_X \|a\|_{L^1 \cap L^p}$. First, the assumption implies $\sup_{t>0} \left\| \frac{1}{z_m(t)} \right\|_{L^\infty} \leq 2$. Hence, applying Lemma 2.1 one can obtain

$$\begin{cases} \|B[u_{m+1}]\|_{X_\infty^1} \lesssim \|u_m\|_{X_\infty^1} (\|u_m\|_{X_\infty^2} + \|u_m\|_{X_\infty^3}) \\ \|B[u_{m+1}]\|_{X_\infty^2} \lesssim \|u_m\|_{X_\infty^1} \|u_m\|_{X_\infty^2} \\ \|B[u_{m+1}]\|_{X_\infty^3} \lesssim \|u_m\|_{X_\infty^1} \|u_m\|_{X_\infty^3}. \end{cases}$$

Combining this with Lemma 3.1, one has

$$\|u_{m+1}\|_{X_\infty} \leq C_X \|a\|_{L^1 \cap L^p} + C_X^* |\theta| \|u_m\|_{X_\infty}^2$$

with the constant C_X^* depending on n, p, q and r . Then, taking small δ so that

$$\delta \leq \frac{1}{4C_X C_X^* |\theta|} \tag{5}$$

we see $\|u_{m+1}\|_{X_\infty} \leq 2C_X \|a\|_{L^1 \cap L^p}$. Therefore, $\{u_m\}_m$ is bounded in X_∞ . For simplicity, let $K_X = \sup_{m \in \mathbb{N} \cup \{0\}} \|u_m\|_{X_\infty} \leq 2C_X \|a\|_{L^1 \cap L^p}$.

To show that $\{u_m\}_m$ is also a Cauchy sequence in X_∞ , we divide $u_{m+1}(t) - u_m(t)$ into three parts;

$$u_{m+1}(t) - u_m(t) = \theta \int_0^t e^{(t-\tau)\Delta} \nabla \cdot (A_m(\tau) + B_m(\tau) + C_m(\tau)) d\tau$$

where

$$\begin{cases} A_m(\tau) = -\frac{u_m(\tau) - u_{m-1}(\tau)}{z_{m-1}(\tau)} \nabla z_{m-1}(\tau) \\ B_m(\tau) = \frac{u_m(\tau) (z_m(\tau) - z_{m-1}(\tau))}{z_{m-1}(\tau) z_m(\tau)} \nabla z_{m-1}(\tau) \\ C_m(\tau) = -\frac{u_m(\tau)}{z_m(\tau)} \nabla (z_m(\tau) - z_{m-1}(\tau)). \end{cases}$$

From Lemma 2.1, it follows

$$\|u_{m+1} - u_m\|_{X_\infty^1} \lesssim |\theta| \int_0^\infty \|A_m(\tau) + B_m(\tau) + C_m(\tau)\|_{L^q \cap L^r} d\tau.$$

and

$$\begin{cases} \|A_m(\tau)\|_{L^q \cap L^r} \lesssim \|u_m(\tau) - u_{m-1}(\tau)\|_{L^\infty} (\|u_{m-1}\|_{X_\infty^2} + \|u_{m-1}\|_{X_\infty^3}) \\ \|B_m(\tau)\|_{L^q \cap L^r} \lesssim \|u_m(\tau)\|_{L^\infty} \|u_m - u_{m-1}\|_{X_\infty^1} (\|u_{m-1}\|_{X_\infty^2} + \|u_{m-1}\|_{X_\infty^3}) \\ \|C_m(\tau)\|_{L^q \cap L^r} \lesssim \|u_m(\tau)\|_{L^\infty} (\|u_m - u_{m-1}\|_{X_\infty^2} + \|u_m - u_{m-1}\|_{X_\infty^3}), \end{cases}$$

which imply

$$\|u_{m+1} - u_m\|_{X_\infty^1} \lesssim |\theta| K_X (1 + K_X) \|u_m - u_{m-1}\|_{X_\infty}.$$

The similar argument yields

$$\|u_{m+1} - u_m\|_{X_\infty} \lesssim |\theta| K_X (1 + K_X) \|u_m - u_{m-1}\|_{X_\infty}.$$

Since $K \leq 1/2$, one obtains $\|u_{m+1} - u_m\|_{X_\infty} \leq C_X^{**} |\theta| K \|u_m - u_{m-1}\|_{X_\infty}$ where the constant C_X^{**} depends on n, p, q and r . Taking small δ so that

$$\delta \leq \frac{1}{4C_X C_X^{**} |\theta|}, \quad (6)$$

we see then $\{u_m\}_m$ is a Cauchy sequence in X_∞ .

To end this proof, we provide a proof of the continuity of U_∞ and ∇U . For $\tilde{t} > t \geq 0$, since $\int_t^{\tilde{t}} \|u(\tau)\|_{L^\infty} d\tau = \int_t^{\tilde{t}} \|e^{\tau\Delta} a\|_{L^\infty} d\tau - |\theta|(\text{I} + \text{II})$ where

$$\begin{aligned} \text{I} &\leq \int_0^t \int_{t-\sigma}^{\tilde{t}-\sigma} \left\| e^{\tau\Delta} \nabla \left(u(\sigma) \frac{\nabla z(\sigma)}{z(\sigma)} \right) \right\|_{L^\infty} d\tau d\sigma \quad \text{and} \\ \text{II} &\leq \int_t^{\tilde{t}} \int_0^{\tilde{t}-\sigma} \left\| e^{\tau\Delta} \nabla \left(u(\sigma) \frac{\nabla z(\sigma)}{z(\sigma)} \right) \right\|_{L^\infty} d\tau d\sigma, \end{aligned}$$

Lebesgue's dominated convergence theorem ensures

$$U_\infty \in C([0, \infty); L^\infty) \text{ and } U_\infty(t) \rightarrow 0 \text{ as } t \searrow 0.$$

The continuity $\nabla U \in C([0, \infty); L^\infty \cap L^q)$ and the convergence $\nabla U(t) \rightarrow 0$ in $L^\infty \cap L^q$ as $t \searrow 0$ are also verified by the same argument above. The proof is completed.

Remark 3.1. For any $\alpha \in (0, 1 - n/r)$,

$$\begin{aligned} \int_0^\infty \|u(t)\|_{\dot{B}_{\infty, \infty}^\alpha} dt &\lesssim \|a\|_{\dot{B}_{\infty, 1}^{\alpha-2}} + |\theta| \|u\|_{X_1} (\|u\|_{X_2} + \|u\|_{X_3}) \\ &\lesssim \|a\|_{L^1 \cap L^p} + |\theta| \|u\|_X^2. \end{aligned}$$

4 Proof of Theorem 2

Let

$$\|u\|_{Y_t} = \|u\|_{Y_t^1} + \|u\|_{Y_t^2} + \|u\|_{Y_t^3} + \|u\|_{Y_t^4}.$$

(i): We verify that $\{u_m\}_m$ is also a Cauchy sequence in Y_T with some T . We have $\|u_0\|_{Y_T} \leq C_Y \|a\|_{L^\infty}$. By Lemma 2.2 and

$$\left\| \frac{1}{z_m(t)} \right\|_{\dot{B}_{\infty,\infty}^s} \approx \left\| \frac{1}{z_m(t)} \right\|_{\dot{C}^s} \lesssim \int_0^t \|u_m(\tau)\|_{\dot{B}_{\infty,\infty}^s} d\tau, \quad (7)$$

one can show that if $\sup_{t < T} \left\| \frac{1}{z_m(t)} \right\|_{L^\infty} \lesssim 1$,

$$\begin{cases} \|B[u_{m+1}]\|_{Y_T^1} \lesssim T \|u_m\|_{Y_T^1} \|u_m\|_{Y_T^3} \\ \|B[u_{m+1}]\|_{Y_T^2} \lesssim T \|u_m\|_{Y_T^2} \|u_m\|_{Y_T^3} + T \|u_m\|_{Y_T^1} \left(T \|u_m\|_{Y_T^2} \|u_m\|_{Y_T^3} + \|u_m\|_{Y_T^4} \right) \\ \|B[u_{m+1}]\|_{Y_T^3} \lesssim T \|u_m\|_{Y_T^2} \|u_m\|_{Y_T^3} + T \|u_m\|_{Y_T^1} \left(T \|u_m\|_{Y_T^2} \|u_m\|_{Y_T^3} + \|u_m\|_{Y_T^4} \right) \\ \|B[u_{m+1}]\|_{Y_T^4} \lesssim T \|u_m\|_{Y_T^2} \|u_m\|_{Y_T^3} + T \|u_m\|_{Y_T^1} \left(T \|u_m\|_{Y_T^2} \|u_m\|_{Y_T^3} + \|u_m\|_{Y_T^4} \right), \end{cases}$$

which imply

$$\|u_{m+1}\|_{Y_T} \leq C_Y \|a\|_{L^\infty} + C_Y^* |\theta| T (1 + T \|u_m\|_{Y_T}) \|u_m\|_{Y_T}^2.$$

Here, Lemma 2.2 and (7) have been applied for Y_T^2 , and Lemma 2.4 and (7) for Y_T^3 . Therefore, taking

$$T \leq \frac{1}{2C_Y \|a\|_{L^\infty}} \quad \text{and} \quad T \leq \frac{1}{8C_Y C_Y^* |\theta| \|a\|_{L^\infty}} \quad (8)$$

and assuming $\|u_m\|_{Y_T} \leq 2C_Y \|a\|_{L^\infty}$, we have $\|u_{m+1}\|_{Y_T} \leq 2C_Y \|a\|_{L^\infty}$. Indeed, from this assumption, $|z_m(t, x)| \geq 1/2$. Let $K_Y(T) = \sup_{m \in \{0\} \cup \mathbb{N}} \|u_m\|_{Y_T} \leq 2C_Y \|a\|_{L^\infty}$.

And then, we have

$$\begin{cases} \|A_m(t) + B_m(t) + C_m(t)\|_{L^\infty} \lesssim t^{1/2} K_Y(t) (1 + t K_Y(t)) \|u_m - u_{m-1}\|_{Y_t} \\ \|A_m(t) + B_m(t) + C_m(t)\|_{\dot{B}_{\infty,\infty}^s} \lesssim t^{(1-s)/2} K_Y(t) (1 + t K_Y(t))^2 \|u_m - u_{m-1}\|_{Y_t}. \end{cases}$$

Here, Lemma 2.2 and

$$\left\| \frac{1}{z_{m-1}(t)z_m(t)} \right\|_{\dot{B}_{\infty,\infty}^s} \lesssim \left\| \frac{1}{z_{m-1}(t)} \right\|_{\dot{B}_{\infty,\infty}^s} + \left\| \frac{1}{z_m(t)} \right\|_{\dot{B}_{\infty,\infty}^s} \lesssim t^{1-s/2} K_Y(t)$$

have been used in the second estimate. Then, we have

$$\|u_{m+1} - u_m\|_{Y_T} \leq C_Y^{**} |\theta| K_Y(T) T \|u_m - u_{m-1}\|_{Y_T}.$$

If

$$T \leq \frac{1}{10C_Y C_Y^{**} |\theta| \|a\|_{L^\infty}}, \quad (9)$$

$\{u_m\}_m$ is a Cauchy sequence in Y_T .

(ii): We shall show $\{u_m\}_m$ is also a Cauchy sequence in $C([0, T]; L^1(\mathbb{R}^n))$ with some T . Let $m \in \mathbb{N} \cup \{0\}$ and we assume $\|u_m\|_{Z_T} \leq 2\|a\|_{L^1}$. Of course, $\|u_0\|_{Z_T} \leq \|a\|_{L^1}$. Because if T fulfills (8) one has

$$\|u_{m+1}\|_{Z_T} \leq \|a\|_{L^1} + C_Y^{**} |\theta| T \|u_m\|_{Z_T} \|u_m\|_{Y_T^3} \quad (10)$$

with a constant C_Y^{**} depending on n only, taking

$$T \leq \frac{1}{2C_Y C_Y^{**} |\theta| \|a\|_{L^\infty}}, \quad (11)$$

we then have $\|u_{m+1}\|_{Z_T} \leq 2\|a\|_{L^1}$. Hence, we write $K_Z(T) = \sup_{m \in \mathbb{N} \cup \{0\}} \|u_m\|_{Z_T} \leq 2\|a\|_{L^1}$ with T fulfills (8) and (11). To show $u_m \in C([0, \infty); L^1(\mathbb{R}^n))$ we write that for small $\varepsilon > 0$

$$\begin{aligned} u_{m+1}(t + \varepsilon) - u_{m+1}(t) &= e^{t\Delta} (e^{\varepsilon\Delta} a - a) \\ &\quad + \theta \int_0^t \nabla e^{(t-\tau)\Delta} (e^{\varepsilon\Delta} - I_d) \left(u_m(\tau) \frac{\nabla z_m(\tau)}{z_m(\tau)} \right) d\tau \\ &\quad - \theta \int_t^{t+\varepsilon} e^{(t+\varepsilon-\tau)\Delta} \nabla \left(u_m(\tau) \frac{\nabla z_m(\tau)}{z_m(\tau)} \right) d\tau. \end{aligned}$$

Combining this with Lebesgue's dominated convergence theorem ensures that the right continuity of u_{m+1} in the topology of L^1 . On the other hand, since

$$\begin{aligned} u_{m+1}(t) - u_{m+1}(t - \varepsilon) &= e^{(t-\varepsilon)\Delta} (e^{\varepsilon\Delta} a - a) \\ &\quad + \theta \int_0^{t-\varepsilon} \nabla e^{(t-\varepsilon-\tau)\Delta} (e^{\varepsilon\Delta} - I_d) \left(u_m(\tau) \frac{\nabla z_m(\tau)}{z_m(\tau)} \right) d\tau \\ &\quad - \theta \int_{t-\varepsilon}^t e^{(t-\tau)\Delta} \nabla \left(u_m(\tau) \frac{\nabla z_m(\tau)}{z_m(\tau)} \right) d\tau, \end{aligned}$$

we have the left continuity of u_{m+1} from the Lebesgue's theorem again.

Next, we observe

$$\|u_{m+1} - u_m\|_{Z_T} \lesssim |\theta| \sup_{t < T} \int_0^t (t - \tau)^{-1/2} \|A_m(\tau) + B_m(\tau) + C_m(\tau)\|_{L^1} d\tau$$

and

$$\begin{cases} \|A_m(\tau)\|_{L^1} \lesssim \tau^{1/2} \|u_m - u_{m-1}\|_{Z_T} \|u_{m-1}\|_{Y_T^3} \\ \|B_m(\tau)\|_{L^1} \lesssim \tau^{3/2} \|u_m\|_{Z_T} \|u_{m-1}\|_{Y_T^3} \|u_m - u_{m-1}\|_{Y_T^1} \\ \|C_m(\tau)\|_{L^1} \lesssim \tau^{1/2} \|u_m\|_{Z_T} \|u_m - u_{m-1}\|_{Y_T^3} \end{cases}$$

and then for T satisfying (11) we have

$$\begin{aligned} \|u_{m+1} - u_m\|_{Z_T} &\leq C_Z |\theta| K_Y(T) T \|u_m - u_{m-1}\|_{Z_T} \\ &\quad + C_Z |\theta| K_Y(T) K_Z(T) T^2 \|u_m - u_{m-1}\|_{Y_T} + C_Z |\theta| K_Z(T) T \|u_m - u_{m-1}\|_{Y_T}. \end{aligned}$$

where the constant C_Z depends on n only. Here, taking

$$T \leq \frac{1}{10C_Y C_Z (1 + |\theta|) \|a\|_{L^\infty}} \quad \text{and} \quad T \leq \frac{1}{10C_Z (1 + |\theta|) \|a\|_{L^1}}, \quad (12)$$

and combining these with (9), we see

$$\|u_{m+1} - u_m\|_{Y_T \cap Z_T} \leq 4/5 \|u_m - u_{m-1}\|_{Y_T \cap Z_T},$$

which means that $\{u_m\}_m$ is a Cauchy sequence in $Y_T \cap Z_T$.

5 Proof of Theorem 1.3

We give the proof in the case (i), only. Let $w(t) = u(t) - v(t)$. By using estimates in the proof of Theorem 1.2, we can see that there exists small $t \in (0, T)$ so that

$$\|w\|_{Y_t} \leq 1/2 \|w\|_{Y_t}.$$

Let $T^* = \sup\{t \in (0, T); u(t, x) = v(t, x) \text{ a.e. } x \in \mathbb{R}^n\}$ and we assume that $T^* < T$, otherwise the proof is completed. Because $u(T^*) = v(T^*) \in L^\infty$, the above argument ensures that there exists $t_* > 0$ so that $u(t, x) = v(t, x)$ for $t \in [0, T^* + t_*]$ and a.e. $x \in \mathbb{R}^n$. This contradicts the definition of T^* and then we have $T^* = T$.

Acknowledgements

The research of this paper has been done during the stay of the second author at University of Bonn. The second author was supported by Research Fellowships of Japan Society for the Promotion of Science. The first and second authors were partially supported by the JSPS Strategic Young Researcher Overseas Visits Program for Accelerating Brain Circulation "Deepening and Evolution of Mathematics and Physics, Building of International Network Hub based on OCAMI (Osaka City University Advanced Mathematical Institute)". They also acknowledge the hospitality of University of Bonn. The authors are grateful to Professors J. Ahn and K. Kang for sending us their preprint.

References

- [1] J. Ahn and K. Kang, *On a Keller-Segel system with logarithmic sensitivity and non-diffusive chemical*, Preprint.
- [2] J.-M. Bony, *Calcul symbolique et propagation des singularites pour les equations aux derivees partielles non lineaires*, Ann. Sci. Ecole Norm. Sup. **14**, (4) (1981) 209-246.
- [3] H.Q. Bui, *Weighted Besov and Triebel spaces: Interpolation by the real method*, Hiroshima Math. J. **12** (1982), 581-605.
- [4] J.-Y. Chemin, *Perfect incompressible fluids*, Oxford Lecture Series in Mathematics and its Applications, **14**. The Clarendon Press, Oxford University Press, New York, (1998)
- [5] L. Corrias, B. Perthame and H. Zaag, *A chemotaxis model motivated by angiogenesis*, C.R. Acad. Sci. Paris, Ser. I **336** (2003), 141-146.
- [6] L. Corrias, B. Perthame and H. Zaag, *Global solutions of some chemotaxis and angiogenesis system in High space dimensions*, Milan J. Math. **72** (2004), 1-28.
- [7] K. Kang, A. Stevens and J.J.L. Velázquez, *Qualitative behavior of a Keller-Segel model with non-diffusive memory*, Comm. Partial Differential Equations, **35** (2010), no. 2, 245-274.
- [8] E.F. Keller and L.A. Segel, *Initiation of slime mold aggregation viewed as an instability*, J. Theor. Biol. **26** (1970), 399-416.
- [9] H. Kozono, T. Ogawa and Y. Taniuchi, *Navier-Stokes equations in the Besov space near L^∞ and BMO* Kyushu J. Math., **57** (2003), 303-324.
- [10] H.A. Levine and B.D. Sleeman, *A system of reaction diffusion equations arising in the theory of reinforced random walks*, SIAM J. Appl. Math. **57** (1997), no.3, 683-730.
- [11] D. Li, T. Li and K. Zhao, *On a hyperbolic-parabolic system modeling chemotaxis*, Math. Models Methods Appl. Sci. **21** (2011), no.8, 1631-1650.
- [12] T. Nagai, *Structure of solutions to a chemotaxis system in one space dimension*, Methods Appl. Anal. **12** (2005), no. 3, 221-238.
- [13] Y. Meyer, *Wavelet, Paraproduct and Navier-Stokes Equations*, Current developments in mathematics. International Press, (1996), 105-212.
- [14] H.G. Othmer and A. Stevens, *Aggregation, blow-up and collapse. The ABC's of taxis in reinforced random walks*, SIAM J. Appl. Math. **57** (1997), no.4, 1044-1081.
- [15] M. Rasle, *Sur une équation intégro-différentielle non linéaire issue de la biologie*, J. Differential Equations **32** (1979), no. 3, 420-453.
- [16] A. Stevens, *Trail following and aggregation of myxobacteria*, J. of Biological system **3** (1995), 1059-1068.

- [17] A. Stevens and J.J.L. Velázquez, *Asymptotic analysis of a chemotaxis system with non-diffusive memory*, Preprint.
- [18] H. Triebel, *Theory of function spaces*, Birkhäuser, Basel (1983).
- [19] H. Triebel, *Characterizations of Besov-Hardy-Sobolev spaces; A unified approach*, J. Approx. Theory **52** (1988), no.2, 162-203.
- [20] Y. Yang, H. Chen and W. Liu, *On existence of global solutions and blow-up to a system of a the reaction-diffusion equations modeling chemotaxis*, SIMA J. Math. Anal. **33** (2001), no.4, 763-785.