

HAMILTONIAN NON-DISPLACEABILITY OF GAUSS IMAGES OF ISOPARAMETRIC HYPERSURFACES

メタデータ	言語: English 出版者: OCAMI 公開日: 2019-09-20 キーワード (Ja): キーワード (En): isoparametric hypersurface, Gauss image, complex hyperquadric, Lagrangian submanifold, Hamiltonian non-displaceability, Floer homology 作成者: 入江, 博, Ma, Hui, 宮岡, 礼子, 大仁田, 義裕 メールアドレス: 所属: Ibaraki University, Tsinghua University, Tohoku University, Osaka City University
URL	https://ocu-omu.repo.nii.ac.jp/records/2016839

HAMILTONIAN NON-DISPLACEABILITY OF GAUSS IMAGES OF ISOPARAMETRIC HYPERSURFACES

HIROSHI IRIYEH, HUI MA, REIKO MIYAOKA, AND YOSHIHIRO
OHNITA

Citation	OCAMI Preprint Series
Issue Date	2015
Type	Preprint
Textversion	Author
Rights	For personal use only. No other uses without permission.
Relation	This is the pre-peer reviewed version of the following article: Bulletin of the London Mathematical Society, vol.48, issue 5, p802-812, which has been published in final form at https://doi.org/10.1112/blms/bdw040 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

From: Osaka City University Advanced Mathematical Institute

<http://www.sci.osaka-cu.ac.jp/OCAMI/publication/preprint/preprint.html>

HAMILTONIAN NON-DISPLACEABILITY OF GAUSS IMAGES OF ISOPARAMETRIC HYPERSURFACES

HIROSHI IRIYEH, HUI MA, REIKO MIYAOKA, AND YOSHIHIRO OHNITA

ABSTRACT. In this article we study the Hamiltonian non-displaceability of Gauss images of isoparametric hypersurfaces in the spheres as Lagrangian submanifolds embedded in complex hyperquadrics.

1. INTRODUCTION

The purpose of this article is to build a bridge between submanifold theory and symplectic geometry. Here, by submanifolds we mean the family of isoparametric hypersurfaces, originated from geometric optics in Italy, which is systematically developed by Élie Cartan. The latter is the Lagrangian intersection theory proposed by A. Weinstein and V.I. Arnold. Needless to say, A. Floer's contribution is quite important [10]. Recently, there has been intensive progress for the Lagrangian Floer theory of toric fibers. In order to proceed the study, we need, on one hand, a lot of concrete examples, and on the other hand, the common properties such examples share.

The isoparametric hypersurfaces are compact oriented hypersurfaces embedded in the standard sphere with constant principal curvatures. They provide a rich class containing infinitely many both homogeneous and non-homogeneous examples. Their topological and differential geometric data are well investigated, and manageable in various calculations. Münzner shows that the number g of distinct constant principal curvatures $\kappa_1 > \cdots > \kappa_g$ must be 1, 2, 3, 4 or 6, and the multiplicities m_i of κ_i are related by $m_i = m_{i+2} \pmod{g}$ [19]. Without loss of generality, we may assume $m_1 \leq m_2$. There are infinitely many non-homogeneous examples with $g = 4$, [24], [9]. The theory has already been applied to various mathematical problems (e.g., [28], [11], [27]).

Through the Gauss map \mathcal{G} , isoparametric hypersurfaces N^n in S^{n+1} yield a nice class of Lagrangian submanifolds $L^n = \mathcal{G}(N^n)$ embedded in the complex hyperquadric $Q_n(\mathbb{C})$ ($n \geq 2$), which is a rank two Hermitian symmetric space of compact type. B. Palmer [25] pointed out the minimality of L , and the second and the fourth authors discussed the Hamiltonian stability and related properties of L [15]–[18].

As a first step to obtain the Floer homology of L , which has been achieved in the cases $g = 1, 2$ [21], [14], we investigate the Hamiltonian non-displaceability of

2010 *Mathematics Subject Classification.* Primary 53C40; Secondary 53C42, 53D12.

Key words and phrases. isoparametric hypersurface, Gauss image, complex hyperquadric, Lagrangian submanifold, Hamiltonian non-displaceability, Floer homology.

The authors were partly supported by JSPS Grant-in-Aid for Young Scientists (B) No. 24740049, NSFC grant No. 11271213, JSPS Grant-in-Aid for Scientific Research (B) No. 15H03616, JSPS Grant-in-Aid for Scientific Research (S) No. 23224002 and (C) No. 15K04851, respectively.

L . Here, a compact Lagrangian submanifold L embedded in a symplectic manifold (M, ω) is said to be *Hamiltonian non-displaceable*, if $L \cap \varphi(L) \neq \emptyset$ for any φ belonging to the group $\text{Ham}(M, \omega)$ of all Hamiltonian diffeomorphisms of M .

Our main result is as follows:

Theorem 1.1. *Let N^n be a compact oriented isoparametric hypersurface of the standard sphere $S^{n+1}(1)$ in \mathbb{R}^{n+2} . Then its Gauss image $L^n = \mathcal{G}(N^n)$ is Hamiltonian non-displaceable in $(Q_n(\mathbb{C}), \omega_{\text{std}})$ except for the following few remaining cases:*

$$\begin{aligned} (g, n, m_1, m_2) &= (3, 3, 1, 1), & N &= \frac{SO(3)}{\mathbb{Z}_2 + \mathbb{Z}_2}, \\ (g, n, m_1, m_2) &= (4, 2k + 2, 1, k), & N &= \frac{SO(2) \times SO(k+2)}{\mathbb{Z}_2 \times SO(k)} \quad (k \geq 1), \\ (g, n, m_1, m_2) &= (6, 6, 1, 1), & N &= \frac{SO(4)}{\mathbb{Z}_2 + \mathbb{Z}_2}, \end{aligned}$$

where ω_{std} denotes the standard induced Kähler form of $Q_n(\mathbb{C}) \subset \mathbb{C}P^{n+1}$.

Here we should mention the preceding results and a remark. We know that any Gauss image $L = \mathcal{G}(N)$ is monotone. A compact monotone Lagrangian submanifold L in (M, ω) is said to be *wide* if the Floer homology $HF(L)$ satisfies

$$HF(L) \cong H_*(L; \mathbb{Z}_2) \otimes \Lambda,$$

and *narrow* if $HF(L) = 0$. See Sections 2 and 3 for details. Notice here if L is wide, then L is Hamiltonian non-displaceable. Y.G. Oh [21] showed that L is wide for any real form L of a compact irreducible Hermitian symmetric space M (see also [14]). In particular, all real forms of $Q_n(\mathbb{C})$, consisting of the Lagrangian sphere S^n and the real quadric $(S^k \times S^{n-k})/\mathbb{Z}_2$ ($1 \leq k \leq n-1$), are wide. Note that they coincide with the Gauss image of isoparametric hypersurfaces with $g = 1$ and 2 , respectively. Moreover, when $g = 3$ and $n = 6, 12$ or 24 , the Gauss image L is wide, although L is not a real form (Corollary 4.4). It is an interesting future problem to decide L is wide or not in the remaining cases.

The paper is organized as follows: In Section 2, we prepare fundamental properties of the Gauss images of isoparametric hypersurfaces in the standard sphere, especially the formula of their minimal Maslov number (Proposition 2.1). In Section 3, we recall Damian's lifted Floer homology and its spectral sequence [7], which is our main tool in the later sections. In Section 4, we examine the topology of Gauss images when $g = 3$. As a result we obtain that the Gauss image with $g = 3$ is a \mathbb{Z}_2 -homology sphere, and it is wide if $n = 6, 12$ or 24 . In Section 5 and Section 6, we prove the Hamiltonian non-displaceability of Gauss images with $g = 4$ or 6 and $m_1 \geq 2$, respectively. In the final section, we propose several open problems and a conjecture.

Throughout this article any manifold is smooth and connected.

Acknowledgments. This work was done mostly during the authors' short visits to RIMS in Kyoto, Japan in 2014 and 2015, and TSIMF in Sanya, China, in 2014. The authors would like to express their sincere gratitude to both institutions for the excellent research circumstance and generous support.

2. GAUSS IMAGES OF ISOPARAMETRIC HYPERSURFACES

Let N^n be an oriented hypersurface embedded in the unit standard sphere $S^{n+1}(1) \subset \mathbb{R}^{n+2}$. Denote by \mathbf{x} the position vector of points of N and \mathbf{n} the

unit normal vector field of N in $S^{n+1}(1)$. It is a fundamental fact in differential geometry that the *Gauss map* defined by

$$\mathcal{G} : N^n \ni p \longmapsto \mathbf{x}(p) \wedge \mathbf{n}(p) \cong [\mathbf{x}(p) + \sqrt{-1}\mathbf{n}(p)] \in \widetilde{Gr}_2(\mathbb{R}^{n+2}) \cong Q_n(\mathbb{C})$$

is always a Lagrangian immersion into the complex hyperquadric $Q_n(\mathbb{C})$. Here the complex hyperquadric $Q_n(\mathbb{C})$ is identified with the real Grassmann manifold $\widetilde{Gr}_2(\mathbb{R}^{n+2})$ of oriented 2-dimensional vector subspaces of \mathbb{R}^{n+2} , which has a symmetric space expression $SO(n+2)/(SO(2) \times SO(n))$.

It follows from [25] that the Gauss map $\mathcal{G} : N^n \rightarrow Q_n(\mathbb{C})$ from an isoparametric hypersurface N^n in $S^{n+1}(1)$ is a *minimal Lagrangian immersion* into $Q_n(\mathbb{C})$. Moreover, the ‘‘Gauss image’’ of \mathcal{G} is a compact minimal Lagrangian submanifold $L^n = \mathcal{G}(N^n) \cong N^n/\mathbb{Z}_g$ embedded in $Q_n(\mathbb{C})$, where $\mathcal{G} : N^n \rightarrow \mathcal{G}(N^n) = L^n$ is the covering map with the Deck transformation group \mathbb{Z}_g [15], [16], [23].

Let us briefly recall the notion of monotone Lagrangian submanifold. Given a Lagrangian manifold L in a symplectic manifold (M, ω) , two homomorphisms $I_{\mu, L} : \pi_2(M, L) \rightarrow \mathbb{Z}$ and $I_\omega : \pi_2(M, L) \rightarrow \mathbb{R}$ are defined as follows. For a smooth map $u : (D^2, \partial D^2) \rightarrow (M, L)$ in the class $A \in \pi_2(M, L)$, there is a unique trivialization, up to homotopy, of the pull-back bundle $u^*TM \cong D^2 \times \mathbb{C}^n$ as a symplectic bundle. This gives a map \tilde{u} from $S^1 = \partial D^2$ to $\Lambda(\mathbb{C}^n)$, the set of Lagrangian vector subspaces in \mathbb{C}^n . By using the Maslov class $\mu \in H^1(\Lambda(\mathbb{C}^n), \mathbb{Z})$, one can define $I_{\mu, L}(A) := \mu(\tilde{u})$. Next I_ω is defined by $I_\omega(A) := \int_{D^2} u^*\omega$. Then a Lagrangian submanifold L in M is said to be *monotone*, if there exists a constant $\lambda > 0$ such that

$$I_{\mu, L} = \lambda I_\omega.$$

Denote by $N_L \in \mathbb{Z}_+$ the positive generator of the image of $I_{\mu, L}$. We call N_L the *minimal Maslov number* of L .

We obtain the following properties of the Gauss image:

Proposition 2.1 ([23], [17]). *The Gauss image $L^n = \mathcal{G}(N^n)$ of an isoparametric hypersurface N^n in S^{n+1} is a compact monotone Lagrangian submanifold embedded in $Q_n(\mathbb{C})$, and its minimal Maslov number N_L is given by*

$$N_L = \frac{2n}{g} = \begin{cases} m_1 + m_2, & \text{if } g \text{ is even,} \\ 2m_1, & \text{if } g \text{ is odd.} \end{cases} \quad (2.1)$$

The Gauss image $\mathcal{G}(N^n)$ is orientable if and only if $2n/g$ is even.

Remark 2.1. It is well-known that $1 \leq N_L \leq 2n$ for a closed monotone Lagrangian submanifold L in $Q_n(\mathbb{C})$. The upper bound is sharp since the Lagrangian sphere $S^n \subset Q_n(\mathbb{C})$ attains $N_L = 2n$. The Gauss images of isoparametric hypersurfaces with $g = 1$ and $g = 2$ are S^n and $(S^k \times S^{n-k})/\mathbb{Z}_2$ ($1 \leq k \leq n-1$), respectively. They provide real forms of $Q_n(\mathbb{C})$, which are totally geodesic. Notice that the minimal Maslov number of $(S^k \times S^{n-k})/\mathbb{Z}_2$ ($1 \leq k \leq n-1$) is equal to n . We observe from (2.1) that Gauss images of isoparametric hypersurfaces with $g \geq 3$ provide examples of Lagrangian submanifolds in $Q_n(\mathbb{C})$ with relatively large minimal Maslov numbers (for the case of $\mathbb{C}P^n$, see [13]). On the other hand, $N_L = 2$ is taken only when $m_1 = m_2 = 1$, and such isoparametric hypersurfaces are homogeneous [6], [26], [8].

When $g \geq 3$, all Gauss images with $N_L = 2$ are given respectively [17] by

$$L = \begin{cases} \frac{SO(3)}{(\mathbb{Z}_2 + \mathbb{Z}_2)\mathbb{Z}_3}, & (g, m_1, m_2) = (3, 1, 1), \\ \frac{SO(2) \times SO(3)}{(\mathbb{Z}_2 \times 1)\mathbb{Z}_4}, & (g, m_1, m_2) = (4, 1, 1), \\ \frac{SO(4)}{(\mathbb{Z}_2 + \mathbb{Z}_2)\mathbb{Z}_6}, & (g, m_1, m_2) = (6, 1, 1). \end{cases}$$

3. DAMIAN'S LIFTED FLOER HOMOLOGY

In this section we recall necessary facts from the Floer theory for monotone Lagrangian submanifolds.

3.1. Lagrangian Floer homology. Let (M, ω) be a closed symplectic manifold and let $L \subset (M, \omega)$ be a closed connected monotone embedded Lagrangian submanifold with $N_L \geq 2$. Consider a Hamiltonian isotopy $\{\varphi_t\}_{0 \leq t \leq 1}$ defined by a time-dependent Hamiltonian $H : [0, 1] \times M \rightarrow \mathbb{R}$. The time-one map φ_1 is called a *Hamiltonian diffeomorphism* of M . Assume that $L \cap \varphi_1(L) \neq \emptyset$. The Floer complex $CF(L)$ is defined as a free module over \mathbb{Z}_2 generated by all points of the intersection $L \cap \varphi_1(L)$. Consider a time-dependent family $J = \{J_t\}_{0 \leq t \leq 1}$ of almost complex structures on M compatible with ω . Floer's boundary operator $\partial_J : CF(L) \rightarrow CF(L)$ is defined by counting (modulo 2) the isolated *J-holomorphic strips* $v : \mathbb{R} \times [0, 1] \rightarrow M$ connecting pairs of points of $L \cap \varphi_1(L)$ with boundary in $L \cup \varphi_1(L)$. For a generic choice of (H, J) , the homology $HF(L) := H_*(CF(L), \partial_J)$ is well-defined and called the *Floer homology* of L with \mathbb{Z}_2 -coefficient, which is invariant under the Hamiltonian isotopies of L (see [10], [20] and [22] for more details).

Let us regard the intersection $\mathcal{C} := L \cap \varphi_1(L)$ as points in L . For any two points $p, q \in \mathcal{C}$, consider an isolated *J-holomorphic strip* $v : \mathbb{R} \times [0, 1] \rightarrow M$ from p to q , which defines a path γ in L from p to q by $\gamma(s) := v(s, 0)$. We denote by Γ all the paths γ given by this procedure. Notice that to reconstruct the Floer complex $(CF(L), \partial_J)$ it is enough to know the above collection (\mathcal{C}, Γ) of points and paths.

3.2. Damian's spectral sequence. Now we recall the definition of Damian's lifted Floer homology for monotone Lagrangian submanifolds (see [7] for details). Let (M, ω) be a closed symplectic manifold and $L \subset M$ be a closed embedded monotone Lagrangian submanifold with $N_L \geq 3$. We start with the data (\mathcal{C}, Γ) and fix an arbitrary covering $\pi : \bar{L} \rightarrow L$. For any point $p \in \mathcal{C}$, denote by $\{p_i\}_{i \in I}$ the elements of the fiber $\pi^{-1}(p)$. Consider the set $\bar{\Gamma}$ of all the lifts of the paths of Γ to the covering space \bar{L} . For any points p_i, q_j ($i, j \in I$), the cardinality of elements in $\bar{\Gamma}$ which connect p_i with q_j is finite. We denote by $n(p_i, q_j)$ its parity. Let $CF^{\bar{L}}(L)$ be the free \mathbb{Z}_2 -module generated by $\bigcup_{p \in \mathcal{C}} \pi^{-1}(p)$. One can define the boundary operator $\partial^{\bar{L}}$ on $CF^{\bar{L}}(L)$ by the formula

$$\partial^{\bar{L}}(p_i) = \sum_{\pi(q_j)=q \in \mathcal{C}} n(p_i, q_j) q_j.$$

Damian [7, Proposition 2.6] proved that $(CF^{\bar{L}}(L), \partial^{\bar{L}})$ is a complex and its homology $HF^{\bar{L}}(L) := H_*(CF^{\bar{L}}(L), \partial^{\bar{L}})$ is called the *lifted Floer homology* of L , which is also invariant under the Hamiltonian isotopies of L .

The following spectral sequence is an adaptation of Biran's construction (see [2, Theorem 5.2.A]) to the lifted Floer homology, which is the main technical tool used in the sequel sections.

Theorem 3.1 (Damian's spectral sequence, [7, Theorem 2.9]). *Denote by $\Lambda = \mathbb{Z}_2[T, T^{-1}]$ the algebra of Laurent polynomials over \mathbb{Z}_2 and $\Lambda^i \subset \Lambda$ the subspace of homogeneous elements of degree i . There exists a spectral sequence $\{E_r^{p,q}, d_r\}$ which satisfies the following properties:*

- (1) $E_0^{p,q} = C_{p+q-pN_L}^{\bar{L}} \otimes \Lambda^{pN_L}$, $d_0 = [\partial_0^{\bar{L}}] \otimes 1$.
- (2) $E_1^{p,q} = H_{p+q-pN_L}(\bar{L}, \mathbb{Z}_2) \otimes \Lambda^{pN_L}$, $d_1 = [\partial_1^{\bar{L}}] \otimes T^{-N_L}$, where

$$[\partial_1^{\bar{L}}] : H_{p+q-pN_L}(\bar{L}; \mathbb{Z}_2) \rightarrow H_{p+q-1-(p-1)N_L}(\bar{L}; \mathbb{Z}_2)$$

is induced by $\partial_1^{\bar{L}}$.

- (3) For any $r \geq 1$, $E_r^{p,q}$ has the form $E_r^{p,q} = V_r^{p,q} \otimes \Lambda^{pN_L}$ with $d_r = \delta_r \otimes T^{-rN_L}$, where each $V_r^{p,q}$ is a vector space over \mathbb{Z}_2 and $\delta_r : V_r^{p,q} \rightarrow V_r^{p-r, q+r-1}$ is a homomorphism defined for every p, q and satisfies $\delta_r \circ \delta_r = 0$. Moreover,

$$V_{r+1}^{p,q} = \frac{\text{Ker}(\delta_r : V_r^{p,q} \rightarrow V_r^{p-r, q+r-1})}{\text{Im}(\delta_r : V_r^{p+r, q-r+1} \rightarrow V_r^{p,q})}.$$

In particular, $V_0^{p,q} = C_{p+q-pN_L}^{\bar{L}}$, $V_1^{p,q} = H_{p+q-pN_L}(\bar{L}; \mathbb{Z}_2)$, $\delta_1 = [\partial_1^{\bar{L}}]$.

- (4) $E_r^{p,q}$ collapses at $(\nu + 1)$ -step and for any $p \in \mathbb{Z}$, $\bigoplus_{q \in \mathbb{Z}} E_\infty^{p,q} \cong HF^{\bar{L}}(L)$, where $\nu = \lfloor \frac{\dim L + 1}{N_L} \rfloor$.

Back to the Gauss image, since $\nu = \lfloor \frac{\dim L + 1}{N_L} \rfloor = \lfloor \frac{(n+1)g}{2n} \rfloor$, one can easily get

Corollary 3.2. *For a Gauss image $L^n = \mathcal{G}(N^n) \subset Q_n(\mathbb{C})$, $g \geq 3$ and any $p, q \in \mathbb{Z}$, we have*

- (1) $E_2^{p,q} = E_\infty^{p,q}$ if and only if $g = 3$ and $(m_1, m_2) = (2, 2), (4, 4), (8, 8)$.
- (2) $E_3^{p,q} = E_\infty^{p,q}$ if and only if $g = 3$, $(m_1, m_2) = (1, 1)$ or $g = 4$.
- (3) $E_4^{p,q} = E_\infty^{p,q}$ if and only if $g = 6$, $(m_1, m_2) = (1, 1)$ or $(2, 2)$.

4. TOPOLOGY OF GAUSS IMAGES OF ISOPARAMETRIC HYPERSURFACES WITH $g = 3$

An isoparametric hypersurface with $g = 3$, the so called Cartan hypersurface, is given by a tube around the standard embedding of the projective plane $\mathbb{F}P^2$ in S^{3d+1} , where \mathbb{F} is the division algebra $\mathbb{R}, \mathbb{C}, \mathbb{H}$ or \mathbb{O} , and $d = 1, 2, 4$ or 8 [6]. The multiplicity satisfies $m = m_1 = m_2 = d$, and $n = 3m$. In this section, we first observe the topology of Gauss images of isoparametric hypersurfaces with $g = 3$, then apply Biran and Cornea's general fact to obtain their wideness except for the case $n = 3$.

Lemma 4.1. *The Gauss image L^3 of $g = 3$ is a \mathbb{Z}_2 -homology sphere.*

Proof. Recall that all homogeneous isoparametric hypersurfaces $N^n \subset S^{n+1}(1)$ can be obtained as principal orbits of compact Riemannian symmetric pairs (U, K) of rank 2, and that isoparametric hypersurfaces with $g = 3$ must be homogenous. For the case when $(g, m, n) = (3, 1, 3)$, in [15, p.780] we have a homogenous space expression $N = K/K_0 \cong \frac{SO(3)}{\mathbb{Z}_2 \oplus \mathbb{Z}_2} \cong SU(2)/\tilde{K}_0$, and $L = K/K_{[a]} \cong SU(2)/\tilde{K}_{[a]}$, where $\tilde{K}_{[a]}$ is generated by \tilde{K}_0 and

$$\pm \begin{pmatrix} \frac{1+\sqrt{-1}}{2} & \frac{1+\sqrt{-1}}{2} \\ -1+\sqrt{-1} & 1-\sqrt{-1} \end{pmatrix} =: \pm A_0, \quad \pm \begin{pmatrix} \frac{-1+\sqrt{-1}}{2} & \frac{1+\sqrt{-1}}{2} \\ -1+\sqrt{-1} & -1-\sqrt{-1} \end{pmatrix} =: \pm A_0^2,$$

with $A_0^3 = -\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. By a direct computation, we obtain $[\tilde{K}_{[a]}, \tilde{K}_{[a]}] = \tilde{K}_0$. Hence

$$H_1(L^3; \mathbb{Z}) \cong \pi_1(L^3)/[\pi_1(L^3), \pi_1(L^3)] \cong \tilde{K}_{[a]}/[\tilde{K}_{[a]}, \tilde{K}_{[a]}] \cong \tilde{K}_{[a]}/\tilde{K}_0 \cong \mathbb{Z}_3.$$

Since L^3 is orientable, by the Poincaré duality and universal coefficient theorem, we obtain

$$H_k(L^3; \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & \text{if } k = 0, 3, \\ \mathbb{Z}_3 & \text{if } k = 1, \\ 0 & \text{if } k = 2. \end{cases}$$

Therefore, by the universal coefficients theorem, L^3 is a \mathbb{Z}_2 -homology sphere. \square

Lemma 4.2. *The Gauss image $L^n = \mathcal{G}(N^n)$ of a Cartan isoparametric hypersurface $N^n \subset S^{n+1}(1)$ (i.e., $g = 3$ and $n = 3, 6, 12, 24$) is a \mathbb{Z}_2 -homology sphere.*

Proof. The \mathbb{Z}_2 -homology of N^n is known [19] as

$$H_k(N; \mathbb{Z}_2) \cong \begin{cases} \mathbb{Z}_2, & \text{for } k = 0, n, \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2, & \text{for } k = m, 2m, \\ 0, & \text{otherwise,} \end{cases}$$

where $n = 3m$ and $m \in \{1, 2, 4, 8\}$. Since $\mathcal{G} : N^n \rightarrow L^n$ is a covering map and \mathbb{Z}_3 acts on N^n as a Deck transformation, Theorem 2.4 in [5, p.120] implies

$$H_k(L^n; \mathbb{Z}_2) = H_k(N/\mathbb{Z}_3; \mathbb{Z}_2) \cong H_k(N; \mathbb{Z}_2)^{\mathbb{Z}_3} = 0,$$

for $k \neq 0, m, 2m, 3m (= n)$.

On the other hand, when $m \in \{2, 4, 8\}$, n is even and $\chi(L) = \chi(N^n)/3 = 6/3 = 2$ holds. Putting $l := \dim_{\mathbb{Z}_2} H_m(L; \mathbb{Z}_2)$, we obtain $2 = \chi(L) = 1 + l + l + 1 = 2 + 2l$, which yields $l = 0$. Hence L is a \mathbb{Z}_2 -homology sphere.

Combing with Lemma 4.1, we complete the proof. \square

Let us give several applications of Lemma 4.2. Firstly we recall Biran and Cornea's result [3] stated in [4, Proposition 1.6 (3)]:

Proposition 4.3. *Let L be a closed monotone Lagrangian submanifold embedded in a closed symplectic manifold (M, ω) . If $H_i(L; \mathbb{Z}_2) = 0$ for any $i \in \mathbb{Z}$ with $i \equiv -1 \pmod{N_L}$, then $HF(L) \cong H_*(L; \mathbb{Z}_2) \otimes \Lambda$.*

By Proposition 2.1, Lemma 4.2 and Proposition 4.3, we obtain

Corollary 4.4. *Let $L^n = \mathcal{G}(N^n) \subset Q_n(\mathbb{C})$ be the Gauss image of an isoparametric hypersurface N with $g = 3$ and $n = 6, 12, 24$. Then L is wide. Hence, $\#(L \cap \varphi(L)) \geq 2$ for any $\varphi \in \text{Ham}(Q_n(\mathbb{C}), \omega_{\text{std}})$ such that L intersects with $\varphi(L)$ transversally.*

Secondly, recall Biran and Khanevsky's result:

Proposition 4.5 ([4, Corollary 1.5]). *Let L^n be a closed monotone Lagrangian submanifold embedded in $Q_n(\mathbb{C})$ with $N_L \geq 2$ and $n \geq 2$. If $HF(L) \neq 0$, then $L \cap S^n \neq \emptyset$, where S^n denotes the totally geodesic Lagrangian sphere of $Q_n(\mathbb{C})$.*

Combining Corollary 4.4 with Proposition 4.5, we obtain

Corollary 4.6. *Let $L^n = \mathcal{G}(N^n) \subset Q_n(\mathbb{C})$ be as in Corollary 4.4. Then $\varphi(L) \cap S^n \neq \emptyset$ for any $\varphi \in \text{Ham}(Q_n(\mathbb{C}), \omega_{\text{std}})$.*

For any n -dimensional submanifold R in $Q_n(\mathbb{C})$, it holds

$$\int_{SO(n+2)} \#(\phi(R) \cap S^n) d\mu_{SO(n+2)}(\phi) \leq 2 \frac{\text{Vol}(SO(n+2))}{\text{Vol}(S^n)} \text{Vol}(R),$$

where $\phi \in SO(n+2) \subset \text{Ham}(Q_n(\mathbb{C}), \omega_{\text{std}})$ ((6.7) in [14]). Finally, we give an estimate of the volume of Gauss images deformed by Hamiltonian isotopies of $Q_n(\mathbb{C})$. Namely, applying above to $R = \varphi(L)$, we obtain

Corollary 4.7. *For a Lagrangian submanifold L in Corollary 4.4,*

$$\frac{1}{2} \text{Vol}(S^n) \leq \text{Vol}(\varphi(L))$$

holds for any $\varphi \in \text{Ham}(Q_n(\mathbb{C}), \omega_{\text{std}})$.

- Remark 4.1.** (1) From Gorodski and Podesta's classification [12] of all compact tight \mathbb{Z}_2 -homology Lagrangian spheres in simply connected compact homogeneous Kähler manifolds, we know that any compact tight \mathbb{Z}_2 -homology Lagrangian sphere in $Q_n(\mathbb{C})$ is the standard real form S^n up to congruence.
- (2) Biran and Cornea [3] proved that any even dimensional closed Lagrangian submanifold L embedded in $Q_n(\mathbb{C})$ with $H_1(L; \mathbb{Z}) = 0$ must be a \mathbb{Z}_2 -homology sphere. Notice that the Gauss images L with $g = 3$ are \mathbb{Z}_2 -homology spheres with $H_1(L; \mathbb{Z}) \cong \mathbb{Z}_3$.

5. HAMILTONIAN NON-DISPLACEABILITY OF GAUSS IMAGES OF ISOPARAMETRIC HYPERSURFACES WITH $g = 4$

Now we consider the Gauss images L of isoparametric hypersurfaces with $g = 4$. Since $N_L = \frac{2n}{4} = \frac{n}{2}$, $N_L \geq 3$ leads to $n \geq 6$. Thus except for the only one case $n = 4$ when $L = \frac{SO(2) \times SO(3)}{(\mathbb{Z}_2 \times 1)\mathbb{Z}_4}$, the argument in Subsection 3.2 can work and the lifted Floer homology $HF^{\bar{L}}(L)$ is well-defined for any covering \bar{L} of L .

Proposition 5.1. *Let $L = \mathcal{G}(N) \subset Q_n(\mathbb{C})$ be the Gauss image of an isoparametric hypersurface N with $g = 4$ and $m_1 \geq 2$. Then $HF^{\bar{L}}(L) \neq 0$ for $\bar{L} = N$.*

Proof. Consider the \mathbb{Z}_4 -covering $\bar{L} = N^n$ over L^n . We argue by contradiction. Assume that $HF^{\bar{L}}(L) = 0$. Since $\nu = \lfloor \frac{\dim L + 1}{N_L} \rfloor = 2$, Damian's spectral sequence $\{E_r^{p,q}, d_r\}$ collapses at 3-step. Then

$$0 = E_3^{0,q} = V_3^{0,q} = \frac{\text{Ker}(\delta_2 : V_2^{0,q} \rightarrow V_2^{-2,q+1})}{\text{Im}(\delta_2 : V_2^{2,q-1} \rightarrow V_2^{0,q})}$$

and thus we have the following exact sequence

$$V_2^{2,q-1} \rightarrow V_2^{0,q} \rightarrow V_2^{-2,q+1} \quad (5.1)$$

for any $q \in \mathbb{Z}$. Since

$$\begin{aligned} V_2^{2,q-1} &= \frac{\text{Ker}([\partial_1^{\bar{L}}] : H_{q+1-2N_L}(\bar{L}; \mathbb{Z}_2) \rightarrow H_{q-N_L}(\bar{L}; \mathbb{Z}_2))}{\text{Im}([\partial_1^{\bar{L}}] : H_{q+2-3N_L}(\bar{L}; \mathbb{Z}_2) \rightarrow H_{q+1-2N_L}(\bar{L}; \mathbb{Z}_2))}, \\ V_2^{-2,q+1} &= \frac{\text{Ker}([\partial_1^{\bar{L}}] : H_{q-1+2N_L}(\bar{L}; \mathbb{Z}_2) \rightarrow H_{q-2+3N_L}(\bar{L}; \mathbb{Z}_2))}{\text{Im}([\partial_1^{\bar{L}}] : H_{q+N_L}(\bar{L}; \mathbb{Z}_2) \rightarrow H_{q-1+2N_L}(\bar{L}; \mathbb{Z}_2))}, \end{aligned}$$

we see that $V_2^{2,q-1} = V_2^{-2,q+1} = 0$ when $2 \leq q \leq n-2$. Then it follows from (5.1) that

$$0 = V_2^{0,q} = \frac{\text{Ker}([\partial_1^{\bar{L}}] : H_q(\bar{L}; \mathbb{Z}_2) \rightarrow H_{q-1+N_L}(\bar{L}; \mathbb{Z}_2))}{\text{Im}([\partial_1^{\bar{L}}] : H_{q+1-N_L}(\bar{L}; \mathbb{Z}_2) \rightarrow H_q(\bar{L}; \mathbb{Z}_2))}$$

holds for $2 \leq q \leq n-2$.

Putting $q = N_L = m_1 + m_2$, we have the exact sequence

$$H_1(\bar{L}; \mathbb{Z}_2) \rightarrow H_{m_1+m_2}(\bar{L}; \mathbb{Z}_2) \rightarrow H_{2(m_1+m_2)-1}(\bar{L}; \mathbb{Z}_2). \quad (5.2)$$

Recall that by Münzner's result [19] the \mathbb{Z}_2 -homology of N^n is given by

$$H_k(N; \mathbb{Z}_2) \cong \begin{cases} \mathbb{Z}_2, & \text{for } k = 0, m_1, m_2, 2m_1 + m_2, m_1 + 2m_2, n, \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2, & \text{for } k = m_1 + m_2, \\ 0, & \text{otherwise.} \end{cases}$$

By the assumption that $m_1 \geq 2$, (5.2) leads to the exact sequence $0 \rightarrow \mathbb{Z}_2 \oplus \mathbb{Z}_2 \rightarrow 0$, which is a contradiction. \square

Corollary 5.2. *Under the same assumption as in Proposition 5.1, the Gauss image $L^n \subset (Q_n(\mathbb{C}), \omega_{\text{std}})$ is Hamiltonian non-displaceable.*

Remark 5.1. Note that any non-homogeneous isoparametric hypersurface in the standard sphere satisfies the assumption of Proposition 5.1. By R. Takagi's result [26] we know that an isoparametric hypersurface N^n with $g = 4$ and $m_1 = 1$ must be $N^n = \frac{SO(2) \times SO(k+2)}{\mathbb{Z}_2 \times SO(k)}$ ($n = 2k + 2, m_2 = k$), in particular homogeneous.

6. HAMILTONIAN NON-DISPLACEABILITY OF THE GAUSS IMAGE OF AN ISOPARAMETRIC HYPERSURFACE WITH $g = 6$

When $g = 6$, $N_L \geq 3$ implies $m_1 = m_2 = 2$ [1]. Thus we focus on the Gauss image $L^{12} = \mathcal{G}(N^{12})$ of the isoparametric hypersurface $N^{12} \subset S^{13}(1)$ with $g = 6$ and $m_1 = m_2 = 2$. Compared with the case when $g = 4$, this case is more complicated because $\{E_r^{p,q}, d_r\}$ collapses at 4-step.

Proposition 6.1. *Let $L^{12} = \mathcal{G}(N^{12}) \subset Q_{12}(\mathbb{C})$ be the Gauss image of an isoparametric hypersurface with $g = 6$ and $m_1 = m_2 = 2$. Then $HF^{\bar{L}}(L) \neq 0$ for $\bar{L} = N$. In particular, L is Hamiltonian non-displaceable in $Q_{12}(\mathbb{C})$.*

Proof. Take the \mathbb{Z}_6 -covering $\bar{L} = N^{12}$ over L^{12} . Since $\nu = \lceil \frac{12+1}{4} \rceil = 3$, Damian's spectral sequence $\{E_r^{p,q}, d_r\}$ collapses at 4-step. Assume that $HF^{\bar{L}}(L) = 0$. Then we have

$$0 = E_4^{0,q} = V_4^{0,q} = \frac{\text{Ker}(\delta_3 : V_3^{0,q} \rightarrow V_3^{-3,q+2})}{\text{Im}(\delta_3 : V_3^{3,q-2} \rightarrow V_3^{0,q})}$$

and the following exact sequence

$$V_3^{3,q-2} \rightarrow V_3^{0,q} \rightarrow V_3^{-3,q+2} \quad (6.1)$$

for any $q \in \mathbb{Z}$.

Notice that

$$\begin{aligned} V_3^{3,q-2} &= \frac{\text{Ker}(\delta_2 : V_2^{3,q-2} \rightarrow V_2^{1,q-1})}{\text{Im}(\delta_2 : V_2^{5,q-3} \rightarrow V_2^{3,q-2})}, \\ V_2^{3,q-2} &= \frac{\text{Ker}([\partial_1^{\bar{L}}] : H_{q-11}(\bar{L}; \mathbb{Z}_2) \rightarrow H_{q-8}(\bar{L}; \mathbb{Z}_2))}{\text{Im}([\partial_1^{\bar{L}}] : H_{q-14}(\bar{L}; \mathbb{Z}_2) \rightarrow H_{q-11}(\bar{L}; \mathbb{Z}_2))}. \end{aligned}$$

Consequently, $V_3^{3,q-2} = 0$ when $q \leq 10$. Similarly, from

$$\begin{aligned} V_3^{-3,q+2} &= \frac{\text{Ker}(\delta_2 : V_2^{-3,q+2} \rightarrow V_2^{-5,q+3})}{\text{Im}(\delta_2 : V_2^{-1,q+1} \rightarrow V_2^{-3,q+2})}, \\ V_2^{-3,q+2} &= \frac{\text{Ker}([\partial_1^{\bar{L}}] : H_{q+11}(\bar{L}; \mathbb{Z}_2) \rightarrow H_{q+14}(\bar{L}; \mathbb{Z}_2))}{\text{Im}([\partial_1^{\bar{L}}] : H_{q+8}(\bar{L}; \mathbb{Z}_2) \rightarrow H_{q+11}(\bar{L}; \mathbb{Z}_2))}, \end{aligned}$$

we see that $V_3^{-3,q+2} = 0$ if $q \geq 2$. Thus (6.1) leads to $V_3^{0,q} = 0$ for $2 \leq q \leq 10$. Due to

$$0 = V_3^{0,q} = \frac{\text{Ker}(\delta_2 : V_2^{0,q} \rightarrow V_2^{-2,q+1})}{\text{Im}(\delta_2 : V_2^{2,q-1} \rightarrow V_2^{0,q})},$$

we get the exact sequence $V_2^{2,q-1} \rightarrow V_2^{0,q} \rightarrow V_2^{-2,q+1}$ for any $2 \leq q \leq 10$. Thanks to

$$\begin{aligned} V_2^{2,q-1} &= \frac{\text{Ker}([\partial_1^{\bar{L}}] : H_{q-7}(\bar{L}; \mathbb{Z}_2) \rightarrow H_{q-4}(\bar{L}; \mathbb{Z}_2))}{\text{Im}([\partial_1^{\bar{L}}] : H_{q-10}(\bar{L}; \mathbb{Z}_2) \rightarrow H_{q-7}(\bar{L}; \mathbb{Z}_2))}, \\ V_2^{-2,q+1} &= \frac{\text{Ker}([\partial_1^{\bar{L}}] : H_{q+7}(\bar{L}; \mathbb{Z}_2) \rightarrow H_{q+10}(\bar{L}; \mathbb{Z}_2))}{\text{Im}([\partial_1^{\bar{L}}] : H_{q+4}(\bar{L}; \mathbb{Z}_2) \rightarrow H_{q+7}(\bar{L}; \mathbb{Z}_2))}, \end{aligned}$$

we derive $V_2^{0,6} = 0$. Now because

$$0 = V_2^{0,6} = \frac{\text{Ker}([\partial_1^{\bar{L}}] : H_6(\bar{L}; \mathbb{Z}_2) \rightarrow H_9(\bar{L}; \mathbb{Z}_2))}{\text{Im}([\partial_1^{\bar{L}}] : H_3(\bar{L}; \mathbb{Z}_2) \rightarrow H_6(\bar{L}; \mathbb{Z}_2))},$$

we get the exact sequence $H_3(\bar{L}; \mathbb{Z}_2) \rightarrow H_6(\bar{L}; \mathbb{Z}_2) \rightarrow H_9(\bar{L}; \mathbb{Z}_2)$, which contradicts the facts in [19] that $H_3(\bar{L}; \mathbb{Z}_2) \cong H_9(\bar{L}; \mathbb{Z}_2) = 0$ and $H_6(\bar{L}; \mathbb{Z}_2) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$. \square

Therefore we complete the proof of Theorem 1.1.

7. OPEN PROBLEMS AND A CONJECTURE

The following are open problems for further research:

- (1) We cannot apply the same method above to determine the Hamiltonian non-displaceability of Gauss images of the following isoparametric hypersurfaces:
 - (i) $g = 3$, $(m_1, m_2) = (1, 1)$, $n = 3$;
 - (ii) $g = 4$, $(m_1, m_2) = (1, k)$, $n = 2(1 + k)$;
 - (iii) $g = 6$, $(m_1, m_2) = (1, 1)$, $n = 6$.
Consider how we can do in these cases.
- (2) Determine the Floer homology $HF(L)$ of Gauss images for $(g, n) = (3, 3)$ and $g = 4, 6$.
- (3) Describe $H_*(L; \mathbb{Z}_2)$ of the Gauss images $L = N/\mathbb{Z}_g$ for $g = 4, 6$ explicitly.

According to the present work and a private communication with Hajime Ono, he and the authors give

Conjecture. Any compact connected minimal Lagrangian submanifold in an irreducible Hermitian symmetric space of compact type is Hamiltonian non-displaceable.

REFERENCES

- [1] U. Abresch, *Isoparametric hypersurfaces with four or six distinct principal curvatures. Necessary conditions on the multiplicities*. Math. Ann. **264** (1983), 283–302.
- [2] P. Biran, *Lagrangian non-intersections*, Geom. Funct. Anal. **16** (2006), 279–326.
- [3] P. Biran and O. Cornea, *Rigidity and uniruling for Lagrangian submanifolds*, Geom. Topol. **13** (2009), 2881–2989.
- [4] P. Biran and M. Khanevsky, *A Floer-Gysin exact sequence for Lagrangian submanifolds*, Comment. Math. Helv. **88** (2013), 899–952.
- [5] G.E. Bredon, *Introduction to Compact Transformation Groups*. Pure and Applied Mathematics, Vol. 46. Academic Press, New York-London, 1972.
- [6] É. Cartan, *Sur des familles remarquables d’ hypersurfaces isoparamétriques dans les espaces sphériques*, Math. Z. **45** (1939), 335–367.
- [7] M. Damian, *Floer homology on the universal cover, Audin’s conjecture and other constraints on Lagrangian submanifolds*, Comment. Math. Helv. **87** (2012), 433–462.
- [8] J. Dorfmeister and E. Neher, *Isoparametric hypersurfaces, case $g = 6$, $m = 1$* , Comm. Algebra **13** (1985), 2299–2368.
- [9] D. Ferus, H. Karcher and H. F. Münzner, *Cliffordalgebren und neue isoparametrische Hyperflächen*, Math. Z. **177** (1981), 479–502.
- [10] A. Floer, *Morse theory for Lagrangian intersections*, J. Differential Geom. **28** (1988), 513–547.
- [11] J.Q. Ge and Y.Q. Xie, *Gradient map of isoparametric polynomial and its application to Ginzburg-Landau system*, J. Func. Anal., **258** (2010), 1682–1691.
- [12] C. Gorodski and F. Podestà, *Tight Lagrangian homology spheres in compact homogeneous Kähler manifolds*, Israel J. Math. **206** (2015), no. 1, 413–429.
- [13] H. Iriyeh, *Symplectic topology of Lagrangian submanifolds of CP^n with intermediate minimal Maslov numbers*, arXiv:1401.0777v1, 2014.
- [14] H. Iriyeh, T. Sakai and H. Tasaki, *Lagrangian Floer homology of a pair of real forms in Hermitian symmetric spaces of compact type*, J. Math. Soc. Japan **65** (2013), 1135–1151.
- [15] H. Ma and Y. Ohnita, *On Lagrangian submanifolds in complex hyperquadrics and isoparametric hypersurfaces in spheres*, Math. Z. **261** (2009), 749–785.
- [16] H. Ma and Y. Ohnita, *Differential geometry of Lagrangian submanifolds and Hamiltonian variational problems*, Harmonic maps and differential geometry, Contemp. Math., **542**, 115–134 Amer. Math. Soc., Providence, RI, 2011.
- [17] H. Ma and Y. Ohnita, *Hamiltonian stability of the Gauss images of homogeneous isoparametric hypersurfaces. I*, J. Differential Geom. **97** (2014), 275–348.
- [18] H. Ma and Y. Ohnita, *Hamiltonian stability of the Gauss images of homogeneous isoparametric hypersurfaces. II*, Tohoku Math. J. **67** (2015), 195–246
- [19] H. F. Münzner, *Isoparametrische Hyperflächen in Sphären, and II*, Math. Ann. **251** (1980), 57–71, **256** (1981), 215–232.
- [20] Y.-G. Oh, *Floer cohomology of Lagrangian intersections and pseudo-holomorphic discs I*, Comm. Pure Appl. Math. **46** (1993), 949–994; Addendum **48** (1995) 1299–1302.
- [21] Y.-G. Oh, *Floer cohomology of Lagrangian intersections and pseudo-holomorphic discs III*, H. Hofer, et al. (Eds.), The Floer Memorial Volume, Birkhäuser, Basel, 1995, 555–573.
- [22] Y.-G. Oh, *Floer cohomology, spectral sequences, and the Maslov class of Lagrangian embeddings*, Int. Math. Res. Not. **7** (1996), 305–346.
- [23] Y. Ohnita, *Geometry of Lagrangian submanifolds and isoparametric hypersurfaces*, Proceedings of The Fourteenth International Workshop on Differential Geometry and Related Fields, **14** (2010), 43–67, ed.by Y.-J. Suh, NIMS, KMS and GRG.
- [24] H. Ozeki and M. Takeuchi, *On some types of isoparametric hypersurfaces in spheres I and II*, Tohoku Math. J. **27** (1975), 515–559, **28** (1976), 7–55.
- [25] B. Palmer, *Hamiltonian minimality and Hamiltonian stability of Gauss maps*, Diff. Geom. and its Appl. **7** (1997), 51–58.
- [26] R. Takagi, *A class of hypersurfaces with constant principal curvatures in a sphere*, J. Differential Geom. **11** (1976), 225–233.
- [27] Z.Z. Tang and W.J. Yan, *Isoparametric foliation and Yau conjecture on the first eigenvalue*, J. Differential Geom. **94** (2013), 521–540.

- [28] Q. M. Wang, *On the topology of Clifford isoparametric hypersurfaces*, J. Differential Geom., **27** (1988), 55–66.

MATHEMATICS AND INFOMATICS, COLLEGE OF SCIENCE, IBARAKI UNIVERSITY, MITO, 310-8512, JAPAN.

E-mail address: `hiroshi.irie.math@vc.ibaraki.ac.jp`

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084, P.R. CHINA

E-mail address: `hma@math.tsinghua.edu.cn`

MATHEMATICAL INSTITUTE, TOHOKU UNIVERSITY, SENDAI 980-8578, JAPAN

E-mail address: `r-miyaok@m.tohoku.ac.jp`

OSAKA CITY UNIVERSITY ADVANCED MATHEMATICAL INSTITUTE & DEPARTMENT OF MATHEMATICS, OSAKA CITY UNIVERSITY, OSAKA, 558-8585, JAPAN

E-mail address: `ohnita@sci.osaka-cu.ac.jp`