

BAND SURGERY ON KNOTS AND LINKS, III

メタデータ	言語: English 出版者: OCAMI 公開日: 2019-09-20 キーワード (Ja): キーワード (En): Band surgery, coherent band-Gordian distance, H(2)-unknotting number, H(2)-Gordian distance 作成者: 金信, 泰造 メールアドレス: 所属: Osaka City University
URL	https://ocu-omu.repo.nii.ac.jp/records/2016840

BAND SURGERY ON KNOTS AND LINKS, III

TAIZO KANENOBU

Citation	OCAMI Preprint Series
Issue Date	2015
Type	Preprint
Textversion	Author
Rights	© 2016 World Scientific Publishing Company. For personal use only. No other uses without permission.
Relation	Preprint of an article submitted for consideration in Journal of Knot Theory and Its Ramifications © 2016 World Scientific Publishing Company. https://www.worldscientific.com/worldscinet/jktr .
Is version of	https://doi.org/10.1142/S0218216516500565

From: Osaka City University Advanced Mathematical Institute

<http://www.sci.osaka-cu.ac.jp/OCAMI/publication/preprint/preprint.html>

BAND SURGERY ON KNOTS AND LINKS, III

TAIZO KANENOBU

ABSTRACT. We give two criteria of links concerning a band surgery: The first one is a condition on the determinants of links which are related by a band surgery using Nakanishi's criterion on knots with Gordian distance one. The second one is a criterion on knots with $H(2)$ -Gordian distance two by using a special value of the Jones polynomial, where an $H(2)$ -move is a band surgery preserving a component number. Then, we give an improved table of $H(2)$ -Gordian distances between knots with up to seven crossings, where we add Zeković's result.

1. INTRODUCTION

Let L be a link in S^3 and $b : I \times I \rightarrow S^3$ an embedding such that $L \cup b(I \times I) = b(I \times \partial I)$, where I is the unit interval $[0, 1]$. Then we may obtain a new link $M = (L \setminus b(I \times \partial I)) \cup b(\partial I \times I)$, which is called a link obtained from L by the *band surgery* along the band B , where $B = b(I \times I)$; see Fig. 1. If L and M are oriented links, and a band surgery preserves the orientations of L and M , the band surgery is said to be *coherent*, otherwise *incoherent*. If L and M are unoriented links, and a band surgery preserves the number of components, then it is called the $H(2)$ -*move*.

Since any oriented link can be deformed into the trivial knot by a sequence of coherent band surgeries, we define the *coherent band-Gordian distance* between two oriented links L and M to be the least number of coherent band surgeries needed to deform L into M , which we denote by $d_{\text{cb}}(L, M)$. Similarly, since any knot can be deformed into the trivial knot by a sequence of $H(2)$ -moves, we define the $H(2)$ -*Gordian distance* between two knots J and K , which we denote by $d_2(J, K)$. In particular, the $H(2)$ -*unknotting number* of a knot K , $u_2(K)$, is the $H(2)$ -Gordian distance between K and the trivial knot.

In this paper we give two criteria of links concerning a band surgery: The first one is a condition on the determinant of a knot or link which is obtained from an unknotting number one knot by a band surgery (Theorem 2.2). This is easily obtained by using a condition on the determinant of a knot obtained from an unknotting number one knot by a crossing change due to Nakanishi [10, 11] (Proposition 2.1). The idea of the proof is similar to that of Theorem 4.2 in [8], which gives a condition on the determinant of a link or knot obtained from a 2-bridge knot by a band surgery. This uses a condition on the determinant

Date: February 10, 2016.

1991 Mathematics Subject Classification. Primary 57M25; Secondary 57M27.

Key words and phrases. Band surgery, coherent band-Gordian distance, $H(2)$ -unknotting number, $H(2)$ -Gordian distance.




FIGURE 1. The link M is obtained from L by a band surgery along the band B , and vice versa.

of a knot obtained from a 2-bridge knot by a crossing change due to Murakami [9]. Using Theorem 2.2 we give tables of the values for which the determinant of a link L does not take such that either $d_2(K, L) = 1$ or $d_{cb}(K, L) = 1$, where K is an unknotting number one knot with determinant ≤ 115 (Tables 1 and 2). They yield a table of pairs of an unknotting number one knot J and a knot K with $d_2(J, K) > 1$ (Table 3), and a table of pairs of an unknotting number one knot J and a 2-component link L with $d_{cb}(J, L) > 1$, where the crossing numbers of J , K and L are ≤ 8 (Table 4). As corollaries of Theorem 2.2, we obtain a condition for an unknotting number one knot to have $H(2)$ -unknotting number two (Corollaries 2.4 and 2.5).

The second one is a criterion on knots with $H(2)$ -Gordian distance two by using a special value of the Jones polynomial (Theorem 3.1), which extends some criteria given in [6, 8].

As an application, we give tables of $H(2)$ -Gordian distances between knots with up to 7 crossings (Tables 6 and 7). They improve the tables compiled in [6], where there remain 60 pairs of knots whose $H(2)$ -Gordian distances are unsettled. Among these pairs we decide the $H(2)$ -Gordian distances for 20 pairs of knots using the criteria above together with those given in [8]. Further, we can decide for 8 pairs of knots by virtue of the paper of Zeković [14]. She has given a new method for searching pairs of knots related by either a crossing change or an $H(2)$ -move. Then she gave tables of pairs of knots with Gordian distance one and those with $H(2)$ -Gordian distance one with at most 9 crossings.

This paper is organized as follows: In Sec. 2 we give a condition on the determinant of a knot or link which is obtained from an unknotting number one knot by a band surgery, which is deduced from Nakanishi's criterion. Then we give a condition for an unknotting number one knot to have $H(2)$ -unknotting number two in terms of the determinant of a knot. In Sec. 3 we give a criterion of a pair of knots with $H(2)$ -Gordian distance two using the special value of the Jones polynomial. In Sec. 4 we give the tables of the $H(2)$ -Gordian distances between knots with up to seven crossings, which improves those in [6].

Notation. For knots and links we use Rolfsen notations [12, Appendix C]. For a knot or link L , we denote by $L!$ its mirror image.

2. DETERMINANT OF A LINK OBTAINED FROM AN UNKNOTTING NUMBER ONE KNOT BY A BAND SURGERY

The following criterion is due to Nakanishi, which is implied from Proposition 13 in [11] and has been essentially given in Theorem 3 in [10].

Proposition 2.1. *Let K be an unknotting number one knot. If a knot J is obtained from K by a crossing change, then there exists an integer s such that:*

$$(1) \quad \det J \equiv \pm s^2 \pmod{\det K}.$$

Using this proposition, we may deduce the following.

Theorem 2.2. *Suppose that a knot or link L is obtained from an unknotting number one knot K by a coherent or incoherent band surgery. Then there exists an integer s such that:*

$$(2) \quad 2 \det L \equiv \pm s^2 \pmod{\det K}.$$

The proof is similar to that of Theorem 4.2 in [8]. In order to prove this, we use the Jones polynomial [5]. We define the *Jones polynomial* $V(L; t) \in \mathbf{Z}[t^{\pm 1/2}]$ of an oriented link L by the following formulas:

$$(3) \quad V(U; t) = 1;$$

$$(4) \quad t^{-1}V(L_+; t) - tV(L_-; t) = (t^{1/2} - t^{-1/2})V(L_0; t);$$

where U is the unknot and L_+ , L_- , L_0 are three oriented links that are identical except near one point where they are as in Fig. 2; we call an ordered set (L_+, L_-, L_0) a *skein triple*.



FIGURE 2. A skein triple (L_+, L_-, L_0) , and a knot L_∞ .

If L_+ and L_- are knots, then L_0 is a 2-component link and we may consider another knot L_∞ which is of the diagram of Fig. 2. Then L_+/L_- and L_∞ are related by an incoherent band surgery, and we have the following relation [3, Theorem 2]:

$$(5) \quad V(L_+; t) - tV(L_-; t) + t^{3\lambda}(t-1)V(L_\infty; t) = 0,$$

where λ is the linking number of L_0 .

For a c -component link L , $i^{c-1}V(L; -1)$ is an integer and the determinant $\det L$ is given by $\det L = |V(L; -1)|$. Putting $t = -1$ in Eqs. (4) and (5), we obtain

$$(6) \quad -V(L_+; -1) + V(L_-; -1) = 2iV(L_0; -1);$$

$$(7) \quad V(L_+; -1) + V(L_-; -1) = 2(-1)^\lambda V(L_\infty; -1).$$

Proof of Theorem 2.2. Suppose that a 2-component link L is obtained from an unknotting number one knot J by a coherent band surgery. Then there exists a knot K such that (J, K, L) is a skein triple; cf. Lemma 2.1(i) in [8]. Then by Proposition 2.1 there exists an integer s with Eq. (1). From Eq. (6) we have $-V(J; -1) + V(K; -1) = 2iV(L; -1)$, and so

since $\det J = |V(J; -1)|$ and $\det K = |V(K; -1)|$, we obtain Eq. (2). For the case where L and J are related by an incoherent band surgery we use Eq. (7). \square

For an unknotting number one knot K with $\det K \leq 115$ we list the impossible values of $\det L$, where L is a knot or link with $d_2(K, L) = 1$ or $d_{cb}(K, L) = 1$ in Tables 1 and 2. They are obtained by using the Mathematica program. Notice that if K is a prime knot with up to 10 crossings, then $\det K \leq 111$. Therefore, Tables 1 and 2 yield Tables 3 and 4. In Table 3 the symbol \times means that the unknotting number one knots J in the row and the knots K in the column are not related by an incoherent band surgery, which also implies that the pairs $(J!, K!)$, $(J, K!)$, $(J!, K)$ are not related by an incoherent band surgery. For example, the pairs of knots $(6_1, 7_4)$, $(6_1!, 7_4!)$, $(6_1, 7_4!)$, $(6_1!, 7_4)$ are not related by an incoherent band surgery. In Table 4 the symbol \times means that the unknotting number one knots J in the row and the oriented links L in the column with any orientation are not related by a coherent band surgery, which implies that the pairs of oriented knots and links (J, L) , $(J!, L!)$, $(J, L!)$, $(J!, L)$ with any orientation are not related by a coherent band surgery.

Example 2.3. $d_2(5_2, 8_{17}) = d_2(7_1, 8_{17}) = 2$. Note that 8_{17} is negative-amphicheiral. First, by Table 3 we see $d_2(5_2, 8_{17}) > 1$ and $d_2(7_1, 8_{17}) > 1$. Notice that previously known methods in [6] (Theorems 4.1, 5.5, 8.1) and [8] (Theorems 4.2, 5.2(iii), 7.2) do not prove this; see Table 5. Conversely, we have

$$(8) \quad d_2(5_2, 8_{17}) \leq d_2(5_2, 6_3) + d_2(6_3, 8_{17}) = 2;$$

$$(9) \quad d_2(7_1, 8_{17}) \leq d_2(7_1, 4_1) + d_2(4_1, 8_{17}) = 2.$$

In fact, the knot 8_{17} is transformed into 4_1 and 6_3 by the $H(2)$ -moves using the band and the crossing as shown in Fig. 3, respectively, which imply $d_2(8_{17}, 4_1) = 1$ and $d_2(8_{17}, 6_3) = 1$.



FIGURE 3. The knot 8_{17} is transformed into 4_1 and 6_3 by $H(2)$ -moves.

Since the $H(2)$ -unknotting number of an unknotting number one knot is at most two by Theorem 3.1 in [7], Theorem 2.2 implies:

Corollary 2.4. *Let K be an unknotting number one knot and let $d = \det K$. Suppose that for any integer x*

$$(10) \quad x^2 \not\equiv \pm 2 \pmod{d},$$

i.e., both 2 and -2 are quadratic non-residues modulo d . Then $u_2(K) = 2$.

TABLE 1. Values for which $\det L$ does not take with $d_2(K, L) = 1$ or $d_{cb}(K, L) = 1$, K being an unknotting number one knot (I).

$\det K$	$\det L \neq$
5, 15, 35, 55, 95, 115	1, 4 (mod 5)
9, 27	3, 6 (mod 9)
13, 39, 91	1, 3, 4, 9, 10, 12 (mod 13)
17, 51	3, 5, 6, 7, 10, 11, 12, 14 (mod 17)
21	1, 4, 5, 16, 17, 20 (mod 21)
25, 75	1, 4, 5, 6, 9, 10, 11, 14, 15, 16, 19, 20, 21, 24 (mod 25)
29, 87	1, 4, 5, 6, 7, 9, 13, 16, 20, 22, 23, 24, 25, 28 (mod 29)
33	5, 7, 10, 13, 14, 19, 20, 23, 26, 28 (mod 33)
37, 111	1, 3, 4, 7, 9, 10, 11, 12, 16, 21, 25, 26, 27, 28, 30, 33, 34, 36 (mod 37)
41	3, 6, 7, 11, 12, 13, 14, 15, 17, 19, 22, 24, 26, 27, 28, 29, 30, 34, 35, 38 (mod 41)
45	1, 3, 4, 6, 9, 11, 12, 14, 15, 16, 19, 21, 24, 26, 29, 30, 31, 33, 34, 36, 39, 41, 42, 44 (mod 45)
49	7, 14, 21, 28, 35, 42 (mod 49)
53	1, 4, 6, 7, 9, 10, 11, 13, 15, 16, 17, 24, 25, 28, 29, 36, 37, 38, 40, 42, 43, 44, 46, 47, 49, 52 (mod 53)
57	5, 10, 11, 13, 17, 20, 22, 23, 26, 31, 34, 35, 37, 40, 44, 46, 47, 52 (mod 57)
61	1, 3, 4, 5, 9, 12, 13, 14, 15, 16, 19, 20, 22, 25, 27, 34, 36, 39, 41, 42, 45, 46, 47, 48, 49, 52, 56, 57, 58, 60 (mod 61)
63	1, 3, 4, 5, 6, 12, 15, 16, 17, 20, 21, 22, 24, 25, 26, 30, 33, 37, 38, 39, 41, 42, 43, 46, 47, 48, 51, 57, 58, 59, 60, 62 (mod 63)
65	1, 3, 4, 6, 9, 10, 11, 12, 14, 16, 17, 19, 21, 22, 23, 24, 25, 26, 27, 29, 30, 31, 34, 35, 36, 38, 39, 40, 41, 42, 43, 44, 46, 48, 49, 51, 53, 54, 55, 56, 59, 61, 62, 64 (mod 65)
69	2, 7, 8, 10, 19, 22, 26, 28, 29, 32, 34, 35, 37, 40, 41, 43, 47, 50, 59, 61, 62, 67 (mod 69)

Let d ($0 < d \leq 115$) be an integer such that both 2 and -2 are quadratic non-residues modulo d . Then from Tables 1 and 2 we have:

$$(11) \quad d = 5, 13, 15, 21, 25, 29, 35, 37, 39, 45, 53, 55, 61, 63, 65, 69, 75, 77, 85, 87, 91, 93, 95, 101, 105, 109, 111, 115.$$

Consequently, the following unknotting number one knots have $H(2)$ -unknotting number two; see [1, 2, 7] for the table of $H(2)$ -unknotting numbers of knots with up to nine crossings.

$$(12) \quad 4_1, 6_3, 7_7; 8_l, l = 1, 9, 13, 17, 21; 9_m, m = 2, 12, 14, 24, 30, 33, 39; 10_n; n = 9, 10, 18, 26, 32, 33, 59, 60, 71, 82, 84, 88, 95, 104, 107, 113, 114, 119, 129, 132, 136, 137, 141, 156, 159, 164.$$

TABLE 2. Values for which $\det L$ does not take with $d_2(K, L) = 1$ or $d_{cb}(K, L) = 1$, K being an unknotting number one knot (II).

$\det K$	$\det L \neq$
73	5, 7, 10, 11, 13, 14, 15, 17, 20, 21, 22, 26, 28, 29, 30, 31, 33, 34, 39, 40, 42, 43, 44, 45, 47, 51, 52, 53, 56, 58, 59, 60, 62, 63, 66, 68 (mod 73)
77	1, 4, 6, 9, 10, 13, 15, 16, 17, 19, 23, 24, 25, 36, 37, 40, 41, 52, 53, 54, 58, 60, 61, 62, 64, 67, 68, 71, 73, 76 (mod 77)
81	3, 6, 12, 15, 21, 24, 27, 30, 33, 39, 42, 48, 51, 54, 57, 60, 66, 69, 75, 78 (mod 81)
85	1, 3, 4, 5, 6, 7, 9, 10, 11, 12, 14, 16, 19, 20, 21, 22, 23, 24, 26, 27, 28, 29, 31, 34, 36, 37, 39, 40, 41, 44, 45, 46, 48, 49, 51, 54, 56, 57, 58, 59, 61, 62, 63, 64, 65, 66, 69, 71, 73, 74, 75, 76, 78, 79, 80, 81, 82, 84 (mod 85)
89	3, 6, 7, 12, 13, 14, 15, 19, 23, 24, 26, 27, 28, 29, 30, 31, 33, 35, 37, 38, 41, 43, 46, 48, 51, 52, 54, 56, 58, 59, 60, 61, 62, 63, 65, 66, 70, 74, 75, 76, 77, 82, 83, 86 (mod 89)
93	1, 4, 7, 10, 11, 16, 17, 19, 23, 25, 26, 28, 29, 40, 44, 49, 53, 64, 65, 67, 68, 70, 74, 76, 77, 82, 83, 86, 89, 92 (mod 93)
97	5, 7, 10, 13, 14, 15, 17, 19, 20, 21, 23, 26, 28, 29, 30, 34, 37, 38, 39, 40, 41, 42, 45, 46, 51, 52, 55, 56, 57, 58, 59, 60, 63, 67, 68, 69, 71, 74, 76, 77, 78, 80, 82, 83, 84, 87, 90, 92 (mod 97)
99	3, 5, 6, 7, 10, 12, 13, 14, 15, 19, 20, 21, 23, 24, 26, 28, 30, 33, 38, 39, 40, 42, 43, 46, 47, 48, 51, 52, 53, 56, 57, 59, 60, 61, 66, 69, 71, 73, 75, 76, 78, 79, 80, 84, 85, 86, 87, 89, 92, 93, 94, 96 (mod 99)
101	1, 4, 5, 6, 9, 13, 14, 16, 17, 19, 20, 21, 22, 23, 24, 25, 30, 31, 33, 36, 37, 43, 45, 47, 49, 52, 54, 56, 58, 64, 65, 68, 70, 71, 76, 77, 78, 79, 80, 81, 82, 84, 85, 87, 88, 92, 95, 96, 97, 100 (mod 101)
105	1, 4, 5, 6, 9, 11, 14, 16, 17, 19, 20, 21, 22, 24, 25, 26, 29, 31, 34, 36, 37, 38, 39, 41, 43, 44, 46, 47, 49, 51, 54, 56, 58, 59, 61, 62, 64, 66, 67, 68, 69, 71, 74, 76, 79, 80, 81, 83, 84, 85, 86, 88, 89, 91, 94, 96, 99, 100, 101, 104 (mod 105)
109	1, 3, 4, 5, 7, 9, 12, 15, 16, 20, 21, 22, 25, 26, 27, 28, 29, 31, 34, 35, 36, 38, 43, 45, 46, 48, 49, 60, 61, 63, 64, 66, 71, 73, 74, 75, 78, 80, 81, 82, 83, 84, 87, 88, 89, 93, 94, 97, 100, 102, 104, 105, 106, 108 (mod 109)
113	3, 5, 6, 10, 12, 17, 19, 20, 21, 23, 24, 27, 29, 33, 34, 35, 37, 38, 39, 40, 42, 43, 45, 46, 47, 48, 54, 55, 58, 59, 65, 66, 67, 68, 70, 71, 73, 74, 75, 76, 78, 79, 80, 84, 86, 89, 90, 92, 93, 94, 96, 101, 103, 107, 108, 110 (mod 113)

The numbers in (11) have a divisor congruent to 5 modulo 8. Indeed, using Lemma 2.6 below, Corollary 2.4 is restated as follows.

Corollary 2.5. *Let K be an unknotting number one knot. If the determinant of K is a multiple of $8k + 5$ for some k , $k = 0, 1, 2, \dots$, then $u_2(K) = 2$.*

Lemma 2.6. *A positive odd integer d is a multiple of $8k + 5$ for some k , $k = 0, 1, 2, \dots$ if and only if both 2 and -2 are quadratic non-residues modulo d .*

TABLE 3. Knots J and K with $d_2(J, K) > 1$, J being of unknotting number one.

K	$\det K$	J						
		4_1	6_1	6_3	7_7	8_9	8_{13}	8_{17}
U	1	×		×	×	×	×	×
$3_1, 8_{19}$	3		×	×				×
$4_1, 5_1$	5				×	×	×	
$5_2, 7_1$	7						×	×
$6_1, 3_1\#3_1, 8_{20}$	9	×		×		×	×	×
$6_2, 7_2$	11	×				×		×
$6_3, 7_3, 8_1$	13						×	
$7_4, 3_1\#4_1, 8_{21}, 3_1\#5_1$	15		×			×		
$7_5, 8_2, 8_3$	17			×	×			
$7_6, 8_4$	19	×				×		
$7_7, 8_5, 3_1\#5_2$	21	×	×			×		×
$8_6, 8_7$	23			×			×	
$8_8, 8_9, 4_1\#4_1$	25			×	×		×	×
$8_{10}, 8_{11}$	27			×				×
$8_{12}, 8_{13}$	29	×		×		×		
8_{14}	31	×				×		
8_{15}	33		×				×	×
8_{16}	35			×		×	×	
8_{17}	37				×			
	39	×	×			×		
	41	×			×	×		×
	43			×	×			
8_{18}	45					×	×	

In order to prove Lemma 2.6 we use the Jacobi symbol; cf. [4]. For an integer m and an odd prime number p the Legendre symbol (m/p) is defined by

$$(13) \quad \left(\frac{m}{p}\right) = \begin{cases} 1 & \text{if } m \text{ is a quadratic residue modulo } p \text{ and } m \not\equiv 0 \pmod{p}; \\ -1 & \text{if } m \text{ is a quadratic non-residue modulo } p. \end{cases}$$

For an odd positive integer n with prime factorization $n = p_1 p_2 \cdots p_r$, where p_i is a prime number, the Jacobi symbol (m/n) is defined by

$$(14) \quad \left(\frac{m}{n}\right) = \prod_{i=1}^r \left(\frac{m}{p_i}\right).$$

where (m/p_i) is the Legendre symbol.

If p_i is a quadratic residue modulo n for each i , then by the Chinese remainder theorem m is a quadratic residue modulo n . If m is a quadratic residue modulo n , then the Jacobi

TABLE 4. 2-component links L and unknotting number one knots J with $d_{\text{cb}}(J, L) > 1$.

L	$\det L$	J						
		4_1 8_{21}	6_1 8_{11}	6_3 8_1	7_7	8_9	8_{13}	8_{17}
U^2	0							
$H_-(= 2_1^2)$	2							
$T_4 = 4_1^2, 7_7^2$	4	×		×	×	×	×	×
$3_1 \# H_-, T_6 (= 6_1^2)$	6	×	×			×	×	
$5_1^2, 7_8^2, T_8 (= 8_1^2), 8_{15}^2$	8							
$6_2^2, 4_1 \# H_-, 5_1 \# H_-$	10			×		×		×
$6_3^2, 3_1 \# T_4, 8_{16}^2$	12		×	×				×
$7_1^2, 5_2 \# H_-$	14	×		×		×		
$7_3^2, 7_4^2, 8_2^2, 8_{12}^2$	16	×		×	×	×	×	×
7_2^2	18							
$7_5^2, 8_6^2$	20				×	×	×	
8_3^2	22			×	×		×	
$7_6^2, 8_4^2$	24	×	×			×	×	
8_5^2	26	×			×	×		×
$8_9^2, 8_{11}^2$	28						×	×
8_7^2	30		×	×		×	×	×
$8_{10}^2, 8_{12}^2$	32							
8_8^2	34	×				×	×	×
8_{14}^2	36	×		×		×	×	×
	38			×	×		×	×
8_{13}^2	40			×		×		×

TABLE 5. Invariants of the knots $5_2, 7_1,$ and 8_{17} .

K	$\sigma(K)$	$V(K; -1)$	$\text{Arf}(K)$	$u_2(K)$
5_2	2	-7	0	1
7_1	6	-7	0	1
8_{17}	0	37	1	2

symbol (m/n) is 1. We use the following formulas of the Jacobi symbol; cf. [13, p. 84].

$$(15) \quad \left(\frac{2}{n}\right) = \begin{cases} 1 & \text{if } n \equiv 1, 7 \pmod{8}; \\ -1 & \text{if } n \equiv 3, 5 \pmod{8}, \end{cases}$$

$$(16) \quad \left(\frac{-2}{n}\right) = \begin{cases} 1 & \text{if } n \equiv 1, 3 \pmod{8}; \\ -1 & \text{if } n \equiv 5, 7 \pmod{8}. \end{cases}$$

Proof of Lemma 2.6. If d is a multiple of $8k + 5$ for some k , $k = 0, 1, 2, \dots$, then both 2 and -2 are quadratic non-residues modulo d . In fact, the Jacobi symbols $(\pm 2/(8k + 5))$ are -1 by Eqs. (15) and (16).

Suppose that any divisor of d is not congruent to 5 modulo 8. Let $d = p_1 p_2 \dots p_m$ be a prime factorization; each p_i is an odd prime number. Then we have two cases:

- (i) $p_i \equiv 1, 3 \pmod{8}$ for each i ;
- (ii) $p_i \equiv 1, 7 \pmod{8}$ for each i .

In fact, if $p_i \equiv 3, p_j \equiv 7 \pmod{8}$, then $p_i p_j \equiv 5 \pmod{8}$. Then in Case (i) by Eq. (16) -2 is a quadratic residue modulo p_i for each i , and so -2 is a quadratic residue modulo d . Similarly, in Case (ii) 2 is a quadratic residue modulo d . This completes the proof. \square

3. CRITERION ON KNOTS WITH $H(2)$ -GORDIAN DISTANCE TWO

In [6] we have compiled a table of $H(2)$ -Gordian distances between knots with up to seven crossings, where we use several criteria to give lower bounds. In [8] we gave further criteria for giving a lower bound of the $H(2)$ -Gordian distance. In Theorems 7.1 and 7.3 in [6], the individual 6 pairs of knots are proved to have $H(2)$ -Gordian distance two by using the special value of the Jones polynomial, which is generalized to Theorem 5.2(iii) in [8].

We give a further criterion of a pair of knots with $H(2)$ -Gordian distance two:

Theorem 3.1. *Let K and K' be knots with $d_2(K, K') = 2$ and $V(K; \omega) = V(K'; \omega) = \pm(i\sqrt{3})^\delta$. If either*

- (i) $\sigma(K) - \sigma(K') \equiv 0 \pmod{8}$ and $\text{Arf}(K) \neq \text{Arf}(K')$, or
- (ii) $\sigma(K) - \sigma(K') \equiv 4 \pmod{8}$ and $\text{Arf}(K) = \text{Arf}(K')$,

then

$$(17) \quad V(K; -1) \equiv V(K'; -1) \pmod{3^{\delta+1}}.$$

Proof. Let J be a knot which is obtained from both K and K' by an $H(2)$ -move; $d_2(J, K) = d_2(J, K') = 1$. First, we show $\sigma(J) \equiv \sigma(K) \pm 2 \pmod{8}$. Indeed, if we assume $\sigma(J) \equiv \sigma(K) \pmod{4}$, then by Lemma 6.1 in [6] we have:

- If $\sigma(K) - \sigma(K') \equiv 0 \pmod{8}$, then $\text{Arf}(K) = \text{Arf}(K')$.
- If $\sigma(K) - \sigma(K') \equiv 4 \pmod{8}$, then $\text{Arf}(K) \neq \text{Arf}(K')$.

This contradicts our assumption.

Next, we show $V(J; \omega) = \pm v_0$, where $v_0 = V(K; \omega) = V(K'; \omega)$. By [6, Theorem 5.3] we have $V(J; \omega) \in \{\pm v_0, \pm i\sqrt{3}^{\pm 1} v_0\}$. Assume $V(J; \omega)/v_0 = \epsilon i\sqrt{3}$, $\epsilon = \pm 1$. Then by Theorem 5.5 in [6] we obtain:

- (a) If $\sigma(J) - \sigma(K) \equiv 2\epsilon \pmod{8}$, then $\text{Arf}(J) = \text{Arf}(K)$.
- (b) If $\sigma(J) - \sigma(K) \equiv -2\epsilon \pmod{8}$, then $\text{Arf}(J) \neq \text{Arf}(K)$.
- (c) If $\sigma(J) - \sigma(K') \equiv 2\epsilon \pmod{8}$, then $\text{Arf}(J) = \text{Arf}(K')$.
- (d) If $\sigma(J) - \sigma(K') \equiv -2\epsilon \pmod{8}$, then $\text{Arf}(J) \neq \text{Arf}(K')$.

Therefore, K and K' do not satisfy the conditions (i) nor (ii). Similarly, we may prove $V(J; \omega) \neq \pm i\sqrt{3}^{-1} v_0$.

Thus we have $V(J; \omega) = \eta v_0$, $\eta = \pm 1$. Then by Theorem 5.2(iii) in [8] we obtain $\eta V(J; -1) \equiv V(K; -1) \equiv V(K'; -1) \pmod{3^{\delta+1}}$, completing the proof. \square

Example 3.2. Let $K = 6_1!$ and $K' = 7_7$. Then $V(K; \omega) = V(K'; \omega) = -i\sqrt{3}$, $\sigma(K) = \sigma(K') = 0$, $\text{Arf}(K) = 0$, $\text{Arf}(K') = 1$, and $9 = V(K; -1) \not\equiv V(K'; -1) = 21 \pmod{9}$. Thus by Theorem 3.1 we obtain $d_2(K, K') \neq 2$, which implies $d_2(K, K') = 3$ by [6, Table 3].

4. $H(2)$ -GORDIAN DISTANCES OF KNOTS WITH UP TO SEVEN CROSSINGS

In Tables 6 and 7 we list the $H(2)$ -Gordian distances between knots with up to seven crossings, which improves those in [6], where the meanings of the marks are as follows:

- The marks 1-2, 2-3, 1-3 mean 1 or 2, 2 or 3, 1 or 2 or 3, respectively.
- The mark 1^z) means that the distance is confirmed to be 1 by Fig. 20 in [14], and 2^z) means that the distance is decided to be 2 from the inequality $d_2(7_1, 7_7) \leq d_2(7_1, 6_3) + d_2(6_3, 7_7) = 2$.
- The mark 2^m) means that the distance is decided to be 2 by using Theorem 4.2 (Example 4.5) in [8].
- The mark 2^{mn}) means that the distance is decided to be 2 by using either by Theorem 4.2 (Example 4.5) in [8] or Theorem 2.2 (Table 3).
- The mark 2^v) means that the distance is decided to be 2 by using Theorem 5.8 (Example 5.10) in [8].
- The mark 2^{mnv}) means that the distance is decided to be 2 by using by either Theorem 4.2 (Example 4.5) in [8], Theorem 2.2 (Table 3), or Theorem 5.8 (Example 5.10) in [8].
- The mark 3^v) means that the distance is decided to be 3 by using Theorem 3.1 (Example 3.2).

ACKNOWLEDGEMENTS

The author was partially supported by KAKENHI, Grant-in-Aid for Scientific Research (C) (No. 21540092), Japan Society for the Promotion of Science. He also would like to thank Professor Masaaki Furusawa for his suggestion for the proof of Lemma 2.6.

REFERENCES

1. Tetsuya Abe and Taizo Kanenobu, *Unoriented band surgery on knots and links*, Kobe J. Math. **31** (2014), no. 1-2, 21–44. MR 3307286
2. Yuanyuan Bao, *$H(2)$ -unknotting operation related to 2-bridge links*, Topology Appl. **159** (2012), no. 8, 2158–2167. MR 2902750
3. Joan S. Birman and Taizo Kanenobu, *Jones' braid-plat formula and a new surgery triple*, Proc. Amer. Math. Soc. **102** (1988), no. 3, 687–695. MR 929004 (89c:57003)
4. Kenneth Ireland and Michael Rosen, *A classical introduction to modern number theory*, second ed., Graduate Texts in Mathematics, vol. 84, Springer-Verlag, New York, 1990. MR 1070716 (92e:11001)
5. V. F. R. Jones, *Hecke algebra representations of braid groups and link polynomials*, Ann. of Math. (2) **126** (1987), no. 2, 335–388. MR 908150 (89c:46092)
6. Taizo Kanenobu, *$H(2)$ -Gordian distance of knots*, J. Knot Theory Ramifications **20** (2011), no. 6, 813–835. MR 2812265 (2012e:57011)

TABLE 6. $H(2)$ -Gordian distances of knots with up to 6 crossings.

	3_1	4_1	5_1	5_2	6_1	6_2	6_3	$3_1\#3_1$	$3_1\#\#3_1$
U	1	2	1	1	1	1	2	2	2
3_1	0	1	2	2	2	1	2	1	1
$3_1!$	2	1	2	1	2	2	2	1	1
4_1		0	2-3	1	2	2	1	2	2
5_1			0	2	1-2	1	2	3	2
$5_1!$			1 ^{z)}	2	1	1-2	2	3	2
5_2				0	1	2	1	2	2
$5_2!$				2	2	1	1	2	2
6_1					0	1-2	2	2-3	1
$6_1!$					1-2	2	2	2-3	1
6_2						0	1	2	2
$6_2!$						2	1	2	2
6_3							0	2	3
$3_1\#3_1$								0	2
$3_1\#\#3_1!$								1-2	2
$3_1!\#3_1$								2	0

7. Taizo Kanenobu and Yasuyuki Miyazawa, *H(2)-unknotting number of a knot*, Commun. Math. Res. **25** (2009), no. 5, 433–460. MR 2573402 (2010i:57019)
8. Taizo Kanenobu and Hiromasa Moriuchi, *Links which are related by a band surgery or crossing change*, Bol. Soc. Mat. Mex. (3) **20** (2014), no. 2, 467–483. MR 3264627
9. Hitoshi Murakami, *Some metrics on classical knots*, Math. Ann. **270** (1985), no. 1, 35–45.
10. Yasutaka Nakanishi, *A note on unknotting number. II*, J. Knot Theory Ramifications **14** (2005), no. 1, 3–8. MR 2124552 (2006b:57006)
11. Yasutaka Nakanishi and Yuki Okada, *Differences of Alexander polynomials for knots caused by a single crossing change*, Topology Appl. **159** (2012), no. 4, 1016–1025. MR 2876708 (2012m:57014)
12. Dale Rolfsen, *Knots and links*, Mathematics Lecture Series, vol. 7, Publish or Perish, Inc., Houston, TX, 1990, Corrected reprint of the 1976 original. MR 1277811 (95c:57018)
13. Teiji Takagi, *Shotou Seisûron Kôgi [Lectures on elementary number theory]*, second ed., Kyoritsu Shuppan, Tokyo, 1971, Japanese.
14. Ana Zeković, *Computation of Gordian distances and H_2 -Gordian distances of knots*, Yugosl. J. Oper. Res. **25** (2015), no. 1, 133–152. MR 3331990

DEPARTMENT OF MATHEMATICS, OSAKA CITY UNIVERSITY, SUGIMOTO, SUMIYOSHI-KU, OSAKA 558-8585, JAPAN

E-mail address: kanenobu@sci.osaka-cu.ac.jp

