

UNIQUE TORIC STRUCTURE ON A FANO BOTT MANIFOLD

メタデータ	言語: English 出版者: OCAMI 公開日: 2022-11-09 キーワード (Ja): キーワード (En): 作成者: Cho, Yunhyung, Lee, Eunjeong, 栴田, 幹也, 朴, 善正 メールアドレス: 所属: Sungkyunkwan University, Institute for Basic Science, Osaka City University, KAIST
URL	https://ocu-omu.repo.nii.ac.jp/records/2016898

Unique toric structure on a Fano Bott manifold

Yunhyung Cho, Eunjeong Lee, Mikiya Masuda, Seonjeong Park

Citation	OCAMI Preprint Series. 2020; 10
Issue Date	2020-07-29
Type	Preprint
Textversion	Author

From: Osaka City University Advanced Mathematical Institute

<http://www.sci.osaka-cu.ac.jp/OCAMI/publication/preprint/preprint.html>

UNIQUE TORIC STRUCTURE ON A FANO BOTT MANIFOLD

YUNHYUNG CHO, EUNJEONG LEE, MIKIYA MASUDA, AND SEONJEONG PARK

ABSTRACT. We prove that if there exists a c_1 -preserving graded ring isomorphism between integral cohomology rings of two Fano Bott manifolds, then they are isomorphic as toric varieties. As a consequence, we give an affirmative answer to McDuff's question on the uniqueness of a toric structure on a Fano Bott manifold.

1. INTRODUCTION

To each symplectic manifold (M, ω) , one can associate the Hamiltonian diffeomorphism group $\text{Ham}(M, \omega)$. It is a normal subgroup of the symplectomorphism group $\text{Symp}(M, \omega)$ and governs all possible Hamiltonian Lie group actions on (M, ω) . The group $\text{Ham}(M, \omega)$ is infinite dimensional and non-compact in general, and it might possess more than one maximal tori with distinct conjugacy classes. It turned out that the number of conjugacy classes of maximal tori in $\text{Ham}(M, \omega)$, allowing conjugations by elements of $\text{Symp}(M, \omega)$, is finite; it was proved by Karshon–Kessler–Pinsonnault [KKP07] and Pinsonnault [Pin08] in dimension four, and by McDuff [McD11, Proposition 3.1] in any dimension.

Recall that a symplectic form ω is called *monotone* if $c_1(M) := c_1(TM, J) = \lambda \cdot [\omega]$ for some $\lambda > 0$ and an ω -compatible almost complex structure J on M . Throughout this paper, we always assume that $\lambda = 1$ unless stated otherwise. In this paper, we concern the following question posed by McDuff [McD11].

Question 1.1 (McDuff). [McD11, Question 1.11] *Let (M, ω) be a $2n$ -dimensional closed monotone symplectic manifold. If T_1 and T_2 are n -tori in $\text{Ham}(M, \omega)$, are (M, ω, T_1) and (M, ω, T_2) equivariantly symplectomorphic? Equivalently, is the number of conjugacy classes of n -tori in $\text{Ham}(M, \omega)$ precisely one?*

Due to the Delzant's theorem [Del88], any closed symplectic toric manifold is equivariantly symplectomorphic to a smooth projective toric variety equipped with a torus invariant Kähler form. When a symplectic form is monotone, it is equivariantly symplectomorphic to a smooth Fano toric variety by the Kleiman's ampleness criterion [Kle66, Theorem 1 in Section III-1] and its moment polytope becomes a *reflexive*¹ polytope.

In the algebro-geometric aspects, Question 1.1 asks whether a smooth Fano toric variety may have more than one toric structure. McDuff [McD11] gave an affirmative answer to Question 1.1 when $M = \mathbb{C}P^k \times \mathbb{C}P^m$, and Fanoe [Fan14] generalized McDuff's result to the case of $\mathbb{C}P^k$ -bundle over $\mathbb{C}P^m$. To the best of the authors' knowledge, Question 1.1 is still open.

¹A polytope is *reflexive* if it is integral and has a unique interior lattice point such that the affine distance from the point to each facet is equal to one.

This paper is addressed to Question 1.1 in case that M is a Bott manifold. A *Bott tower*, first introduced by Grossberg and Karshon [GK94], is an iterated $\mathbb{C}P^1$ -bundle starting with a point

$$\mathcal{B}_n \xrightarrow{\pi_n} \mathcal{B}_{n-1} \xrightarrow{\pi_{n-1}} \cdots \longrightarrow \mathcal{B}_1 = \mathbb{C}P^1 \xrightarrow{\pi_1} \mathcal{B}_0 = \{\text{a point}\}$$

where each \mathcal{B}_i is obtained by projectivizing the direct sum of the trivial line bundle $\underline{\mathbb{C}}$ and a complex line bundle ξ_i over \mathcal{B}_{i-1} , i.e., $\mathcal{B}_i = P(\underline{\mathbb{C}} \oplus \xi_i)$. The total space \mathcal{B}_n is called a *Bott manifold*.

We may equip a natural complex structure on a Bott manifold by taking each ξ_i as a holomorphic line bundle so that \mathcal{B}_n becomes a complex manifold with a natural $(\mathbb{C}^*)^n$ -action constructed in an iterative way using a toric structure of a base space and a \mathbb{C}^* -action on a fiber at each stage, see [Oda78, Section I-7.6']. Indeed, \mathcal{B}_n is a smooth projective toric variety.

Any Bott manifold \mathcal{B}_n can be characterized by an $n \times n$ lower triangular integer matrix called a *Bott matrix*. Roughly speaking, a Bott manifold \mathcal{B}_n is a smooth projective toric variety, where the corresponding fan is combinatorially equivalent to the normal fan of the n -cube. After fixing one reference maximal cone of the fan and making it into the “first quadrant” via some basis change by multiplying a suitable element of $\mathrm{GL}(n, \mathbb{Z})$, we obtain residual n (column) vectors which form a Bott matrix. Note that a Bott matrix presentation of a Bott manifold is not unique and it depends on the choice of a reference cone and an element of $\mathrm{GL}(n, \mathbb{Z})$. In Section 3, we illustrate some operations (temporarily denoted by **Op.1** and **Op.2** in this paper) on the set of Bott matrices and describe cohomology ring isomorphisms induced by the operations that will be crucially used in the proof of our main theorem.

Not all Bott manifolds are Fano. For instance, there are only two Fano Bott manifolds in dimension four: $\mathbb{C}P^1 \times \mathbb{C}P^1$ and a Hirzebruch surface $P(\underline{\mathbb{C}} \oplus \mathcal{O}(1))$. Recently, Suyama [Suy19] classified all Fano Bott manifolds in terms of Bott matrices, see Section 2 for the detail. Now we state our main theorem which says that every Fano Bott manifold is characterized (as a toric variety) by its integral cohomology ring and the first Chern class.

Theorem 1.2 (Theorem 4.2). *Let X and Y be Fano Bott manifolds. If there exists a c_1 -preserving graded ring isomorphism*

$$\varphi: H^*(X; \mathbb{Z}) \rightarrow H^*(Y; \mathbb{Z}),$$

then X and Y are isomorphic² as toric varieties, i.e., the fans associated to X and Y are unimodularly equivalent.

We call a monotone symplectic manifold (M, ω) a *monotone Bott manifold* if M is diffeomorphic to a Bott manifold. Using Theorem 1.2, we obtain a positive answer to Question 1.1.

Corollary 1.3. *Any monotone Bott manifold has a unique toric structure.*

Proof. Suppose that (M, ω) is a $2n$ -dimensional monotone Bott manifold and T_1, T_2 are n -tori in $\mathrm{Ham}(M, \omega)$. From the Delzant’s theorem [Del88], each T_i -action makes M into a toric Fano variety

²We say that two toric varieties X and Y are *isomorphic as toric varieties* if there exists a toric isomorphism ϕ from X to Y , i.e., $\phi(t \cdot x) = \phi(t) \cdot \phi(x)$ for every element t in the torus $T_X \subset X$ and $x \in X$. In fact, it was proved in [Ber03] that there exists a toric isomorphism if X and Y are isomorphic as abstract varieties.

(which we denote by X_i) with T_i -invariant complex structure J_i on M . Note that each J_i can be chosen to be ω -compatible so that $c_1(X_1) = c_1(X_2) = [\omega] \in H^2(M; \mathbb{Z})$.

On the other hand, it follows from [MP08, Corollary 3.5 and Theorem 5.5] that any smooth projective toric variety whose integral cohomology ring is isomorphic to that of a Bott manifold is in fact isomorphic to a Bott manifold as a toric variety. (See Remark 2.3 for the detail.) Thus we may assume that X_1 and X_2 are Fano Bott manifolds. Then the identity map $H^*(M; \mathbb{Z}) \rightarrow H^*(M; \mathbb{Z})$ induces a graded ring isomorphism

$$H^*(X_1; \mathbb{Z}) \rightarrow H^*(X_2; \mathbb{Z}), \quad c_1(X_1) = [\omega] \mapsto [\omega] = c_1(X_2).$$

Thus the result follows from Theorem 1.2. □

It is worth mentioning a relation between Theorem 1.2 and a problem posed by the third author and Suh [MS08, Problems 1 and 4] which asks whether two smooth complete toric varieties having isomorphic cohomology rings (as graded rings) are diffeomorphic or not. This problem is now called the *cohomological rigidity* for toric varieties. There are many partial affirmative answers to the problem. For instance, two smooth complete toric varieties with Picard number 2 are diffeomorphic if and only if their integral cohomology rings are isomorphic as graded rings, see [CMS10]. We also refer the reader to [CMS11, BEM⁺17] and references therein for recent accounts of this problem.

Inspired by Theorem 1.2, we pose the following conjecture.

Conjecture 1.4. *Suppose that X and Y are smooth toric Fano varieties. If there exists a c_1 -preserving graded ring isomorphism between their integral cohomology rings, then X and Y are isomorphic as toric varieties.*

Conjecture 1.4 was verified for some other classes of smooth toric Fano varieties. Indeed, the authors confirmed Conjecture 1.4 for smooth toric Fano varieties with Picard number 2, whose proof will be provided in an upcoming manuscript [CLMP20]. Also the third author together with Higashitani and Kurimoto [HKM20] proved Conjecture 1.4 for smooth toric Fano varieties with small dimension ($\dim X_{\mathbb{C}} \leq 4$) or with large Picard number.)

Note that if Conjecture 1.4 is true, then the answer to Question 1.1 is positive. More precisely, if (M, ω, T_1) and (M, ω, T_2) are two toric structures over the same monotone symplectic manifold (M, ω) , then the identity map on $H^*(M; \mathbb{Z})$ satisfies the hypothesis in Conjecture 1.4. Therefore, Conjecture 1.4 can be thought as a stronger version of Question 1.1.

As a final remark, we would like to mention a recent work of Pabiniak and Tolman. In [PT20], they considered the following question which they called *symplectic cohomological rigidity*.

Question 1.5. [PT20, p.3] *Let (M_1, ω_1) and (M_2, ω_2) be symplectic toric manifolds. If there exists a graded ring isomorphism between their integral cohomology rings sending $[\omega_1]$ to $[\omega_2]$, are (M_1, ω_1) and (M_2, ω_2) symplectomorphic?*

They also gave a positive answer to Question 1.5 under the assumptions that ω_1 and ω_2 are rational symplectic forms and that $H^*(M_1; \mathbb{Q}) \cong H^*(M_2; \mathbb{Q}) \cong H^*(\mathbb{C}P^1 \times \cdots \times \mathbb{C}P^1; \mathbb{Q})$.

This paper is organized as follows. In Section 2, we explain the notion of Bott manifolds and also discuss their cohomological properties. In Section 3, we introduce two operations **Op.1** and **Op.2** on Bott matrices and prove that any two Bott matrices which represent isomorphic Bott manifolds are obtained by applying those operations repeatedly. In Section 4, we give the proof of Theorem 1.2.

Acknowledgements. Y. Cho was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIP; Ministry of Science, ICT & Future Planning) (NRF-2020R1C1C1A01010972). E. Lee was supported by IBS-R003-D1. M. Masuda was supported in part by JSPS Grant-in-Aid for Scientific Research 16K05152. S. Park was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Government of Korea (NRF-2018R1A6A3A11047606). This work was partly supported by Osaka City University Advanced Mathematical Institute (MEXT Joint Usage/Research Center on Mathematics and Theoretical Physics).

2. BOTT MANIFOLDS

We begin by recalling the definition of Bott towers and Bott manifolds.

Definition 2.1. [GK94, §2.1] A *Bott tower* \mathcal{B}_\bullet is an iterated $\mathbb{C}P^1$ -bundle starting with a point:

$$(2.1) \quad \begin{array}{ccccccc} \mathcal{B}_n & \xrightarrow{\pi_n} & \mathcal{B}_{n-1} & \xrightarrow{\pi_{n-1}} & \cdots & \xrightarrow{\pi_2} & \mathcal{B}_1 \xrightarrow{\pi_1} \mathcal{B}_0, \\ \parallel & & & & & & \parallel & \parallel \\ P(\underline{\mathbb{C}} \oplus \xi_n) & & & & & & \mathbb{C}P^1 & \{\text{a point}\} \end{array}$$

where each \mathcal{B}_i is the complex projectivization of the Whitney sum of a holomorphic line bundle ξ_i and the trivial bundle $\underline{\mathbb{C}}$ over \mathcal{B}_{i-1} . The total space \mathcal{B}_n is called a *Bott manifold*.

Let γ_j be the tautological line bundle over \mathcal{B}_j and $\gamma_{i,j}$ the pullback of γ_j by the projection $\pi_i \circ \cdots \circ \pi_{j+1}: \mathcal{B}_i \rightarrow \mathcal{B}_j$ for $i > j$. We also define $\gamma_{j,j} := \gamma_j$ for convenience. The Picard group of \mathcal{B}_{i-1} is isomorphic to the free abelian group of rank $i-1$, and is generated by the line bundles $\gamma_{i-1,j}$ for $1 \leq j < i$ by [Har77, Exercise II.7.9]. Therefore, for each $i = 2, \dots, n$, there exist $a_{i,j} \in \mathbb{Z}$ for $1 \leq j < i$ such that

$$(2.2) \quad \xi_i = \bigotimes_{1 \leq j < i} \gamma_{i-1,j}^{\otimes a_{i,j}}.$$


Thus the set of integers $\{a_{i,j}\}_{1 \leq j < i \leq n}$ determines a Bott manifold.

Each projection $\pi_i: \mathcal{B}_i \rightarrow \mathcal{B}_{i-1}$ admits a section induced from the zero section of $\underline{\mathbb{C}} \oplus \xi_i$. This implies that

$$\pi_i^*: H^*(\mathcal{B}_{i-1}; \mathbb{Z}) \rightarrow H^*(\mathcal{B}_i; \mathbb{Z})$$

is an injective ring homomorphism. By abuse of notation, we continue to write $x_j \in H^2(\mathcal{B}_{i-1}; \mathbb{Z})$ for the first Chern class of the dual of $\gamma_{i-1,j}$ for each $i > j$. From (2.2), we obtain

$$c_1(\xi_i) = - \sum_{j=1}^{i-1} a_{i,j} x_j \in H^2(\mathcal{B}_{i-1}).$$


 FIGURE 1. The fan of \mathcal{H}_1 .

On the other hand, a Bott manifold \mathcal{B}_n is a smooth projective toric variety by the construction (cf. [GK94] and [Oda78, Section I-7.6']). If \mathcal{B}_n is obtained from $\{a_{i,j}\}_{1 \leq j < i \leq n}$, then it is known from [Civ05, §3] that its fan has $2n$ rays and their generators are column vectors of the following matrix

$$(2.3) \quad (E \mid A) := \begin{bmatrix} 1 & & & -1 & & & \\ & 1 & & a_{2,1} & -1 & & \\ & & 1 & a_{3,1} & a_{3,1} & -1 & \\ & & & \vdots & \vdots & \ddots & \ddots \\ & & & 1 & a_{n,1} & a_{n,2} & \cdots & a_{n,n-1} & -1 \end{bmatrix},$$

where E is the $n \times n$ identity matrix. We call an integer matrix of the form A in (2.3) a *Bott matrix*. For a given Bott matrix A , we denote by

$$\mathbf{e}_j := j\text{-th standard basis vector}, \quad \mathbf{v}_j := j\text{-th column vector of } A.$$

The fan Σ_A of the Bott manifold \mathcal{B}_n has 2^n number of maximal cones $\Sigma(n) = \{\sigma_I \mid I \subset [n]\}$, where $[n] := \{1, \dots, n\}$ and

$$(2.4) \quad \sigma_I = \text{Cone}(\{\mathbf{e}_j \mid j \in I\} \cup \{\mathbf{v}_j \mid j \in I^c\}).$$

Example 2.2. Let $n = 2$. Then the Bott manifold determined by $\{a_{2,1}\}$ is a Hirzebruch surface $\mathcal{H}_{a_{2,1}} := P(\mathbb{C} \oplus \mathcal{O}(-a_{2,1}))$. There are four maximal cones:

$$\sigma_{\emptyset} = \text{Cone}\{\mathbf{v}_1, \mathbf{v}_2\}, \quad \sigma_{\{1\}} = \text{Cone}\{\mathbf{e}_1, \mathbf{v}_2\}, \quad \sigma_{\{2\}} = \text{Cone}\{\mathbf{e}_2, \mathbf{v}_1\}, \quad \sigma_{\{1,2\}} = \text{Cone}\{\mathbf{e}_1, \mathbf{e}_2\}.$$

We present the fan of \mathcal{H}_1 in Figure 1.

Let \mathcal{M}_n be the set of all Bott matrices of size $n \times n$, i.e., the set of all $n \times n$ lower triangular integer matrix with -1 's on the main diagonal as in (2.3). Since a Bott matrix A determines the fan Σ_A of a Bott manifold, we denote the corresponding Bott manifold by $\mathcal{B}(A)$. Note that it happens that $\mathcal{B}(A)$ and $\mathcal{B}(A')$ are isomorphic as toric varieties even if A and A' are different.

Remark 2.3. For a given smooth projective toric variety M , if its integral cohomology ring is isomorphic to that of a certain Bott manifold \mathcal{B} as graded rings, then the fan of M is combinatorially equivalent to that of \mathcal{B} , i.e., it is combinatorially equivalent to the normal fan of the n -cube (see [MP08, Theorem 5.5]). Moreover, such a toric variety M is again a Bott manifold by [MP08, Corollary 3.5]. Accordingly, any smooth projective toric variety whose integral cohomology ring is isomorphic to that of a Bott manifold is isomorphic to a Bott manifold.

2.1. Cohomology rings. By the Borel–Hirzebruch formula [BH58], the integral cohomology ring of a Bott manifold $\mathcal{B}(A)$ for $A \in \mathcal{M}_n$ is described by

$$\begin{aligned}
 H^*(\mathcal{B}(A); \mathbb{Z}) &\cong \mathbb{Z}[x_1, \dots, x_n] / \langle x_i^2 + c_1(\xi_i)x_i \mid i = 1, \dots, n \rangle \\
 (2.5) \quad &\cong \mathbb{Z}[x_1, \dots, x_n] / \langle x_i^2 - (a_{i,1}x_1 + \dots + a_{i,i-1}x_{i-1})x_i \mid i = 1, \dots, n \rangle \\
 &\cong \mathbb{Z}[x_1, \dots, x_n] / \langle x_i^2 - \alpha_i x_i \mid i = 1, 2, \dots, n \rangle.
 \end{aligned}$$

Here, x_i is the first Chern class of the dual of $\gamma_{n,i}$, and we set

$$\alpha_i := a_{i,1}x_1 + \dots + a_{i,i-1}x_{i-1} \in H^2(\mathcal{B}(A); \mathbb{Z}).$$

Note that x_1, \dots, x_n are of degree two and they generate the cohomology ring $H^*(\mathcal{B}(A); \mathbb{Z})$.

For a \mathbb{Q} -coefficient, we often use the following notation

$$(2.6) \quad y_i := x_i - \frac{1}{2}\alpha_i \in H^2(\mathcal{B}(A); \mathbb{Q}).$$

Note that y_i 's may not be integral classes but they generate $H^*(\mathcal{B}(A); \mathbb{Q})$ as a ring.

Recall from [Oda88, Theorem 3.12] that the total Chern class of a Bott manifold $\mathcal{B}(A)$ is written by

$$c(\mathcal{B}(A)) = \prod_{i=1}^n (1 + x_i)(1 + x_i - \alpha_i) = \prod_{i=1}^n (1 + 2x_i - \alpha_i).$$

Substituting (2.6) in the above, we get

$$(2.7) \quad c(\mathcal{B}(A)) = \prod_{i=1}^n (1 + 2y_i)$$

which implies that

$$(2.8) \quad c_1(\mathcal{B}(A)) = 2 \sum_{i=1}^n y_i \quad \text{and} \quad c_n(\mathcal{B}(A)) = 2^n \prod_{i=1}^n y_i.$$

Remark 2.4. Note that the cohomology ring description in (2.5) can be obtained from the Danilov–Jurkiewicz theorem, see [BP15, Theorem 5.3.1]. It follows that the even Betti number $b_{2i}(\mathcal{B}(A))$ is $\binom{n}{i}$ for $1 \leq i \leq n$. Since $H^*(\mathcal{B}(A); \mathbb{Z})$ is generated by degree two elements $\{x_i \mid 1 \leq i \leq n\}$, we obtain the following.

- (1) Since x_1, \dots, x_n are linearly independent (over \mathbb{Z}), so are y_1, \dots, y_n (over \mathbb{Q}).
- (2) The set $\{x_{i_1} \cdots x_{i_k} \mid 1 \leq i_1 < \dots < i_k \leq n\}$ is a \mathbb{Z} -basis of $H^{2k}(\mathcal{B}(A); \mathbb{Z})$.
- (3) $\prod_{i=1}^n y_i$ is the orientation class of $\mathcal{B}(A)$. Indeed, since the Euler characteristic of $\mathcal{B}(A)$ is 2^n and it agrees with $c_n(\mathcal{B}(A))$ evaluated on the fundamental class of $\mathcal{B}(A)$, $\prod_{i=1}^n y_i$ evaluated on the fundamental class is 1.

In terms of y_i 's, we can obtain a simple description of (any) isomorphisms of $H^*(\mathcal{B}(A); \mathbb{Q})$ as follows. In the following, we denote x_i , y_i and α_i in $H^*(\mathcal{B}(A))$ by x_i^A , y_i^A and α_i^A , respectively.

Proposition 2.5. [CMM15, Proposition 4.1] *For A and A' in \mathcal{M}_n , if we have a graded ring isomorphism*

$$\varphi: H^*(\mathcal{B}(A); \mathbb{Q}) \rightarrow H^*(\mathcal{B}(A'); \mathbb{Q}),$$

then there are nonzero $q_1, \dots, q_n \in \mathbb{Q}$ and a permutation σ on $[n]$ such that

$$\varphi(y_i^A) = q_i y_{\sigma(i)}^{A'} \quad \text{for } i = 1, \dots, n.$$

Theorem 2.6. *Suppose that there is a c_1 -preserving graded cohomology ring isomorphism between two Bott manifolds. Then all q_i 's in Proposition 2.5 are equal to 1. Moreover, it preserves their total Chern classes and hence all the Chern numbers of the two Bott manifolds are the same.*

Proof. Let \mathcal{B} and \mathcal{B}' be Bott manifolds determined by Bott matrices A and A' , respectively. Let φ be the c_1 -preserving graded cohomology ring isomorphism between \mathcal{B} and \mathcal{B}' . By (2.8), we have

$$\varphi \left(2 \sum_{i=1}^n y_i^A \right) = 2 \sum_{i=1}^n y_i^{A'}.$$

On the other hand, it follows from Proposition 2.5 that

$$\varphi \left(2 \sum_{i=1}^n y_i^A \right) = 2 \sum_{i=1}^n q_i y_{\sigma(i)}^{A'}.$$

Comparing these two identities, we obtain

$$2 \sum_{i=1}^n y_i^{A'} = 2 \sum_{i=1}^n q_i y_{\sigma(i)}^{A'}.$$

Here, $y_1^{A'}, \dots, y_n^{A'}$ are linearly independent, so we conclude $q_i = 1$ for any i . This together with (2.7) shows that φ preserves their total Chern classes, proving the former part of the theorem.

The latter part of the theorem follows from the former part and the fact that φ preserves the orientation classes $\prod_{i=1}^n y_i^A$ and $\prod_{i=1}^n y_i^{A'}$ (as well as top Chern classes). \square

Remark 2.7. We note the following.

- (1) Not every graded cohomology ring isomorphism between Bott manifolds is c_1 -preserving. For instance, one can find such an isomorphism for Hirzebruch surfaces \mathcal{H}_0 and \mathcal{H}_2 . See Example 2.8.
- (2) Recall that two Hirzebruch surfaces \mathcal{H}_a and \mathcal{H}_b are isomorphic if and only $|a| = |b|$. However, for any integers a and b with the same parity, there is a c_1 -preserving graded cohomology ring isomorphism between Hirzebruch surfaces \mathcal{H}_a and \mathcal{H}_b . Therefore, the existence of such a cohomology ring isomorphism does not imply that two varieties are isomorphic. We would need to restrict our attention to Fano Bott manifolds to conclude a variety isomorphism. See Remark 4.3.

Example 2.8. It follows from (2.5) that

$$H^*(\mathcal{H}_0; \mathbb{Z}) = \mathbb{Z}[x_1, x_2] / \langle x_1^2, x_2^2 \rangle \quad \text{and} \quad H^*(\mathcal{H}_2; \mathbb{Z}) = \mathbb{Z}[x'_1, x'_2] / \langle (x'_1)^2, x'_2(x'_2 - 2x'_1) \rangle.$$

Note that $c_1(\mathcal{H}_0) = 2x_1 + 2x_2$ and $c_1(\mathcal{H}_2) = 2x'_2$. Then the map $\varphi: H^*(\mathcal{H}_0) \rightarrow H^*(\mathcal{H}_2)$ given by $\varphi(x_1) = x'_1$ and $\varphi(x_2) = x'_1 - x'_2$ is a graded ring isomorphism which does not preserve the first Chern class. On the other hand, the map φ' defined by $\varphi'(x_1) := x'_1$ and $\varphi'(x_2) := x'_2 - x'_1$ is a c_1 -preserving isomorphism from $H^*(\mathcal{H}_0)$ to $H^*(\mathcal{H}_2)$. (Note that $\varphi'(c_1(\mathcal{H}_0)) = \varphi'(2x_1 + 2x_2) = 2x'_2 = c_1(\mathcal{H}_2)$.)

2.2. Fano Bott manifolds. In this subsection, we recall a description of Fano Bott manifolds from [Suy19].

Theorem 2.9. [Suy19, Theorem 8] *A Bott manifold $\mathcal{B}(A)$ is Fano if and only if each column of $A + E$ has values in $\{-1, 0, 1\}$ and it satisfies one of the following:*

- (1) *all entries are zero,*
- (2) *there is at most one 1 and every other entry below the 1 vanishes (if there is 1 on the column),*
- (3) *if there is -1 at the i -th row, then the entries below the -1 coincide with the entries on the i -th column below the diagonal $a_{ii} = -1$.*

Example 2.10. Consider the following Bott matrices.

$$A_1 = \begin{bmatrix} -1 & 0 & 0 \\ 1 & -1 & 0 \\ 2 & 1 & -1 \end{bmatrix}, \quad A_2 = \begin{bmatrix} -1 & 0 & 0 \\ 1 & -1 & 0 \\ 1 & 0 & -1 \end{bmatrix}, \quad A_3 = \begin{bmatrix} -1 & 0 & 0 \\ -1 & -1 & 0 \\ 0 & 1 & -1 \end{bmatrix}, \quad A_4 = \begin{bmatrix} -1 & 0 & 0 \\ -1 & -1 & 0 \\ 1 & 1 & -1 \end{bmatrix}.$$

One can easily check that only A_4 satisfies all the conditions in Theorem 2.9 and so $\mathcal{B}(A_4)$ is the only Fano Bott manifold among $\mathcal{B}(A_k)$'s for $1 \leq k \leq 4$. Indeed, A_4 is (10) on the list of Fano threefolds in the book of Oda [Oda88, Figure 2.6] and (12) on the list of ‘Smooth toric Fano varieties’ [Obr07] in the Graded Ring Database [BK].

3. OPERATIONS ON BOTT MATRICES

For two Bott matrices A and A' in \mathcal{M}_n , we say that A and A' are *isomorphic* if $\mathcal{B}(A)$ and $\mathcal{B}(A')$ are isomorphic as toric varieties. Equivalently, A and A' are isomorphic if the corresponding fans Σ_A and $\Sigma_{A'}$ are unimodularly equivalent, i.e., there is a \mathbb{Z} -linear map in $\mathrm{GL}(n, \mathbb{Z})$ which sends a maximal cone in Σ_A to a maximal cone in $\Sigma_{A'}$. For a given Bott matrix A , there are two natural ways of producing (possibly new) isomorphic Bott matrices as follows.

For $A \in \mathcal{M}_n$ and $I \subset [n]$, we consider the $n \times n$ lower triangular matrix L_I whose j -th column \mathbf{c}_j is defined by

$$\mathbf{c}_j := \begin{cases} \mathbf{e}_j & \text{if } j \in I, \\ \mathbf{v}_j & \text{if } j \in I^c, \end{cases}$$

where \mathbf{v}_j is the j -th column of A .

Proposition 3.1. *For $A \in \mathcal{M}_n$ and $I \subset [n]$, the matrix*

$$A_I := L_I^{-1} \cdot L_{I^c}$$

is also a Bott matrix. Moreover, A and A_I are isomorphic. We denote the operation $A \mapsto A_I$ by “Op.1”.

Proof. Observe that A_I is a lower triangular integer matrix and

$$(A_I)_{ii} = \sum_{j=1}^n (L_I^{-1})_{ij} (L_{I^c})_{ji} = (L_I^{-1})_{ii} (L_{I^c})_{ii} = -1$$

since $(L_I^{-1})_{ij} = 0$ for $j > i$ and $(L_{I^c})_{ji} = 0$ for $i > j$. Thus the first claim easily follows.

For the latter statement, consider a \mathbb{Z} -linear map given by $L_I^{-1} \in \text{GL}(n, \mathbb{Z})$. Then it induces a map

$$\Sigma_A \rightarrow \Sigma_{A_I}$$

which sends each maximal cone $\sigma_J \in \Sigma_A$ to $L_I^{-1}\sigma_J \in \Sigma_{A_I}$ for each $J \subset [n]$ (in particular σ_I to the first quadrant). Therefore, two fans Σ_A and Σ_{A_I} are unimodularly equivalent. \square

Remark 3.2. The column vectors of L_I are the ray generators of the maximal cone σ_I in (2.4). Therefore, the operation **Op.1** is nothing but a procedure of selecting a *reference cone* σ_I and sending it to the *first quadrant* by $L_I^{-1} \in \text{GL}(n, \mathbb{Z})$. Accordingly, if we take a Bott matrix A which defines a *Fano* Bott manifold, all the matrices A_I obtained by the operation **Op.1** define *Fano* Bott manifolds because *Fano* is an intrinsic property of a toric variety.

Depending on A , it could happen that a reordering (by some permutation $\pi \in S_n$) of the standard basis $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ changes A into another Bott matrix (denoted by A_π) by rearranging its column vectors. Equivalently, $A_\pi = P_\pi A P_\pi^{-1}$, where P_π is the row permutation matrix of π , that is, P_π has 1 on $(i, \pi(i))$ -entry for $i = 1, \dots, n$ and all the other entries are zero. We call the operation $A \mapsto A_\pi$ “**Op.2**” when A_π is still a Bott matrix. It is straightforward that A and A_π are isomorphic since Σ_A and Σ_{A_π} are the same up to reordering coordinates.

Example 3.3. Consider $A = \begin{bmatrix} -1 & 0 & 0 \\ 1 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$.

(1) (**Op.1**) For $I = \{1\} \subset [3]$, we have

$$L_I = L_I^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \quad L_{I^c} = \begin{bmatrix} -1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad A_I = \begin{bmatrix} -1 & 0 & 0 \\ -1 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}.$$

(2) (**Op.2**) For $\pi = (2, 3)$, we have

$$A_{(2,3)} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 1 & 0 & -1 \end{bmatrix}.$$

The following proposition tells us that for a given Bott matrix A , all Bott matrices isomorphic to A are produced by applying **Op.1** and **Op.2** to A .

Proposition 3.4. *Two Bott matrices are isomorphic if and only if one is obtained from the other by applying two operations **Op.1** and **Op.2**.*

Proof. The “if” part is straightforward. Thus we only need to prove the “only if” part.

Suppose that A and B are isomorphic Bott matrices, i.e., there exists a \mathbb{Z} -linear map $C \in \text{GL}(n, \mathbb{Z})$ sending maximal cones in Σ_A to maximal cones in Σ_B . The \mathbb{Z} -linear map C sends $\sigma_{[n]}^A \in \Sigma_A$ to $\sigma_I^B \in \Sigma_B$ for some $I \subset [n]$. (We denote by σ_I^A the maximal cone in Σ_A given by $I \subset [n]$.) Since B and $B_I := (L_I^B)^{-1} \cdot L_{I^c}^B$ are isomorphic, we may think of $(L_I^B)^{-1} \cdot C$ as a \mathbb{Z} -linear

map which sends $\sigma_{[n]}^A \in \Sigma_A$ to $\sigma_{[n]}^{B_I} \in \Sigma_{B_I}$. In other words, $(L_I^B)^{-1} \cdot C$ is a reordering of the standard basis and hence it corresponds to **Op.2**. Therefore, B_I is obtained from A by **Op.2**. Accordingly, B is given by A applying **Op.2** and **Op.1** in order and this completes the proof. \square

Note that the rational cohomology class y_i^A given in (2.6) depends on the Bott matrix A . In the rest of the section, we will show that if A and A' are isomorphic, then there is a graded ring isomorphism

$$(3.1) \quad \varphi: H^*(\mathcal{B}(A); \mathbb{Z}) \rightarrow H^*(\mathcal{B}(A'); \mathbb{Z})$$

sending the set $\{2y_i^A\}_{i=1}^n$ to $\{2y_i^{A'}\}_{i=1}^n$. This fact can be obtained as a byproduct of Propositions 3.4, 3.5, and 3.6.

Proposition 3.5. *Let $A \in \mathcal{M}_n$. For $I \subset [n]$, there exists a c_1 -preserving graded ring isomorphism*

$$\varphi_I: H^*(\mathcal{B}(A); \mathbb{Z}) \rightarrow H^*(\mathcal{B}(A_I); \mathbb{Z})$$

such that $\varphi_I(2y_i^A) = 2y_i^{A_I}$ for $i = 1, \dots, n$.

Proof. We first consider the case where $I = \{k\}^c = [n] \setminus \{k\}$ for some $k \in [n]$ so that $A_I = L_{\{k\}^c}^{-1} L_{\{k\}}$. Let \mathbf{a}_i and \mathbf{a}_i^I be the i -th row of $A + E$ and $A_I + E$, respectively. By direct computations, we obtain

- (1) $\mathbf{a}_i^I = \mathbf{a}_i$ for $i < k$;
- (2) $\mathbf{a}_i^I = -\mathbf{a}_k$ for $i = k$; and
- (3) $\mathbf{a}_i^I = \mathbf{a}_i + a_{i,k} \mathbf{a}_k$ for $i > k$.

Then the Bott tower \mathcal{B}_\bullet^I corresponding to A_I is given by

$$\mathcal{B}_i^I = \begin{cases} \mathcal{B}_i & \text{for } i < k, \\ P(\mathbb{C} \oplus \xi_k^{-1}) & \text{for } i = k, \\ P(\mathbb{C} \oplus (\xi_i \otimes \xi_k^{a_{i,k}})) & \text{for } i > k. \end{cases}$$

Here, ξ_1, \dots, ξ_n are the line bundles used to construct the Bott tower $\mathcal{B}_\bullet = \mathcal{B}(A)$.

We define a map φ_I by

$$(3.2) \quad \varphi_I(x_i^A) = \begin{cases} x_i^{A_I} & \text{for } i \neq k, \\ x_k^{A_I} + \sum_{j < k} a_{k,j} x_j^{A_I} & \text{for } i = k. \end{cases}$$

Then, by (2), we obtain that

$$(3.3) \quad \varphi_I(x_k^A) = x_k^{A_I} - \alpha_k^{A_I} \quad \text{and} \quad \varphi_I(\alpha_k^A) = -\alpha_k^{A_I}.$$

We claim that φ_I is well-defined and is indeed a graded ring isomorphism such that $\varphi_I(2y_i^A) = 2y_i^{A_I}$ for every $i = 1, \dots, n$. The well-definedness follows by showing that $\varphi_I(x_i^A(x_i^A - \alpha_i^A)) = 0$ for all i .

- For $i < k$, we get $\varphi_I(x_i^A(x_i^A - \alpha_i^A)) = x_i^{A_I}(x_i^{A_I} - \alpha_i^{A_I}) = 0$;
- for $i = k$, by (3.3), we have

$$\varphi_I(x_k^A(x_k^A - \alpha_k^A)) = (x_k^{A_I} - \alpha_k^{A_I})(x_k^{A_I} - \alpha_k^{A_I} - \varphi_I(\alpha_k^A)) = x_k^{A_I}(x_k^{A_I} - \alpha_k^{A_I}) = 0;$$

- for $i > k$, we have $\varphi_I(x_i^A(x_i^A - \alpha_i^A)) = x_i^{A_I}(x_i^{A_I} - \varphi_I(\alpha_i^A)) = x_i^{A_I}(x_i^{A_I} - \alpha_i^{A_I}) = 0$;

where the second last equality is obtained from (3.2) and (3):

$$(3.4) \quad \varphi_I(\alpha_i^A) = \sum_{j < i} a_{i,j} \varphi_I(x_j^A) = \sum_{j \neq k, j < i} a_{i,j} x_j^{A_I} + a_{i,k} \left(x_k^{A_I} + \sum_{j < k} a_{k,j} x_j^{A_I} \right) = \alpha_i^{A_I} \quad \text{for } i > k.$$

To show $\varphi_I(2y_i^A) = 2y_i^{A_I}$, we only need to check the case when $i = k$ because $\varphi_I(x_i^A) = x_i^{A_I}$ and $\varphi_I(\alpha_i^A) = \alpha_i^{A_I}$ for $i \neq k$ by (3.2) and (3.4). Then, by (3.3), we obtain

$$\varphi_I(2y_k^A) = \varphi_I(2x_k^A - \alpha_k^A) = 2(x_k^{A_I} - \alpha_k^{A_I}) + \alpha_k^{A_I} = 2x_k^{A_I} - \alpha_k^{A_I} = 2y_k^{A_I}$$

and this completes the proof for the case of $I = \{k\}^c$.

For a general $I = \{i_1 < \dots < i_m\} \subset [n]$, using the fact $L_I = L_{\{i_1\}^c} \cdot L_{\{i_2\}^c} \cdots L_{\{i_m\}^c}$, we obtain

$$\varphi_I = \varphi_{\{i_1\}^c} \circ \varphi_{\{i_2\}^c} \circ \dots \circ \varphi_{\{i_m\}^c}.$$

Applying the previous procedure repeatedly, the result follows. One can immediately check that φ_I is c_1 -preserving by (2.6). \square

For the cohomology ring isomorphism between Bott manifolds induced from **Op.2**, we recall the result [CMM15].

Proposition 3.6 ([CMM15, Lemma 6.1]). *Let $A \in \mathcal{M}_n$. For a permutation π on $[n]$, if $A_\pi \in \mathcal{M}_n$, then there is a c_1 -preserving graded ring isomorphism*

$$\varphi_\pi: H^*(\mathcal{B}(A); \mathbb{Z}) \rightarrow H^*(\mathcal{B}(A_\pi); \mathbb{Z})$$

such that $\varphi_\pi(x_i^A) = x_{\pi(i)}^{A_\pi}$ for $i = 1, \dots, n$. Indeed, we have $\varphi_\pi(2y_i^A) = 2y_{\pi(i)}^{A_\pi}$ for $i = 1, \dots, n$.

4. MAIN THEOREM

In this section, we will prove Theorem 1.2. Throughout this section, every cohomology ring will take coefficient in \mathbb{Z} unless otherwise stated.

Before to begin with, we explain some notations used in this section. We use letters \mathcal{B} , \mathcal{B}' and \mathcal{B}'' to indicate Bott manifolds. And also we denote by (x_i, y_i, α_i) , (x'_i, y'_i, α'_i) , and $(x''_i, y''_i, \alpha''_i)$ the elements x , y , and α defined in (2.5) and (2.6) for \mathcal{B} , \mathcal{B}' , and \mathcal{B}'' , respectively.

Lemma 4.1. *Let \mathcal{B} and \mathcal{B}' be Fano Bott manifolds. If there is a c_1 -preserving graded ring isomorphism $\varphi: H^*(\mathcal{B}) \rightarrow H^*(\mathcal{B}')$, then there exists a Fano Bott manifold \mathcal{B}'' together with a c_1 -preserving graded ring isomorphism $\psi: H^*(\mathcal{B}) \rightarrow H^*(\mathcal{B}'')$ such that \mathcal{B}' and \mathcal{B}'' are isomorphic and $\psi(x_1) = x''_1$. In particular, we have $\psi(2y_1) = 2y''_1$.*

Proof. Suppose that Fano Bott manifolds \mathcal{B} and \mathcal{B}' are determined by Bott matrices $A = (a_{ij})$ and $A' = (a'_{ij})$, respectively. By Theorem 2.6, there exists a permutation σ on $[n]$ such that $\varphi(2y_i) = 2y'_{\sigma(i)}$ for each $i = 1, \dots, n$. If $\sigma(1) \neq 1$, then we get $a'_{\sigma(1),j} = 0$ for every $j < \sigma(1)$. Indeed, since $2y_1 = 2x_1$ and $2y'_{\sigma(1)} = 2x'_{\sigma(1)} - \sum_{j < \sigma(1)} a'_{\sigma(1),j} x'_j$, we get

$$2(\varphi(x_1) - x'_{\sigma(1)}) = - \sum_{j < \sigma(1)} a'_{\sigma(1),j} x'_j$$

which is divisible by 2. Since $a'_{\sigma(1),j}$ belongs to $\{0, \pm 1\}$, we conclude that $a'_{\sigma(1),j} = 0$ for every $j < \sigma(1)$ and $\varphi(x_1) = x'_{\sigma(1)}$. This fact tells us that the $\sigma(1)$ -th row of $A' + E$ is zero, and therefore we may apply **Op.2** to A' for the permutation $\pi = s_1 s_2 \dots s_{\sigma(1)-1}$ where s_i denotes the simple transposition $(i, i+1)$. Here, we note that $(\pi \circ \sigma)(1) = 1$.

We consider a Bott matrix $A'' := A'_\pi$ and let \mathcal{B}'' be a Bott manifold associated to A'' . Suppose that \mathcal{B}'' is Fano. Then \mathcal{B}' and \mathcal{B}'' are isomorphic and there is a graded ring isomorphism $\varphi': H^*(\mathcal{B}') \rightarrow H^*(\mathcal{B}'')$ such that $\varphi'(x'_{\sigma(1)}) = x''_{\pi(\sigma(1))} = x''_1$ by Proposition 3.6. Then $\varphi' \circ \varphi$ is the desired isomorphism and it completes the proof. \square

Theorem 4.2 (Theorem 1.2). *Let \mathcal{B} and \mathcal{B}' be Fano Bott manifolds. Assume that there is a c_1 -preserving graded ring isomorphism $\varphi: H^*(\mathcal{B}) \rightarrow H^*(\mathcal{B}')$. Then \mathcal{B} and \mathcal{B}' are isomorphic as toric varieties.*

Proof. Suppose that Fano Bott manifolds \mathcal{B} and \mathcal{B}' are determined by Bott matrices $A = (a_{ij})$ and $A' = (a'_{ij})$, respectively. From Theorem 2.6, there is a permutation σ on $[n]$ such that $\varphi(2y_i) = 2y'_{\sigma(i)}$ for all $i = 1, \dots, n$. We may further assume that $\varphi(x_1) = x'_1$ (or equivalently $\varphi(2y_1) = 2y'_1$) by Lemma 4.1, i.e., $\sigma(1) = 1$.

Now we choose an index $k \in [n]$ (with $k > 1$) such that $\varphi(x_i) = x'_i$ for every $i < k$ and $\varphi(x_k) \neq x'_k$, that is, $\varphi(2y_i) = 2y'_i$ for all $i < k$. Then there are two possibilities:

- (1) $\sigma(k) = k$; or
- (2) $\sigma(k) > k$.

We first consider the case where $\sigma(k) = k$. That is, $\varphi(2y_k) = 2y'_k$ and $\varphi(x_k) \neq x'_k$. Since $\varphi(2y_k) = \varphi(2x_k - \alpha_k) = 2\varphi(x_k) - \varphi(\alpha_k)$ and $\varphi(2y_k) = 2y'_k = 2x'_k - \alpha'_k$, we get

$$(4.1) \quad 2(\varphi(x_k) - x'_k) = \varphi(\alpha_k) - \alpha'_k$$

and hence $\varphi(\alpha_k) - \alpha'_k$ is divisible by 2. Note that $\varphi(\alpha_k) - \alpha'_k = \sum_{j < k} (a_{k,j} - a'_{k,j})x'_j$ by the definition of α_i and α'_i in (2.5) and from the induction hypothesis. Comparing this with the equation (4.1), we have $\varphi(x_k) = x'_k + \sum_{j < k} c_j x'_j$, where

$$(4.2) \quad 2c_j = a_{k,j} - a'_{k,j} \quad \text{for } j = 1, \dots, k-1$$

which implies that

$$(4.3) \quad c_j = a_{k,j} = -a'_{k,j} \quad \text{when } c_j \neq 0$$

by the Fano condition in Theorem 2.9. Note that

$$(4.4) \quad \text{not every } c_j \text{ is zero for } j < k \text{ by our assumption.}$$

Now we claim that the equation (4.3) holds even when $c_j = 0$. By the well-definedness of the isomorphism φ , we have that

$$\begin{aligned} 0 = \varphi(x_k(x_k - \alpha_k)) &= \left(x'_k + \sum_{j < k} c_j x'_j \right) \left(x'_k + \sum_{j < k} c_j x'_j - \sum_{j < k} a_{k,j} x'_j \right) \\ &= (x'_k)^2 + x'_k \left(2 \sum_{j < k} c_j x'_j - \sum_{j < k} a_{k,j} x'_j \right) + \left(\sum_{j < k} c_j x'_j \right) \left(\sum_{j < k} c_j x'_j - \sum_{j < k} a_{k,j} x'_j \right) \\ &= (x'_k)^2 + x'_k \sum_{j < k} (2c_j - a_{k,j}) x'_j + \left(\sum_{j < k} c_j x'_j \right) \left(\sum_{j < k} (c_j - a_{k,j}) x'_j \right) \end{aligned}$$

in $H^*(\mathcal{B}')$. Since

- there is no term having x'_k in $\left(\sum_{j < k} c_j x'_j \right) \left(\sum_{j < k} (c_j - a_{k,j}) x'_j \right)$, and
- $(x'_k)^2 + x'_k \sum_{j < k} (2c_j - a_{k,j}) x'_j$ can be expressed as a linear combination of the linearly independent set $\{x'_k x'_j\}_{j < k}$,

both $(x'_k)^2 + x'_k \sum_{j < k} (2c_j - a_{k,j}) x'_j$ and $\left(\sum_{j < k} c_j x'_j \right) \left(\sum_{j < k} (c_j - a_{k,j}) x'_j \right)$ vanish in $H^*(\mathcal{B}')$. In particular, it follows from the vanishing of the latter term above that

$$0 = \left(\sum_{j < k} c_j x'_j \right) \left(\sum_{j < k} (c_j - a_{k,j}) x'_j \right) = \left(\sum_{\substack{j < k \\ c_j \neq 0}} c_j x'_j \right) \left(- \sum_{\substack{\ell < k \\ c_\ell = 0}} a_{k,\ell} x'_\ell \right) = - \sum_{\substack{j, \ell < k \\ c_j \neq 0 \text{ and } c_\ell = 0}} c_j a_{k,\ell} x'_j x'_\ell$$

in $H^*(\mathcal{B})$, where the second equality follows from (4.3). However, $\{x'_j x'_\ell \mid c_j \neq 0, c_\ell = 0, j, \ell < k\}$ is linearly independent by Remark 2.4 (2), so $a_{k,\ell} = 0$ if $c_\ell = 0$. Moreover, since $2c_j = a_{k,j} - a'_{k,j}$ by (4.2), we conclude that

- if $c_j = 0$, then $a_{k,j} = a'_{k,j} = 0$, and
- if $c_j \neq 0$, then $a_{k,j} = -a'_{k,j}$ by (4.3).

Consequently, we have $c_j = a_{k,j} = -a'_{k,j}$ for every $j < k$. Therefore, we obtain

$$\varphi(x_k) = x'_k + \sum_{j < k} c_j x'_j = x'_k - \sum_{j < k} a'_{k,j} x'_j = x'_k - \alpha'_k.$$

Now, we let \mathcal{B}'' be a Fano Bott manifold whose Bott matrix A'' is obtained from A' by applying **Op.1** with $I = \{k\}^c$. Then there is a c_1 -preserving graded ring isomorphism $\varphi': H^*(\mathcal{B}') \rightarrow H^*(\mathcal{B}'')$ such that $\varphi'(x'_j) = x''_j$ for every $j \neq k$ and $\varphi'(x'_k) = x''_k - \alpha''_k$ by (3.2) and (3.3), and also $\varphi'(2y'_i) = 2y''_i$ for all i by Proposition 3.5. Then, the composition $\psi := \varphi' \circ \varphi$ is a graded ring isomorphism $\psi: H^*(\mathcal{B}) \rightarrow H^*(\mathcal{B}'')$ such that $\psi(2y_i) = 2y''_i$ for every $i = 1, \dots, n$, \mathcal{B}' and \mathcal{B}'' are isomorphic as toric varieties, and $\psi(x_j) = x''_j$ for every $j \leq k$. Indeed, by (3.3), we have

$$\psi(x_k) = \varphi'(x'_k - \alpha'_k) = x''_k - \alpha''_k + \alpha''_k = x''_k.$$

Now, we consider the second case, $\sigma(k) > k$. Since $\varphi(2y_k) = 2\varphi(x_k) - \sum_{j < k} a_{k,j}x'_j$ and $\varphi(2y_k) = 2y'_{\sigma(k)}$, we get

$$(4.5) \quad 2(\varphi(x_k) - x'_{\sigma(k)}) = \sum_{j < k} a_{k,j}x'_j - \sum_{\ell < \sigma(k)} a'_{\sigma(k),\ell}x'_\ell.$$

Because $a_{k,j}$ and $a_{\sigma(k),\ell}$ belong to $\{0, 1, -1\}$ and the left hand side of (4.5) is divisible by 2, we have

$$a'_{\sigma(k),\ell} = 0 \quad \text{for } k \leq \ell < \sigma(k).$$

Indeed, the $\sigma(k)$ -th row of A' has consecutive zeros from $(\sigma(k), k)$ to $(\sigma(k), \sigma(k) - 1)$. Therefore by applying **Op.2** to A' for the permutation $\pi = s_k \cdots s_{\sigma(k)-1}$, we get a new Bott matrix A'' such that $\mathcal{B}(A'')$ is Fano and it is isomorphic to \mathcal{B}' and there is a graded ring isomorphism $\varphi': H^*(\mathcal{B}') \rightarrow H^*(\mathcal{B}'')$ satisfying

$$\varphi'(x'_i) = \begin{cases} x''_i & \text{for } i < k \text{ or } i > \sigma(k), \\ x''_{i+1} & \text{for } k \leq i \leq \sigma(k) - 1, \\ x''_k & \text{for } i = \sigma(k) \end{cases}$$

by Proposition 3.6. Since $\pi \circ \sigma(i) = i$ for every $i \leq k$, the composition $\varphi' \circ \varphi$ is a c_1 -preserving graded ring isomorphism $\varphi' \circ \varphi: H^*(\mathcal{B}) \rightarrow H^*(\mathcal{B}'')$ satisfying

$$\varphi' \circ \varphi(x_i) = x''_i \quad (i < k) \quad \text{and} \quad \varphi' \circ \varphi(2y_k) = \varphi'(2y'_{\sigma(k)}) = 2y_{\pi \circ \sigma(k)} = 2y''_k.$$

Hence this case reduces to the first case.

We may repeat the above argument as many times as necessary. Since the indices are bounded above by n , this process must stop. Therefore, we have a Fano Bott manifold $\tilde{\mathcal{B}}$ isomorphic to \mathcal{B}' whose Bott matrix is exactly the same as that of \mathcal{B} , i.e., \mathcal{B} and \mathcal{B}' are isomorphic. This finishes the proof. \square

Remark 4.3. We cannot extend Theorem 1.2 to weak Fano Bott manifolds. Note that the Hirzerbuch surfaces \mathcal{H}_0 and \mathcal{H}_2 are weak Fano Bott manifolds. As we saw in Example 2.8, \mathcal{H}_0 and \mathcal{H}_2 are not isomorphic but there is a c_1 -preserving graded cohomology ring isomorphism between them.

Remark 4.4. One may wonder whether we can extend Theorem 1.2 to Bott manifolds whose Bott matrices have entries 0, 1, or -1 . However, the set of such Bott matrices is not closed under the operation **Op.1**. For example, consider a matrix

$$A = \begin{bmatrix} -1 & 0 & 0 \\ 1 & -1 & 0 \\ 1 & 1 & -1 \end{bmatrix}.$$

Then, we have that

$$A_\emptyset = A^{-1} = \begin{bmatrix} -1 & 0 & 0 \\ -1 & -1 & 0 \\ -2 & -1 & -1 \end{bmatrix}$$

whose entry has $-2 \notin \{0, 1, -1\}$. Note that the set of Bott matrices obtained from Fano Bott manifolds is closed under the operation **Op.1** as we mentioned in Remark 3.2.

Remark 4.5. For $n = 3$, there are five Bott matrices associated to Fano Bott manifolds up to isomorphisms.

Oda's list	6	7	8	9	10
Øbro's list	21	11	18	17	12
Bott matrix	$\begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$	$\begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 1 & 1 & -1 \end{bmatrix}$	$\begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 1 & -1 & -1 \end{bmatrix}$	$\begin{bmatrix} -1 & 0 & 0 \\ 1 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$	$\begin{bmatrix} -1 & 0 & 0 \\ 1 & -1 & 0 \\ 0 & 1 & -1 \end{bmatrix}$

Here, we refer the lists provided by Oda [Oda88, Figure 2.6] and Øbro [Øbr07] in the Graded Ring Database [BK]. However, the Bott manifolds corresponding to the second and the third matrices are diffeomorphic since their cohomology rings are isomorphic as graded rings. Indeed, the degrees of the corresponding Bott manifolds are different, so there does not exist c_1 -preserving cohomology ring isomorphism between them. Note that the family of Bott manifolds of complex dimension at most 4 is cohomologically rigid. See [CMS10, Cho15].

REFERENCES

- [BEM⁺17] Victor M. Buchstaber, Nikolay Yu. Erokhovets, Mikiya Masuda, Taras E. Panov, and Seonjeong Park, *Cohomological rigidity of manifolds defined by 3-dimensional polytopes*, Uspekhi Mat. Nauk **72** (2017), no. 2(434), 3–66.
- [Ber03] Florian Berchtold, *Lifting of morphisms to quotient presentations*, Manuscripta Math. **110** (2003), no. 1, 33–44.
- [BH58] Armand Borel and Friedrich Hirzebruch, *Characteristic classes and homogeneous spaces. I*, Amer. J. Math. **80** (1958), 458–538.
- [BK] Gavin Brown and Alexander Kasprzyk, *Graded ring database*, <http://www.grdb.co.uk/forms/toricsmooth>.
- [BP15] Victor M. Buchstaber and Taras E. Panov, *Toric topology*, Mathematical Surveys and Monographs, vol. 204, American Mathematical Society, Providence, RI, 2015.
- [Cho15] Suyoung Choi, *Classification of Bott manifolds up to dimension 8*, Proc. Edinb. Math. Soc. (2) **58** (2015), no. 3, 653–659.
- [Civ05] Yusuf Civan, *Bott towers, crosspolytopes and torus actions*, Geom. Dedicata **113** (2005), 55–74.
- [CLMP20] Yunhyung Cho, Eunjeong Lee, Mikiya Masuda, and Seonjeong Park, *On Fano generalized Bott manifolds*, in preparation.
- [CMM15] Suyoung Choi, Mikiya Masuda, and Satoshi Murai, *Invariance of Pontrjagin classes for Bott manifolds*, Algebr. Geom. Topol. **15** (2015), no. 2, 965–986.
- [CMS10] Suyoung Choi, Mikiya Masuda, and Dong Youp Suh, *Topological classification of generalized Bott towers*, Trans. Amer. Math. Soc. **362** (2010), no. 2, 1097–1112.
- [CMS11] ———, *Rigidity problems in toric topology: a survey*, Tr. Mat. Inst. Steklova **275** (2011), 188–201.
- [Del88] Thomas Delzant, *Hamiltoniens périodiques et images convexes de l'application moment*, Bull. Soc. Math. France **116** (1988), no. 3, 315–339.
- [Fan14] Andrew Fanoe, *Toric structures on bundles of projective spaces*, J. Symplectic Geom. **12** (2014), no. 4, 685–724.
- [GK94] Michael Grossberg and Yael Karshon, *Bott towers, complete integrability, and the extended character of representations*, Duke Math. J. **76** (1994), no. 1, 23–58.
- [Har77] Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York-Heidelberg, 1977, Graduate Texts in Mathematics, No. 52.
- [HKM20] Akihiro Higashitani, Kazuki Kurimoto, and Mikiya Masuda, *Cohomological rigidity for toric Fano manifolds of small dimensions or large Picard numbers*, in preparation.

- [KKP07] Yael Karshon, Liat Kessler, and Martin Pinsonnault, *A compact symplectic four-manifold admits only finitely many inequivalent toric actions*, J. Symplectic Geom. **5** (2007), no. 2, 139–166.
- [Kle66] Steven L. Kleiman, *Toward a numerical theory of ampleness*, Ann. of Math. (2) **84** (1966), 293–344.
- [McD11] Dusa McDuff, *The topology of toric symplectic manifolds*, Geom. Topol. **15** (2011), no. 1, 145–190.
- [MP08] Mikiya Masuda and Taras E. Panov, *Semi-free circle actions, Bott towers, and quasitoric manifolds*, Mat. Sb. **199** (2008), no. 8, 95–122.
- [MS08] Mikiya Masuda and Dong Youp Suh, *Classification problems of toric manifolds via topology*, Toric topology, Contemp. Math., vol. 460, Amer. Math. Soc., Providence, RI, 2008, pp. 273–286.
- [Obr07] Mikkel Øbro, *An algorithm for the classification of smooth Fano polytopes*, preprint, arXiv:0704.0049 (2007).
- [Oda78] Tadao Oda, *Torus embeddings and applications*, Tata Institute of Fundamental Research Lectures on Mathematics and Physics, vol. 57, Tata Institute of Fundamental Research, Bombay; by Springer-Verlag, Berlin-New York, 1978, Based on joint work with Katsuya Miyake.
- [Oda88] ———, *Convex bodies and algebraic geometry*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 15, Springer-Verlag, Berlin, 1988, An introduction to the theory of toric varieties, Translated from the Japanese.
- [Pin08] Martin Pinsonnault, *Maximal compact tori in the Hamiltonian group of 4-dimensional symplectic manifolds*, J. Mod. Dyn. **2** (2008), no. 3, 431–455.
- [PT20] Milena Pabiniak and Susan Tolman, *Symplectic cohomological rigidity via toric degenerations*, preprint, arXiv:2002.12434 (2020).
- [Suy19] Yusuke Suyama, *Fano generalized Bott manifolds*, Manuscripta Mathematica (2019).

DEPARTMENT OF MATHEMATICS EDUCATION, SUNGKYUNKWAN UNIVERSITY, SEOUL, REPUBLIC OF KOREA

E-mail address: `yunhyung@skku.edu`

CENTER FOR GEOMETRY AND PHYSICS, INSTITUTE FOR BASIC SCIENCE (IBS), POHANG, REPUBLIC OF KOREA

E-mail address: `eunjeong.lee@ibs.re.kr`

OSAKA CITY UNIVERSITY ADVANCED MATHEMATICS INSTITUTE (OCAMI) & DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA CITY UNIVERSITY, SUMIYOSHIKU, SUGIMOTO, 558-8585, OSAKA, JAPAN

E-mail address: `masuda@osaka-cu.ac.jp`

DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, DAEJEON, REPUBLIC OF KOREA

E-mail address: `seonjeong1124@gmail.com`