

A CLASSIFICATION OF LEFT-INVARIANT SYMPLECTIC STRUCTURES ON SOME LIE GROUPS

メタデータ	言語: English 出版者: OCAMI 公開日: 2022-06-06 キーワード (Ja): キーワード (En): Moduli space, Left-invariant symplectic structures, Symplectic Lie group, Symplectic QR decomposition 作成者: Castellanos Moscoso, Luis Pedro, 田丸, 博士 メールアドレス: 所属: Osaka City University, Osaka City University
URL	https://ocu-omu.repo.nii.ac.jp/records/2016907

A CLASSIFICATION OF LEFT-INVARIANT SYMPLECTIC STRUCTURES ON SOME LIE GROUPS

LUIS PEDRO CASTELLANOS MOSCOSO

AND HIROSHI TAMARU

Citation	OCAMI Preprint Series. 2020, 20-19
Issue Date	2020-12-18
Type	Preprint
Textversion	Author
Rights	For personal use only. No other uses without permission.
Relation	This is a pre-print of an article published in Beiträge zur Algebra und Geometrie / Contributions to Algebra and Geometry. The final authenticated version is available online at: https://doi.org/10.1007/s13366-022-00643-1 .

From: Osaka City University Advanced Mathematical Institute

<http://www.sci.osaka-cu.ac.jp/OCAMI/publication/preprint/preprint.html>

A CLASSIFICATION OF LEFT-INVARIANT SYMPLECTIC STRUCTURES ON SOME LIE GROUPS

LUIS PEDRO CASTELLANOS MOSCOSO AND HIROSHI TAMARU

ABSTRACT. We are interested in the classification of left-invariant symplectic structures on Lie groups. Some classifications are known, especially in low dimensions. In this paper we establish a new approach to classify (up to automorphism and scale) left-invariant symplectic structures on Lie groups. The procedure is based on the moduli space of left-invariant nondegenerate 2-forms. Then we apply our procedure for two particular Lie groups of dimension $2n$ and give classifications of left-invariant symplectic structures on them.

1. INTRODUCTION

In geometry it is an important problem to study whether a given manifold admits some nice geometric structures. In the setting of Lie groups, it is natural to ask about the existence of left-invariant structures. A *symplectic Lie group* is a Lie group G endowed with a left-invariant symplectic form ω (that is, a nondegenerate closed 2-form). The geometry of symplectic Lie groups is an active field of research. There are many interesting results on the structure of symplectic Lie groups and some classifications in low dimensions, but the general picture is far from complete. Conjectures about the existence of isotropic subgroups are still unsolved. Some nice known results include

- (1) Unimodular symplectic Lie groups are solvable. All symplectic Lie groups of dimension 4 are solvable ([5]).
- (2) Some of the known classification in low dimensions include: complete classification for the 4-dimensional case ([14]), filiform Lie algebras up to dimension 10 ([6]), most of solvable Lie algebras up to dimension 6 ([12],[4]).
- (3) Completely solvable symplectic Lie groups have Lagrangian subgroups. If dimension ≤ 6 , every symplectic Lie group has a Lagrangian subgroup. All nilpotent symplectic Lie groups of dimension ≤ 6 have a Lagrangian normal subgroup ([1]).

In [7] we can find a novel method to find nice (e.g., Einstein or Ricci soliton) left-invariant Riemannian metrics. The method is based on the moduli space of left-invariant Riemannian metrics on a Lie group G . Recall that there is a correspondence between left-invariant metrics on G and inner products on its Lie algebra \mathfrak{g} . The moduli space is defined as the orbit space of the action of $\mathbb{R}^\times \text{Aut}(\mathfrak{g})$ on the space $\mathfrak{M}(\mathfrak{g})$ of inner products on \mathfrak{g} . An expression of the moduli space derives a *Milnor-type theorem*. Milnor-type theorems, first introduced in the aforementioned paper, can be thought as a generalization of so called Milnor frames introduced by Milnor in his famous paper [13]. In the setting of Riemannian metrics a Milnor-type theorem gives an orthonormal basis for each inner product $\langle \cdot, \cdot \rangle$ in \mathfrak{g} , such that the bracket relations between the elements of the basis are given in terms of a small

This work was partly supported by Osaka City University Advanced Mathematical Institute (MEXT Joint Usage/Research Center on Mathematics and Theoretical Physics JPMXP0619217849). The second author was supported by JSPS KAKENHI Grant Number 19K21831.

number of parameters. Therefore they can be used, for example, to calculate curvatures in a relatively easy way. In [10] the same ideas are used successfully in the pseudo-Riemannian case.

It would be then natural to try to use the same ideas for symplectic Lie groups. In a similar way as before, the study of symplectic Lie groups reduces to the study of *symplectic Lie algebras* (\mathfrak{g}, ω) , that is Lie algebras \mathfrak{g} endowed with nondegenerate closed 2-forms (or equivalently two-cocycles $\omega \in Z^2(\mathfrak{g})$). We study the moduli space of left-invariant nondegenerate 2-forms, which is the orbit space of the action of $\mathbb{R}^\times \text{Aut}(\mathfrak{g})$ on the space $\Omega(\mathfrak{g})$ of nondegenerate 2-forms on \mathfrak{g} . Then we derive a procedure to obtain Milnor-type theorems. In the setting of symplectic Lie groups a Milnor-type theorem gives a symplectic basis for each nondegenerate 2-form in \mathfrak{g} , such that the bracket relations between the elements of the basis are given in terms of a small number of parameters. We show how these theorems can be used to search for 2-forms that are closed. We also use our method to show the existence of some particular subalgebras. These ideas are developed in Section 3. We would like to emphasize that the approach presented in this paper is different from that in the existing literature. In addition, the method can be used, at least theoretically, for any Lie algebra.

As a first application of this method, in Section 5, we study two particular Lie algebras: the Lie algebra of the real hyperbolic space $\mathfrak{g}_{\mathbb{R}H^{2n}}$, and the direct sum of the 3-dimensional Heisenberg Lie algebra and the abelian Lie algebra $\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3}$. These Lie algebras have particularly big automorphism groups: they are parabolic subgroups of $\text{GL}(2n, \mathbb{R})$. Therefore, we expect the corresponding moduli spaces to be small, which is a reason of our choice of these Lie algebras. In fact, in [15] Wolf proved that for the actions of parabolic subgroups on symmetric spaces of reductive type (which is our case), the number of orbits is finite. For the first Lie algebra, we will show in Section 5.1 that the action considered is transitive, and thus the moduli space consists of just one element. Correspondingly, we obtain that there is only one symplectic structure when $n = 1$ and no symplectic structure if $n > 1$. For the second Lie algebra we will show in Section 5.2 that the action considered has at most five orbits (depending on n). Correspondingly, we obtain that for the second Lie algebra there is only one symplectic structure for all $n > 0$. Furthermore we show that the second Lie algebra admits Lagrangian ideals for any $n > 0$.

As a tool for studying the moduli space we also obtain in Section 4 a slight modification of a decomposition theorem of symplectic matrices called symplectic QR decomposition.

The authors would like to thank Takayuki Okuda, Akira Kubo, Yuichiro Takeuchi, Kaname Hashimoto, Yuji Kondo and Masahiro Kawamata for helpful comments.

2. PRELIMINARIES

In this section, we recall some basic notions on left-invariant symplectic forms on Lie groups, and Lagrangian normal subgroups.

2.1. Left-invariant symplectic 2-forms. Let G be a simply connected Lie group with dimension $2n$ and \mathfrak{g} its corresponding Lie algebra. We are interested in the space of left-invariant symplectic forms on G :

$$\left\{ \omega(\cdot, \cdot) \in \bigwedge^2 T^*G \mid \omega^n \neq 0, \text{left-invariant}, d\omega = 0 \right\}.$$

This is a subset of the set of all nondegenerate left-invariant 2-forms, denoted by

$$\Omega(G) := \left\{ \omega(\cdot, \cdot) \in \bigwedge^2 T^*G \mid \omega^n \neq 0, \text{left-invariant} \right\}.$$

For this set we have the following natural equivalence relation.

Definition 2.1. Let $\omega_1, \omega_2 \in \Omega(G)$. Then, (G, ω_1) and (G, ω_2) are said to be *equivalent up to automorphism* (resp. *equivalent up to automorphism and scale*) if there exists $\phi \in \text{Aut}(G)$ such that $\phi^*\omega_1 = \omega_2$ (resp. if there exist $\phi \in \text{Aut}(G)$ and a constant $c \neq 0$ such that $c \cdot (\phi)^*\omega_1 = \omega_2$).

It is well known that the space $\Omega(G)$ can be identified with the space of nondegenerate 2-forms on \mathfrak{g} , denoted by

$$\Omega(\mathfrak{g}) := \left\{ \omega(\cdot, \cdot) \in \bigwedge^2 \mathfrak{g}^* \mid \omega^n \neq 0 \right\}.$$

For this set we have the following natural equivalence relation.

Definition 2.2. Let $\omega_1, \omega_2 \in \Omega(\mathfrak{g})$. Then, (\mathfrak{g}, ω_1) and (\mathfrak{g}, ω_2) are said to be *equivalent up to automorphism* (resp. *equivalent up to automorphism and scale*) if there exists $\phi \in \text{Aut}(\mathfrak{g})$ such that $\phi^*\omega_1 = \omega_2$ (resp. if there exist $\phi \in \text{Aut}(\mathfrak{g})$ and a constant $c \neq 0$ such that $c \cdot (\phi)^*\omega_1 = \omega_2$).

When the Lie group is simply connected both notions Definition 2.1 and Definition 2.2 of equivalence coincide. This fact allows us to work at the Lie algebra level.

Remark 2.3. If (S, ω_1) and (S, ω_2) are symplectic manifolds and there exists $\phi \in \text{Diff}(S)$ such that $\phi^*\omega_1 = \omega_2$, then (S, ω_1) and (S, ω_2) are said to be symplectomorphically equivalent and ϕ is called a symplectomorphism. Notice that the equivalence relation in Definition 2.1 (and the corresponding notion in Definition 2.2) is stronger, but this would be the usual notion of equivalence in symplectic Lie groups. In fact, in the context of symplectic Lie groups, the map in Definition 2.1 or Definition 2.2 is also sometimes called a symplectomorphism.

Remember that a symplectic vector space is a pair (V, ω) , where V is a vector space and ω is a nondegenerate 2-form. For every $\omega_{\mathfrak{g}} \in \Omega(\mathfrak{g})$ the pair $(\mathfrak{g}, \omega_{\mathfrak{g}})$ is a symplectic vector space. The next is a well known fact.

Proposition 2.4. Let $\omega_{\mathfrak{g}} \in \Omega(\mathfrak{g})$, and $\omega_G \in \Omega(G)$ be the corresponding 2-form on the Lie group. Then ω_G is closed if and only if $\omega_{\mathfrak{g}}$ satisfies, for all $x, y, z \in \mathfrak{g}$

$$d\omega_{\mathfrak{g}}(x, y, z) := \omega_{\mathfrak{g}}(x, [y, z]) + \omega_{\mathfrak{g}}(z, [x, y]) + \omega_{\mathfrak{g}}(y, [z, x]) = 0.$$

A 2-form $\omega_{\mathfrak{g}} \in \Omega(\mathfrak{g})$ that satisfies the previous property will be called a *closed 2-form or symplectic form* on the Lie algebra \mathfrak{g} .

Remark 2.5. The previous condition can be expressed in terms of the cohomology of Lie algebras. One knows that $\omega_{\mathfrak{g}}$ is closed if and only if $\omega \in Z^2(\mathfrak{g})$, where $Z^2(\mathfrak{g})$ is the set of 2-cocycles in the trivial representation over \mathbb{R} .

From the theory of homogeneous spaces we also have the identification

$$\Omega(\mathfrak{g}) \cong \text{GL}(2n, \mathbb{R})/\text{Sp}_n(\mathbb{R}).$$

Here we identify $\mathfrak{g} \cong \mathbb{R}^{2n}$, and then the general linear group $\text{GL}(2n, \mathbb{R})$ acts transitively on $\Omega(\mathfrak{g})$ by

$$g.\omega(\cdot, \cdot) = \omega(g^{-1}(\cdot), g^{-1}(\cdot)) \quad \forall g \in \text{GL}(2n, \mathbb{R}).$$

We also recall that $\text{Sp}_n(\mathbb{R})$ is the symplectic group, that is the group of linear maps which preserve the canonical symplectic 2-form ω_0 in \mathbb{R}^{2n} . If $\{e_1, \dots, e_{2n}\}$ is the canonical basis in \mathbb{R}^{2n} and $\{\varepsilon^1, \dots, \varepsilon^{2n}\}$ is the corresponding dual basis, then the canonical symplectic 2-form is given by

$$\omega_0 := \varepsilon^1 \wedge \varepsilon^{n+1} + \dots + \varepsilon^n \wedge \varepsilon^{2n}.$$

Then the group $\text{Sp}_n(\mathbb{R})$ can be described as

$$\text{Sp}_n(\mathbb{R}) := \{A \in \text{GL}(2n, \mathbb{R}) \mid {}^t A J A = J\}$$

where $J := \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$.

Remark 2.6. The symplectic group $\mathrm{Sp}_n(\mathbb{R})$ is itself a Lie group, but this Lie group is not the definition of “symplectic Lie group”.

We wish to study the set $\Omega(\mathfrak{g})$ and use Proposition 2.4 to search for 2-forms that are closed, but this set can be rather big so we introduce the concept of the *moduli space* in Section 3.

2.2. Subalgebras of symplectic Lie algebras. In this subsection, we recall the definition of some particular subalgebras of symplectic Lie algebras and make a brief comment about their importance. For details, we refer to [1] and references within.

In the context of symplectic Lie algebras, isotropic and Lagrangian subalgebras are of particular importance (at the Lie group level these correspond to isotropic and Lagrangian subgroups, respectively). Given a subspace W of a symplectic vector space (V, ω) , recall that the orthogonal space W^\perp with respect to ω is defined by

$$W^\perp := \{v \in V \mid \omega(v, w) = 0 \text{ for all } w \in W\}.$$

Definition 2.7. Let (\mathfrak{g}, ω) be a symplectic Lie algebra. A subalgebra $\mathfrak{l} \subset \mathfrak{g}$ is said to be *isotropic* (resp. *Lagrangian*) if it satisfies $\mathfrak{l} \subset \mathfrak{l}^\perp$ (resp. $\mathfrak{l} = \mathfrak{l}^\perp$).

For example, the theory of *symplectic reduction* with respect to isotropic ideals has been used to study the structure of symplectic Lie algebras. Lie algebras that do not have a symplectic reduction (called *irreducible*) or those that can be reduced to a trivial Lie algebra (called *completely reducible*) are of particular interest. In particular, a classification exists for irreducible symplectic Lie algebras. Lagrangian ideals also appear in connection with flat Lie algebras, as every symplectic Lie algebra with a Lagrangian ideal can be constructed as Lagrangian extension of a flat Lie algebra.

In low dimensions several results with respect to the existence of isotropic and Lagrangian subalgebras are known, but in general the question is still not satisfactorily answered.

3. THE MODULI SPACES

In this section, we define the moduli space of left-invariant nondegenerate 2-forms on a Lie group, and formulate the procedure to obtain a kind of generalization of the Milnor frames.

3.1. The definition. Consider the automorphism group of \mathfrak{g} defined by

$$\mathrm{Aut}(\mathfrak{g}) := \{\phi \in \mathrm{GL}(2n, \mathbb{R}) \mid \phi[\cdot, \cdot] = [\phi(\cdot), \phi(\cdot)]\}.$$

Also define $\mathbb{R}^\times := \mathbb{R} \setminus 0$. Then we can consider the set

$$\mathbb{R}^\times \mathrm{Aut}(\mathfrak{g}) := \{\phi \in \mathrm{GL}(2n, \mathbb{R}) \mid \phi \in \mathrm{Aut}(\mathfrak{g}), c \in \mathbb{R}^\times\},$$

which is a subgroup of $\mathrm{GL}(2n, \mathbb{R})$. Hence it naturally acts on $\Omega(\mathfrak{g})$. We can then consider the orbit space of this action.

Definition 3.1. The orbit space of the action of $\mathbb{R}^\times \mathrm{Aut}(\mathfrak{g})$ on $\Omega(\mathfrak{g})$ will be called the *moduli space of left-invariant nondegenerate 2-forms* and will be denoted by

$$\mathfrak{P}\Omega(\mathfrak{g}) := \mathbb{R}^\times \mathrm{Aut}(\mathfrak{g}) \backslash \Omega(\mathfrak{g}) := \{\mathbb{R}^\times \mathrm{Aut}(\mathfrak{g}) \cdot \omega \mid \omega \in \Omega(\mathfrak{g})\}.$$

One can easily see that, if $\omega_1, \omega_2 \in \Omega(\mathfrak{g})$ are in the same $\mathbb{R}^\times \mathrm{Aut}(\mathfrak{g})$ -orbit, then they are equivalent up to automorphism and scale. Therefore there is a surjection from the moduli space $\mathfrak{P}\Omega(\mathfrak{g})$ onto the quotient space

$$\Omega(\mathfrak{g}) / \text{“up to automorphism and scale”}.$$

This would be not bijective, since ω and $-\omega$ are possibly not in the same $\mathbb{R}^\times \text{Aut}(\mathfrak{g})$ -orbit. In other words, the moduli space $\mathfrak{P}\Omega(\mathfrak{g})$ gives a finer partition than the latter quotient space. Note that the equivalence up to automorphism and scale preserves the closedness of 2-forms. Therefore, in order to search closed 2-forms, we have only to consider this quotient space, and also the moduli space.

In the latter sections, instead of studying $\Omega(\mathfrak{g})$ directly we will focus on studying the moduli space: we want to find orbits that correspond to closed 2-forms. As mentioned in the introduction, inspired in [7] and [10] we introduce in the next subsection a strategy to study the moduli space and to find the orbits of closed 2-forms.

3.2. Milnor frames procedure. Remember that given a symplectic vector space (V, ω) with $\dim V = 2n$, we can always choose a basis $\{x_1, \dots, x_{2n}\}$ of V such that for $i < j$

$$\omega(x_i, x_j) = \begin{cases} 1 & (\text{if } j = i + n), \\ 0 & (\text{all other cases}). \end{cases}$$

This basis is called a symplectic basis.

Remember that we denote by ω_0 the canonical symplectic form on a Lie algebra $\mathfrak{g} \cong \mathbb{R}^{2n}$. Let $\{e_1, \dots, e_{2n}\}$ denote the corresponding symplectic basis. To simplify the notation let us denote the orbit of $\mathbb{R}^\times \text{Aut}(\mathfrak{g})$ through $\omega \in \Omega(\mathfrak{g})$ by

$$[\omega] := (\mathbb{R}^\times \text{Aut}(\mathfrak{g})) \cdot \omega := \{\phi \cdot \omega \mid \phi \in \mathbb{R}^\times \text{Aut}(\mathfrak{g})\}.$$

Definition 3.2. A subset $U \subset \text{GL}(2n, \mathbb{R})$ is called a *set of representatives* of $\mathfrak{P}\Omega(\mathfrak{g})$ if it satisfies

$$\mathfrak{P}\Omega(\mathfrak{g}) = \{[h \cdot \omega_0] \mid h \in U\}.$$

Remark 3.3. Of course the set of representatives is not unique. In practice we want the set of representatives to be as small as possible.

Let $[[g]]$ denote the double coset of $g \in \text{GL}(2n, \mathbb{R})$ defined by

$$[[g]] := \mathbb{R}^\times \text{Aut}(\mathfrak{g}) g \text{Sp}(2n, \mathbb{R}) := \{\phi g s \mid \phi \in \mathbb{R}^\times \text{Aut}(\mathfrak{g}), s \in \text{Sp}_n(\mathbb{R})\}.$$

By standard theory of double coset spaces, we have a criterion for a set U to be a set of representatives (we refer to [7]).

Lemma 3.4. *Let $U \subset \text{GL}(2n, \mathbb{R})$, and assume that for every $g \in \text{GL}(2n, \mathbb{R})$ there exists $h \in U$ such that $h \in [[g]]$. Then U is a set of representatives of $\mathfrak{P}\Omega(\mathfrak{g})$.*

Now we state a theorem for obtaining Milnor type frames in the symplectic case.

Theorem 3.5. *Let U be a set of representatives of $\mathfrak{P}\Omega(\mathfrak{g})$. Then for every $\omega \in \Omega(\mathfrak{g})$ there exist $k > 0$, $\phi \in \text{Aut}(\mathfrak{g})$ and $h \in U$ such that $\{\phi g e_1, \dots, \phi g e_{2n}\}$ is a symplectic basis with respect to $k\omega$.*

Proof. The proof is similar to [7], Theorem 2.4. Take $\omega \in \Omega(\mathfrak{g})$. Let ω_0 be the canonical symplectic form on \mathfrak{g} . From the definition of a set of representatives there exists $h \in U$ such that $[\omega] = [h \cdot \omega_0]$. Hence, there exist $c \in \mathbb{R}^\times$ and $\phi \in \text{Aut}(\mathfrak{g})$ such that $\omega = (c\phi h) \cdot \omega_0$. Then we can easily check that $\{\phi h e_1, \dots, \phi h e_{2n}\}$ is a symplectic basis with respect to $c^2\omega$. Note that $k := c^2 > 0$, which completes the proof. \square

The basis obtained in this theorem will be called *Milnor frames*. Notice that if U has a small number of parameters, the bracket relations of the Milnor frames will also be given in terms of a small set of parameters. In such cases, it becomes much easier to search for closed 2-forms inside of $\Omega(\mathfrak{g})$. In the same way Milnor frames can also be useful to search for isotropic or Lagrangian subalgebras. We will show the previous procedure in two concrete examples in Section 5. Before working some concrete examples we introduce a tool that will be useful to calculate a smaller set of representatives and to obtain nice Milnor frames.

4. QR SYMPLECTIC DECOMPOSITION

In order to obtain a nice set of representatives, it is useful to have general results for decomposing matrices using symplectic matrices. Some of the known results can be seen in [2] and [3]. In this section we obtain a slight modification.

First of all we set up some notations. We denote by $M_{a \times b}(\mathbb{R})$ the set of all $a \times b$ real matrices, by I_k the $k \times k$ identity matrix, and by tM the transpose of a matrix M .

For $M \in M_{2n \times 2n}(\mathbb{R})$ we frequently use block decompositions. In most of the matrices we use in this paper the size of each block can be understood from the shape of the matrix, if confusion is possible we will describe explicitly the size of the blocks. Also recall that $P \in \text{GL}(n, \mathbb{R})$ is called a permutation matrix if it induces a permutation among the elements in the standard basis $\{e_1, \dots, e_n\}$. For each permutation $\sigma \in S_n$ of degree n , we denote the corresponding permutation matrix by $P_\sigma^{(n)}$, if there is no possible confusion we will simply denote it by P_σ or P . For example, the permutation matrix corresponding to the cyclic permutation $(n, n-1, \dots, 2, 1)$ is given by

$$P_{(n, n-1, \dots, 2, 1)}^{(n)} = \left(\begin{array}{c|c} & I_{n-1} \\ \hline 1 & \end{array} \right).$$

We here make a list of well-known symplectic matrices in terms of the block decompositions.

Proposition 4.1. *The next $2n \times 2n$ matrices are symplectic:*

(1) Type 1

$$\left(\begin{array}{c|c} I_n & C \\ \hline 0 & I_n \end{array} \right) \text{ or } \left(\begin{array}{c|c} I_n & 0 \\ \hline C & I_n \end{array} \right) \quad \text{with } C = {}^tC.$$

(2) Type 2

$$\left(\begin{array}{c|c} A & 0 \\ \hline 0 & {}^tA^{-1} \end{array} \right) \quad \text{for any } A \in \text{GL}(n, \mathbb{R}).$$

(3) Type 3

$$\left(\begin{array}{c|c} P & 0 \\ \hline 0 & P \end{array} \right) \quad \text{for any permutation matrix } P \in \text{GL}(n, \mathbb{R}).$$

Note that Type 3 is just a special case of Type 2 because ${}^tP_\sigma^{-1} = P_\sigma$, but we write it separately because we use it frequently. The next lemma gives a key step to obtain our decomposition theorem.

Lemma 4.2. *For all $M \in \text{GL}(2n, \mathbb{R})$, there exists a symplectic matrix $S \in \text{Sp}_n(\mathbb{R})$ such that the left-upper $n \times n$ block of MS is nonsingular.*

Proof. Take any $M \in \text{GL}(2n, \mathbb{R})$, and denote its block decomposition by

$$(4.1) \quad M = \left(\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right),$$

where A, B, C, D are $n \times n$ matrices. Let $r := \text{rank}(A)$. If $r = n$, then it is enough to put $S := I_{2n}$. Assume that $r < n$. Then we have only to show that there exists $S \in \text{Sp}_n(\mathbb{R})$ such that the left-upper $n \times n$ block of MS has rank bigger than r .

First of all, since $r = \text{rank}(A) < n$, it follows from linear algebra that there exist $g_1, g_2 \in \text{GL}(n, \mathbb{R})$ such that

$$(4.2) \quad g_1 A g_2 = \left(\begin{array}{c|c} I_r & 0 \\ \hline 0 & 0 \end{array} \right).$$

We define the following matrices:

$$(4.3) \quad K := \left(\begin{array}{c|c} g_1 & 0 \\ \hline 0 & I_n \end{array} \right) \in \mathrm{GL}(2n, \mathbb{R}), \quad S_1 := \left(\begin{array}{c|c} g_2 & 0 \\ \hline 0 & {}^t g_2^{-1} \end{array} \right) \in \mathrm{Sp}_n(\mathbb{R}).$$

Then one can directly see that

$$(4.4) \quad M_1 := KMS_1 = \left(\begin{array}{cc|cc} I_r & 0 & * & * \\ 0 & 0 & \alpha & \beta \\ \hline * & * & * & * \\ * & * & * & * \end{array} \right),$$

where $\alpha \in \mathrm{M}_{n-r \times r}(\mathbb{R})$ and $\beta \in \mathrm{M}_{n-r \times n-r}(\mathbb{R})$.

We here claim that we can assume $\beta \neq 0$ without loss of generality. In order to prove this, assume $\beta = 0$. Then one has $\alpha \neq 0$, since M_1 is nonsingular. Therefore there exists $\gamma \in \mathrm{M}_{r \times n-r}(\mathbb{R})$ such that $\alpha\gamma \neq 0$. We define a symplectic matrix of Type 2 by

$$(4.5) \quad S_2 := \left(\begin{array}{cc|cc} I_r & 0 & & \\ -{}^t\gamma & I_{n-r} & & \\ \hline & & I_r & \gamma \\ & & 0 & I_{n-r} \end{array} \right) \in \mathrm{Sp}_n(\mathbb{R}).$$

A direct calculation yields that

$$(4.6) \quad M_1 S_2 = \left(\begin{array}{cc|cc} I_r & 0 & * & * \\ 0 & 0 & \alpha & \alpha\gamma \\ \hline * & * & * & * \\ * & * & * & * \end{array} \right).$$

Recall that $\alpha\gamma \neq 0$, which completes the proof of the claim.

We assume $\beta \neq 0$ from now on. Let us consider a symplectic matrix of Type 1 defined by

$$(4.7) \quad S_3 := \left(\begin{array}{c|c} I_r & \\ \hline I_{n-r} & \\ \hline & I_r \\ & I_{n-r} \end{array} \right) \in \mathrm{Sp}_n(\mathbb{R}).$$

Then one can directly see that

$$(4.8) \quad M_2 := M_1 S_3 = \left(\begin{array}{cc|cc} I_r & 0 & * & * \\ 0 & \beta & \alpha & \beta \\ \hline * & * & * & * \\ * & * & * & * \end{array} \right).$$

Since $\beta \neq 0$, the rank of the left-upper block of M_2 is bigger than r . This is also true for

$$(4.9) \quad K^{-1}M_2 = K^{-1}M_1S_3 = K^{-1}(KMS_1)S_3 = M(S_1S_3),$$

since the multiplication by K^{-1} does not change the rank of the left-upper block. This completes the proof. \square

We are now in the position to give our decomposition theorem. Recall that a matrix $T = (t_{ij}) \in \mathrm{M}_{n \times n}(\mathbb{R})$ is said to be strictly lower triangular (resp. lower triangular) if $t_{ij} = 0$ holds for all $i \leq j$ (resp. $i < j$).

Theorem 4.3. *For all $M \in \mathrm{GL}(2n, \mathbb{R})$, there exist a symplectic matrix $S \in \mathrm{Sp}_n(\mathbb{R})$ and a strictly lower triangular matrix $T \in \mathrm{M}_{n \times n}(\mathbb{R})$ such that*

$$MS = \left(\begin{array}{c|c} I_n & T \\ \hline * & * \end{array} \right).$$

Proof. Take any $M \in \mathrm{GL}(2n, \mathbb{R})$. One then knows by Lemma 4.2 that there exists $S_1 \in \mathrm{Sp}_n(\mathbb{R})$ such that the left-upper $n \times n$ block of MS_1 is nonsingular. Denote by $A \in \mathrm{GL}(n, \mathbb{R})$ the left-upper block of MS_1 , that is,

$$(4.10) \quad M_1 := MS_1 = \left(\begin{array}{c|c} A & * \\ \hline * & * \end{array} \right).$$

Then, by using a symplectic matrix of Type 2, we can transform M_1 into

$$(4.11) \quad M_2 := M_1 \left(\begin{array}{c|c} A^{-1} & 0 \\ \hline 0 & {}^t A \end{array} \right) = \left(\begin{array}{c|c} I_n & B \\ \hline * & * \end{array} \right).$$

For the block B , there exists a symmetric matrix $C \in \mathrm{M}_{n \times n}(\mathbb{R})$ such that $C + B$ is strictly lower triangular. Hence, by using a symplectic matrix of Type 1, we can transform M_2 into

$$(4.12) \quad M_3 := M_2 \left(\begin{array}{c|c} I_n & C \\ \hline 0 & I_n \end{array} \right) = \left(\begin{array}{c|c} I_n & C + B \\ \hline * & * \end{array} \right),$$

which completes the proof. \square

Our result is a modification of the so called QR unitary decomposition (see for example [2] Theorem 2) as follows.

Remark 4.4 (Unitary QR decomposition). For any $A \in \mathrm{M}_{2n \times 2n}(\mathbb{R})$, there always exists a decomposition of the form $A = RQ$, where

$$Q \in \mathrm{Sp}_n(\mathbb{R}) \cap \mathrm{O}(2n, \mathbb{R}), \quad R = \left(\begin{array}{c|c} R_{11} & R_{12} \\ \hline * & * \end{array} \right),$$

with R_{11} being lower triangular and R_{12} strictly lower triangular.

Notice that in our case we are not restricted to unitary matrices and that the matrix we want to decompose is always nonsingular.

5. EXAMPLES

5.1. The Lie group $G_{\mathbb{R}\mathrm{H}^{2n}}$. This group is the semidirect product of the abelian group \mathbb{R} and \mathbb{R}^{2n-1} , where \mathbb{R} acts on \mathbb{R}^{2n-1} by $t.x := e^t x$ ($t \in \mathbb{R}, x \in \mathbb{R}^{2n-1}$). The corresponding Lie algebra of this group is $\mathfrak{g}_{\mathbb{R}\mathrm{H}^{2n}} \cong \mathbb{R}^{2n} = \mathrm{Span}\{e_1, \dots, e_{2n}\}$, with bracket relations given by $[e_1, e_k] = e_k$ where $k = 2, \dots, 2n$. For this Lie algebra it is known that (see [9])

$$(5.1) \quad \mathbb{R}^\times \mathrm{Aut}(\mathfrak{g}_{\mathbb{R}\mathrm{H}^{2n}}) = \left\{ \left(\begin{array}{c|ccc} * & 0 & \cdots & 0 \\ \hline * & & & \\ \vdots & & * & \\ * & & & \end{array} \right) \in \mathrm{GL}(2n, \mathbb{R}) \right\},$$

where the size of this block decomposition is $(1, 2n - 1)$.

Proposition 5.1. *The action of $\mathbb{R}^\times \mathrm{Aut}(\mathfrak{g}_{\mathbb{R}\mathrm{H}^{2n}})$ on $\Omega(\mathfrak{g}_{\mathbb{R}\mathrm{H}^{2n}})$ is transitive. A set of representatives U for this action is given by*

$$U = \{I_{2n}\}.$$

Proof. Take any $g \in \mathrm{GL}(2n, \mathbb{R})$. By Theorem 4.3 there exists a matrix $S \in \mathrm{Sp}_n(\mathbb{R})$ such that

$$(5.2) \quad [[g]] \ni gS = \left(\begin{array}{c|c} I_n & T \\ \hline * & * \end{array} \right) =: g_1,$$

where T is strictly lower triangular. Hence one has $g_1 \in \mathbb{R}^\times \text{Aut}(\mathfrak{g}_{\mathbb{R}H^{2n}})$, and also $g_1^{-1} \in \mathbb{R}^\times \text{Aut}(\mathfrak{g}_{\mathbb{R}H^{2n}})$. This shows that

$$(5.3) \quad [[g]] \ni g_1^{-1} g_1 = I_{2n},$$

which completes the proof. \square

Theorem 5.2 (Milnor-type). *For all $\omega \in \Omega(\mathfrak{g}_{\mathbb{R}H^{2n}})$, there exist $t > 0$ and a symplectic basis $\{x_1, \dots, x_{2n}\} \subset \mathfrak{g}_{\mathbb{R}H^{2n}}$ with respect to $t\omega$ such that*

$$[x_1, x_k] = x_k \quad \text{for } k = 2, \dots, 2n.$$

Proof. By Proposition 5.1, a set of representatives U for the action of $\mathbb{R}^\times \text{Aut}(\mathfrak{g}_{\mathbb{R}H^{2n}})$ on $\Omega(\mathfrak{g}_{\mathbb{R}H^{2n}})$ is given just by the identity I_{2n} . Take any $\omega \in \Omega(\mathfrak{g}_{\mathbb{R}H^{2n}})$. Then, by Theorem 3.5 there exists $\phi \in \text{Aut}(\mathfrak{g}_{\mathbb{R}H^{2n}})$ such that $\{x_1 := \phi e_1, \dots, x_n := \phi e_n\}$ is symplectic with respect to $t\omega$. Hence, we only have to check the bracket relations among them:

$$[x_1, x_k] = [\phi e_1, \phi e_k] = \phi[e_1, e_k] = \phi e_k = x_k \quad \text{for } k = 2, \dots, 2n.$$

This completes the proof. \square

Now we can use Proposition 2.4 to search for closed 2-forms.

Corollary 5.3. *Let $\omega \in \Omega(\mathfrak{g}_{\mathbb{R}H^{2n}})$. Then $d\omega = 0$ if and only if $n = 1$.*

Proof. For any $\omega \in \Omega(\mathfrak{g}_{\mathbb{R}H^{2n}})$, we can find a symplectic basis $\{x_1, \dots, x_{2n}\}$ with respect to $t\omega$ such that the bracket relation are given by Theorem 5.2. Notice that when using the Milnor frames obtained in Theorem 5.2 to search for symplectic forms, the parameter t has no effect on the condition of Proposition 2.4. Therefore, we can take $t = 1$ without loss of generality.

If $n = 1$ we automatically have $d\omega = 0$. If $n > 1$ we have $n + 2 \leq 2n$ and $x_{n+2} \in \mathfrak{g}_{\mathbb{R}H^{2n}}$. Hence

$$\begin{aligned} d\omega(x_1, x_2, x_{n+2}) &= \omega(x_2, [x_{n+2}, x_1]) + \omega(x_{n+2}, [x_1, x_2]) \\ &= -2\omega(x_2, x_{n+2}) = -2 \neq 0. \end{aligned}$$

Therefore, no closed 2-form corresponds to this case. This completes the proof. \square

In terms of the above symplectic basis, one can easily see that $\mathfrak{l} := \text{Span}\{x_2\}$ defines a Lagrangian ideal of $\mathfrak{g}_{\mathbb{R}H^{2n}}$.

5.2. The Lie group $H^3 \times \mathbb{R}^{2n-3}$. This group is the direct product of the 3-dimensional Heisenberg Lie group H^3 and the abelian Lie group \mathbb{R}^{2n-3} . Denote by \mathfrak{h}^3 and \mathbb{R}^{2n-3} the corresponding Lie algebras of H^3 and \mathbb{R}^{2n-3} , respectively. Then the corresponding Lie algebra of $H^3 \times \mathbb{R}^{2n-3}$ is

$$\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3} \cong \mathbb{R}^{2n} = \text{Span}\{e_1, \dots, e_{2n}\}$$

with bracket relations given by $[e_1, e_2] = e_{2n}$. For this Lie algebra we have that (see [9])

$$(5.4) \quad \mathbb{R}^\times \text{Aut}(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3}) = \left\{ \left(\begin{array}{cc|ccc|c} * & * & 0 & \cdots & 0 & 0 \\ * & * & 0 & \cdots & 0 & 0 \\ \hline * & * & & & & 0 \\ \vdots & \vdots & & & & \vdots \\ * & * & & * & & 0 \\ \hline * & * & * & \cdots & * & * \end{array} \right) \in \text{GL}(2n, \mathbb{R}) \right\},$$

where the size of this block decomposition is $(2, 2n - 3, 1)$. In order to obtain a set of representatives, we start with a calculation related to permutation matrices.

Lemma 5.4. *For any matrix of permutation P_σ , we can select*

$$K_1 := \left(\begin{array}{c|c} P_{\sigma_1}^{(2)} & \\ \hline & P_{\sigma_2}^{(n-2)} \end{array} \right), \quad K_2 := \left(\begin{array}{c|c} P_{\sigma_3}^{(n-1)} & \\ \hline & 1 \end{array} \right)$$

such that

$$K_2 P_\sigma^t K_1 \in \{P_{(n,n-1,\dots,2,1)}, I_n\}.$$

Proof. Let P_σ be a permutation matrix. Recall that each column vector of P_σ coincides with some e_k . First of all we consider the case that the matrix P_σ is of the form

$$P_\sigma = \left(\begin{array}{c|ccc} & & & \\ \hline * & * & 0 & \dots & 0 \end{array} \right).$$

We can select a matrix K_1 of the above form such that

$$P'_\sigma = P_\sigma^t K_1 = \left(\begin{array}{c|ccc} & & & \\ \hline 1 & 0 & 0 & \dots & 0 \end{array} \right).$$

Then there exists a matrix K_2 such that $K_2 P'_\sigma = P_{(n,n-1,\dots,2,1)}$.

We next consider the remaining case, that is,

$$P_\sigma = \left(\begin{array}{c|ccc} & & & \\ \hline 0 & 0 & * & \dots & * \end{array} \right).$$

In a similar way as the first case, we select a matrix K_1 such that

$$P'_\sigma = P_\sigma^t K_1 = \left(\begin{array}{c|ccc} & & & \\ \hline 0 & \dots & 0 & 1 \end{array} \right).$$

Finally we select a matrix K_2 such that $K_2 P'_\sigma = I_n$, which completes the proof. \square

In terms of the above lemma, we study double cosets. The following lemma gives a key step to obtain a set of representatives.

Lemma 5.5. *For any $g \in \mathrm{GL}(2n, \mathbb{R})$, there exist $P \in \{I_n, P_{(n,n-1,\dots,2,1)}^{(n)}\}$ and a strictly lower triangular matrix T such that*

$$(5.5) \quad [[g]] \ni \left(\begin{array}{c|c} I_n & T \\ \hline 0 & P \end{array} \right).$$

Proof. Take $g \in \mathrm{GL}(2n, \mathbb{R})$. By Theorem 4.3 there exists a matrix $S \in \mathrm{Sp}_n(\mathbb{R})$ which changes the left-upper block to I_n , that is,

$$(5.6) \quad [[g]] \ni gS = \left(\begin{array}{c|c} I_n & * \\ \hline C & * \end{array} \right) =: g_1.$$

By using a certain matrix in $\mathbb{R}^\times \mathrm{Aut}(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3})$, we have

$$[[g]] \ni \left(\begin{array}{c|c} I_n & 0 \\ \hline -C & I_n \end{array} \right) \left(\begin{array}{c|c} I_n & * \\ \hline C & * \end{array} \right) = \left(\begin{array}{c|c} I_n & * \\ \hline 0 & D \end{array} \right) =: g_2.$$

Note that the matrix D is nonsingular. Then, by a result known as ‘‘LPU decomposition’’ ([8] Theorem 3.5.11), there exist a lower triangular matrix L and an

upper matrix triangular matrix U such that $P' := LDU$ is a permutation matrix. Therefore, we have

$$[[g]] \ni \left(\begin{array}{c|c} {}^tU & \\ \hline & L \end{array} \right) g_2 \left(\begin{array}{c|c} {}^tU^{-1} & \\ \hline & U \end{array} \right) = \left(\begin{array}{c|c} I_n & * \\ \hline 0 & P' \end{array} \right) =: g_3.$$

For the permutation matrix P' , we can choose K_1 and K_2 given in Lemma 5.4 such that $P := K_2 P' K_1 \in \{I_n, P_{(n,n-1,\dots,2,1)}^{(n)}\}$. Note that

$$\left(\begin{array}{c|c} K_1 & \\ \hline & K_2 \end{array} \right) \in \mathbb{R}^\times \text{Aut}(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3}),$$

because of the forms of K_1 and K_2 . Hence, we have

$$[[g]] \ni \left(\begin{array}{c|c} K_1 & \\ \hline & K_2 \end{array} \right) g_3 \left(\begin{array}{c|c} K_1^{-1} & \\ \hline & {}^tK_1 \end{array} \right) = \left(\begin{array}{c|c} I_n & * \\ \hline 0 & P \end{array} \right) =: g_4.$$

Finally, by the same way as in the proof of Theorem 4.3, we can use a symplectic matrix S' of Type 1 to transform g_4 into the desired form

$$[[g]] \ni g_4 S' = \left(\begin{array}{c|c} I_n & T \\ \hline 0 & P \end{array} \right),$$

where T is strictly lower triangular. This completes the proof. \square

We are now in the position to give a set of representatives for the action of $\mathbb{R}^\times \text{Aut}(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3})$. In fact, it consists of two or five points, depending on n .

Proposition 5.6. *A set of representatives U for the action of $\mathbb{R}^\times \text{Aut}(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3})$ on $\Omega(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3})$ is given by the matrices (2) if $n = 2$, and by the matrices (1)–(3) if $n > 2$:*

$$\begin{aligned} (1) & I_{2n} + kE_{2,n+1}, & (2) & \left(\begin{array}{c|c} I_n & 0 \\ \hline 0 & P_{(n,1,\dots,n-1)} \end{array} \right) + kE_{2,n+1} \\ (3) & \left(\begin{array}{c|c} I_n & 0 \\ \hline 0 & P_{(n,1,\dots,n-1)} \end{array} \right) + E_{3,n+1}, \end{aligned}$$

where $k \in \{0, 1\}$.

Proof. Take any $g \in \text{GL}(2n, \mathbb{R})$. It is enough to show that $[[g]]$ contains one of the matrices (1)–(3) if $n > 2$ and one of the matrices (2) if $n = 2$. By Lemma 5.5, there exist $P \in \{I_n, P_{(n,n-1,\dots,2,1)}^{(n)}\}$ and a strictly lower triangular matrix T such that

$$(5.7) \quad [[g]] \ni \left(\begin{array}{c|c} I_n & T \\ \hline 0 & P \end{array} \right) =: g_1.$$

We consider two cases according to the choice of P , and some subcases.

Case 1: $P = P_{(n,n-1,\dots,2,1)}^{(n)}$. We write the matrix $T \in M_{n \times n}(\mathbb{R})$ as

$$T = \left(\begin{array}{c|c} 0 & 0 \\ \hline * & T' \end{array} \right),$$

where $T' \in M_{(n-1) \times (n-1)}(\mathbb{R})$ is strictly lower triangular. We define some matrices:

$$A := \left(\begin{array}{c|c} 0 & 0 \\ \hline -T' & 0 \end{array} \right) \in M_{n \times n}(\mathbb{R}), \quad D := \left(\begin{array}{c|c} I_n & A \\ \hline 0 & I_n \end{array} \right) \in \mathbb{R}^\times \text{Aut}(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3}).$$

We use the matrix D to transform the matrix g_1 into

$$[[g]] \ni Dg_1 = \left(\begin{array}{c|c} I_n & T + AP \\ \hline 0 & P \end{array} \right) =: g_2.$$

By a direct calculation of $T + AP$, one can express g_2 as

$$(5.8) \quad g_2 = \left(\begin{array}{c|c} I_n & 0 \\ \hline 0 & P \end{array} \right) + x_2 E_{2,n+1} + \cdots + x_n E_{n,n+1}.$$

We here consider the following three subcases.

Subcase 1-(i): $x_2 \neq 0$. In this case, we can transform x_2 into 1 as follows. Define the following two diagonal matrices:

$$V_1 := E_{1,1} + (1/x_2)E_{2,2} + E_{3,3} + \cdots + E_{n,n}, \quad V_2 = P({}^tV_1^{-1}){}^tP.$$

Then we can take matrices which makes x_2 into 1, that is,

$$\begin{aligned} [[g]] &\ni \left(\begin{array}{c|c} V_1 & \\ \hline & V_2 \end{array} \right) g_2 \left(\begin{array}{c|c} V_1^{-1} & \\ \hline & {}^tV_1 \end{array} \right) \\ &= \left(\begin{array}{c|c} I_n & 0 \\ \hline 0 & P \end{array} \right) + E_{2,n+1} + x_3E_{3,n+1} + \cdots + x_nE_{n,n+1} =: g_3. \end{aligned}$$

Now define the matrices:

$$T_1 := I_n - x_3E_{3,2} - \cdots - x_nE_{n,2} \in M_{n \times n}(\mathbb{R}), \quad T_2 := P({}^tT_1^{-1}){}^tP.$$

We can take matrices that eliminate the rest of parameters in g_3 , that is,

$$(5.9) \quad [[g]] \ni \left(\begin{array}{c|c} T_1 & \\ \hline & T_2 \end{array} \right) g_3 \left(\begin{array}{c|c} T_1^{-1} & \\ \hline & {}^tT_1 \end{array} \right) = \left(\begin{array}{c|c} I_n & 0 \\ \hline 0 & P \end{array} \right) + E_{2,n+1}.$$

This coincides with the matrix (2) of $k = 1$ in the statement of this proposition.

Subcase 1-(ii): $x_2 = 0$ and $n = 2$. In this case we immediately obtain

$$(5.10) \quad g_2 = \left(\begin{array}{c|c} I_2 & 0 \\ \hline 0 & P_{(2,1)} \end{array} \right),$$

which coincides with the matrix (2) of $k = 0$ in the statement of this proposition.

Subcase 1-(iii): $x = 0$ and $n > 2$. Given a vector ${}^t(x_3, \dots, x_n)$, it is well known that there exists a matrix $U \in \text{GL}(n-2, \mathbb{R})$ such that

$$U \cdot {}^t(x_3, \dots, x_n) = {}^t(k, 0, \dots, 0),$$

where $k \in \{0, 1\}$. Define the matrices:

$$H_1 := \left(\begin{array}{c|c} I_2 & \\ \hline & U \end{array} \right) \in \text{GL}(n, \mathbb{R}), \quad H_2 := P({}^tH_1^{-1}){}^tP.$$

Now we can consider the transformation

$$[[g]] \ni \left(\begin{array}{c|c} H_1 & \\ \hline & H_2 \end{array} \right) g_2 \left(\begin{array}{c|c} H_1^{-1} & \\ \hline & {}^tH_1 \end{array} \right) =: g_3.$$

The effect of the previous transformation is just the action of U on the vector ${}^t(x_3, \dots, x_n)$. Therefore one has

$$(5.11) \quad g_3 = \left(\begin{array}{c|c} I_n & 0 \\ \hline 0 & P \end{array} \right) + kE_{3,n+1},$$

where $k \in \{0, 1\}$. This coincides with the matrix (2) of $k = 0$, or with the matrix (3) in the statement of this proposition.

Case 2: $P = I_n$. We consider the following two subcases.

Subcase 2-(i): $n = 2$. We show that this case is equivalent to Case 1. We have

$$g_1 = \left(\begin{array}{c|c} I_2 & T \\ \hline 0 & I_2 \end{array} \right).$$

Define the matrices:

$$R = \left(\begin{array}{c|c} P_{(2,1)} & \\ \hline & I_2 \end{array} \right) \in \mathbb{R}^\times \text{Aut}(\mathfrak{h}^3 \oplus \mathbb{R}^1), \quad S = \left(\begin{array}{c|c} P_{(2,1)} & \\ \hline & P_{(2,1)} \end{array} \right) \in \text{Sp}_2(\mathbb{R}).$$

Now we can consider the transformation

$$[[g]] \ni Rg_1S = \left(\begin{array}{c|c} I_2 & * \\ \hline 0 & P_{(2,1)} \end{array} \right) =: g_2.$$

In a similar way to the final part of Lemma 5.5 we can use a symplectic matrix S' of Type 1 to transform g_2 into

$$[[g]] \ni g_2 S' = \left(\begin{array}{c|c} I_n & T \\ \hline 0 & P_{(2,1)} \end{array} \right),$$

where T is strictly lower triangular. This matrix corresponds to Case 1. Therefore, for the case of $n = 2$, we have obtained a set of representatives consisting of two points.

Subcase 2-(ii): $n > 2$. We start from the matrix g_1 , and consider the matrix $T = (t_{i,j}) \in M_{n \times n}(\mathbb{R})$. Let us put

$$T' := T - t_{2,1} E_{2,1}.$$

Since T is lower triangular, one has

$$D' := \left(\begin{array}{c|c} I_n & -T' \\ \hline 0 & I_n \end{array} \right) \in \mathbb{R}^\times \text{Aut}(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3}).$$

We use the matrix D' to transform the matrix g_2 into

$$[[g]] \ni D' g_1 = I_{2n} + t_{2,1} E_{2,n+1} =: g_2.$$

Finally, in a similar way to Case 1, if $t_{2,1} \neq 0$ we can transform it easily into 1. After this transformation we obtain

$$(5.12) \quad [[g]] \ni I_{2n} + k E_{2,n+1},$$

where $k \in \{0, 1\}$. This coincides with the matrix (1) in the statement of this proposition, which completes the proof. \square

Given this set of representatives we can give a Milnor-type theorem.

Theorem 5.7 (Milnor-type). *For all $\omega \in \Omega(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3})$, there exist $t > 0$, $k \in \{0, 1\}$ and a symplectic basis $\{x_1, \dots, x_{2n}\}$ of $\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3}$ with respect to $t\omega$ such that the bracket relations are given by one of the following:*

- (1) $[x_1, x_2] = x_{2n}$, $[x_1, x_{n+1}] = kx_{2n}$ (for $n > 2$),
- (2) $[x_1, x_2] = x_{n+1} - kx_2$, $[x_1, x_{n+1}] = kx_{n+1} - k^2 x_2$, or
- (3) $[x_1, x_2] = x_{n+1} - x_3$ (for $n > 2$).

Proof. Let $\{e_1, \dots, e_{2n}\}$ be the canonical basis of $\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3}$, whose bracket relation is given by $[e_1, e_2] = e_{2n}$. In Proposition 5.6 we obtained a set of representatives U for the action of $\mathbb{R}^\times \text{Aut}(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3})$ on $\Omega(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3})$. Take any $\omega \in \Omega(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3})$. Then it follows from Theorem 3.5 that there exist $u \in U$, $t > 0$ and $\phi \in \text{Aut}(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3})$ such that $\{x_1 := \phi u e_1, \dots, x_n := \phi u e_n\}$ is symplectic with respect to $t\omega$. Hence, we only have to check the bracket relations among them. We study each case corresponding to the five matrices in U case by case.

Case 1: we consider the case of $u = I_{2n} + k E_{2,n+1}$ with $k \in \{0, 1\}$ and $n > 2$, which corresponds to the matrices (1) in Proposition 5.6. In this case one has

$$ue_i = \begin{cases} ke_2 + e_{n+1} & (i = n+1), \\ e_i & (i \neq n+1). \end{cases}$$

Therefore, among $\{ue_1, \dots, ue_{2n}\}$, the only possible nonzero bracket relations are

$$\begin{aligned} [ue_1, ue_2] &= [e_1, e_2] = e_{2n} = ue_{2n}, \\ [ue_1, ue_{n+1}] &= [e_1, ke_2 + e_{n+1}] = kue_{2n}. \end{aligned}$$

By applying the automorphism ϕ to both sides, we obtain

$$\begin{aligned} [x_1, x_2] &= [\phi ue_1, \phi ue_2] = \phi[ue_1, ue_2] = \phi ue_{2n} = x_n, \\ [x_1, x_{n+1}] &= [\phi ue_1, \phi ue_{n+1}] = \phi[ue_1, ue_{n+1}] = \phi kue_{2n} = kx_{2n}. \end{aligned}$$

Therefore the bracket relations are given by (1) in the statement of this theorem.

Case 2: we consider the case of

$$\left(\begin{array}{c|c} I_n & 0 \\ \hline 0 & P_{(n,1,\dots,n-1)} \end{array} \right) + kE_{2,n+1} \quad (\text{with } k \in \{0, 1\}),$$

which corresponds to the matrices (2) in Proposition 5.6. In this case one has

$$ue_i = \begin{cases} e_i & (i \leq n), \\ ke_2 + e_{2n} & (i = n+1), \\ e_{i-1} & (i > n+1). \end{cases}$$

Therefore, among $\{ue_1, \dots, ue_{2n}\}$, the only possible nonzero bracket relations are

$$\begin{aligned} [ue_1, ue_2] &= [e_1, e_2] = e_{2n} = ue_{n+1} - kue_2, \\ [ue_1, ue_{n+1}] &= [e_1, ke_2 + e_{n+1}] = kue_{n+1} - k^2ue_2. \end{aligned}$$

By applying the automorphism ϕ to both sides, we obtain the bracket relations given by (2) in the statement of this theorem.

Case 3: we consider the case of $n > 2$ and

$$\left(\begin{array}{c|c} I_n & 0 \\ \hline 0 & P_{(n,1,\dots,n-1)} \end{array} \right) + E_{3,n+1},$$

which corresponds to the matrices (3) in Proposition 5.6. In this case one has

$$ue_i = \begin{cases} e_i & (i \leq n), \\ e_3 + e_{2n} & (i = n+1), \\ e_{i-1} & (i > n+1). \end{cases}$$

Therefore, among $\{ue_1, \dots, ue_{2n}\}$, the only possible nonzero bracket relations are

$$[ue_1, ue_2] = [e_1, e_2] = e_{2n} = ue_{n+1} - ue_3.$$

By applying the automorphism ϕ to both sides, we obtain the bracket relations given by (3). This completes the proof. \square

Now we can use Proposition 2.4 to search for closed 2-forms and, obtain the existence and uniqueness result as follows.

Corollary 5.8. *There exists a unique symplectic structure on $\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3}$ up to automorphism and scale. In fact, $\omega \in \Omega(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3})$ satisfies $d\omega = 0$ if and only if there exist $t > 0$ and a symplectic basis $\{x_1, \dots, x_{2n}\} \subset \mathfrak{h}^3 \oplus \mathbb{R}^{2n-3}$ with respect to $t\omega$ such that*

$$[x_1, x_2] = x_{n+1}.$$

Proof. For any $\omega \in \Omega(\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3})$ we can find a symplectic basis $\{x_1, \dots, x_{2n}\}$ with respect to $t\omega$ such that the bracket relations are given by (1)–(3) in Theorem 5.7. As we did in Proposition 5.3 we take $t = 1$, without loss of generality. We check the closed condition for each case.

Bracket relations (1). In this case we have $n > 2$, and

$$[x_1, x_2] = x_{2n}, \quad [x_1, x_{n+1}] = kx_{2n}.$$

Note that $[x_1, x_n] = 0$ since $n \neq 2$. Then one can see that

$$\begin{aligned} d\omega(x_1, x_2, x_n) &= \omega(x_1, [x_2, x_n]) + \omega(x_n, [x_1, x_2]) + \omega(x_2, [x_n, x_1]) \\ &= \omega(x_n, x_{2n}) = 1 \neq 0. \end{aligned}$$

Therefore, no closed 2-form corresponds to this case.

Bracket relations (2). Recall that it satisfies

$$[x_1, x_2] = x_{n+1} - kx_2, \quad [x_1, x_{n+1}] = kx_{n+1} - k^2x_2.$$

One can easily see that

$$d\omega(x_1, x_{n+1}, x_{n+2}) = \omega(x_{n+2}, [x_1, x_{n+1}]) = k^2.$$

Hence we have only to consider the case of $k = 0$. In this case, the bracket relations reduce to

$$[x_1, x_2] = x_{n+1}.$$

For this bracket relations it is easy to check that $d\omega = 0$. In particular, for each $j \in \{3, \dots, 2n\}$, we have

$$d\omega(x_1, x_2, x_j) = \omega(x_j, [x_1, x_2]) = \omega(x_j, x_{n+1}) = 0.$$

This bracket corresponds to a closed 2-form, as desired.

Bracket relations (3). In this case we have $n > 2$, and

$$[x_1, x_2] = x_{n+1} - x_3.$$

Then one can easily see that

$$d\omega(x_1, x_2, x_{n+3}) = \omega(x_{n+3}, [x_1, x_2]) = 1 \neq 0.$$

Therefore, no closed 2-form corresponds to this case. This completes the proof. \square

In terms of the above symplectic basis, one can easily see the existence of a Lagrangian normal subgroup in $H^3 \times \mathbb{R}^{2n-3}$, equipped with a left-invariant symplectic structure.

Corollary 5.9. *For every symplectic structure on $\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3}$ there exists a Lagrangian ideal.*

Proof. Let ω be a symplectic structure on $\mathfrak{h}^3 \oplus \mathbb{R}^{2n-3}$. Then by Corollary 5.8, there exist $t > 0$ and a symplectic basis $\{x_1, \dots, x_{2n}\}$ with respect to $t\omega$ such that

$$[x_1, x_2] = x_{n+1}.$$

Define $\mathfrak{l} = \{x_2, \dots, x_n, x_{n+1}\}$. Then one can easily check that \mathfrak{l} is a Lagrangian ideal. \square

REFERENCES

- [1] Baues, O., Cortés V.: *Symplectic Lie groups*. Astérisque **379** (2016).
- [2] Benner P., Faßbender H., Yang C.: *Some remarks on the complex J-symmetric eigenproblem*. Linear Algebra Appl. **544** (2018), 407–442.
- [3] Bunse-Gerstner, A.: *Matrix factorizations for symplectic QR-like methods*. Linear Algebra Appl. **83** (1986), 49–77.
- [4] Campoamor-Stursberg, R.: *Symplectic forms on six-dimensional real solvable Lie algebras*. I. Algebra Colloq. **16** (2009), 253–266.
- [5] Chu, B.Y.: *Symplectic homogeneous spaces*. Trans. Amer. Math. Soc. **197** (1974), 145–159.
- [6] Gómez, J. R., Jiménez-Merchén, A., Khakimdjjanov, Y.: *Symplectic structures on the filiform Lie algebras*. J. Pure Appl. Algebra **156** (2001), 15–31.
- [7] Hashinaga, T., Tamaru, H., Terada, K.: *Milnor-type theorems for left-invariant Riemannian metrics on Lie groups* J. Math. Soc. Japan **68** (2016), 669–684.
- [8] Horn, R.A., Horn, R.A., Johnson, C.R.: *Matrix Analysis*. Cambridge University Press (1990).
- [9] Kodama, H., Takahara, A., and Tamaru, H.: *The space of left-invariant metrics on a Lie group up to isometry and scaling*. Manuscripta Math. **135** (2011), 229–243.
- [10] Kubo, A., Onda, K., Taketomi, Y., Tamaru, H.: *On the moduli spaces of left-invariant pseudo-Riemannian metrics on Lie groups*. Hiroshima Math. J. **46** (2016), 357–374.
- [11] Lauret, J.: *Degenerations of Lie algebras and geometry of Lie groups*. Differential Geom. Appl. **18** (2003), 177–194.
- [12] Michel G., Abdelkader B.: *Sur les algèbres de Lie munies d'une forme symplectique*. Rend. Sem. Fac. Sci. Univ. Cagliari **57** (1987) 85–97.
- [13] Milnor, J.: *Curvatures of left invariant metrics on Lie groups*. Advances in Math. **21** (1976), 293–329.
- [14] Ovando, G.: *Four dimensional symplectic Lie algebras*. Beiträge Algebra Geom. **47** (2006), 419–434.
- [15] Wolf, J.A.: *Finiteness of orbit structure for real flag manifolds*. Geometriae Dedicata. **3** (1974), 377–384.

(L. Castellanos) DEPARTMENT OF MATHEMATICS, OSAKA CITY UNIVERSITY, SUGIMOTO, SUMIYOSHI-KU, OSAKA CITY, JAPAN 585-8585

Email address: `caste3.1416@gmail.com`

(H. Tamaru) DEPARTMENT OF MATHEMATICS, OSAKA CITY UNIVERSITY, SUGIMOTO, SUMIYOSHI-KU, OSAKA CITY, JAPAN 585-8585

Email address: `tamaru@sci.osaka-cu.ac.jp`