

Deletion of degQ gene enhances outer membrane vesicle production of *Shewanella oneidensis* cells

| | |
|-------|--|
| メタデータ | 言語: English 出版者: Springer 公開日: 2018-06-11 キーワード: 外膜小胞 作成者: 尾島, 由紘, Mohanadas, Thivagaran, Kitamura, Kosei, 布上, 翔太, 矢嶋, 黎輝, 田谷, 正仁 メールアドレス: 所属: Osaka City University, Osaka University, Osaka University, Osaka University, Osaka University, Osaka University, Osaka University, Osaka University |
| URL | https://ocu-omu.repo.nii.ac.jp/records/2019641 |

Deletion of *degQ* gene enhances outer membrane vesicle production of *Shewanella oneidensis* cells

Ojima Yoshihiro, Mohanadas Thivagaran, Kitamura Kosei,
Nunogami Shota, Yajima Reiki, Taya Masahito

| | |
|--------------------|--|
| Citation | Archives of Microbiology. 199(3); 415-423 |
| Issue Date | 2017-04 |
| Type | Journal Article |
| Textversion | author |
| Rights | This is a post-peer-review, pre-copyedit version of an article published in Archives of Microbiology. The final authenticated version is available online at: https://doi.org/10.1007/s00203-016-1315-4 |
| DOI | 10.1007/s00203-016-1315-4 |

Self-Archiving by Author(s)
Placed on: Osaka City University

Deletion of *degQ* gene enhances outer membrane vesicle
production of *Shewanella oneidensis* cells

Yoshihiro Ojima*[†], Thivagaran Mohanadas, Kosei Kitamura, Shota Nunogami, Reiki

Yajima and Masahito Taya

*Division of Chemical Engineering, Graduate School of Engineering Science, Osaka
University, 1-3 Machikaneyama-cho, Toyonaka, Osaka 560-8531, Japan*

*[†]Present address: Osaka City University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka
558-8585, Japan)*

* Corresponding author: Yoshihiro Ojima

Tel: +81-6-6605-2163; fax: +81-6-6605-3092

e-mail: ojima@bioa.eng.osaka-cu.ac.jp

Keywords: *Shewanella oneidensis*, outer membrane vesicle, *degQ*, periplasmic serine
protease

1 **Abstract**

2 *Shewanella oneidensis* is a Gram-negative facultative anaerobe that can use a wide
3 variety of terminal electron acceptors for anaerobic respiration. In this study, *S.*
4 *oneidensis degQ* gene, encoding a putative periplasmic serine protease was cloned and
5 expressed. The activity of purified DegQ was inhibited by diisopropyl fluorophosphate,
6 a typical serine protease-specific inhibitor, indicating that DegQ is a serine protease.
7 In-frame deletion and subsequent complementation of the *degQ* was carried out to
8 examine the effect of envelope stress on the production of outer membrane vesicles
9 (OMVs). Analysis of periplasmic proteins from the resulting *S. oneidensis* strain
10 showed that deletion of *degQ* induced protein accumulation, and resulted in a
11 significant decrease in protease activity within the periplasmic space. OMVs from the
12 wild-type and mutant strains were purified and observed by transmission electron
13 microscopy. Sodium dodecyl sulfate-polyacrylamide gel electrophoresis analysis of the
14 OMVs showed a prominent band at ~37 kDa. Nano liquid chromatography-tandem
15 mass spectrometry analysis identified three outer membrane porins (SO3896, SO1821
16 and SO3545) as dominant components of the band, suggesting that these proteins could
17 be used as indices for comparing OMV production by *S. oneidensis* strains. Quantitative
18 evaluation showed that *degQ*-deficient cells had a five-fold increase in OMV production
19 compared with wild-type cells. Thus, the increased OMV production following the
20 deletion of DegQ in *S. oneidensis* may be responsible for the increase in envelope stress.

1 **Introduction**

2 Many Gram-negative bacteria form outer membrane vesicles (OMVs) as part of
3 their natural growth cycle (Rothfield and Pearlman-Kothencz 1969; Wensink and
4 Witholt 1981; Nowotny et al. 1982; Beveridge 1999). OMVs are spherical bilayered
5 proteolipids with a diameter of 20–250 nm, and are compositionally similar to the outer
6 membranes of bacteria, which contain outer membrane proteins, lipids, periplasmic
7 proteins, lipopolysaccharides, RNA, and DNA (Knox et al. 1966; Kadurugamuwa and
8 Beveridge 1996). Gram-negative bacteria living in fresh water environments and as
9 pathogens in animal hosts produce OMVs (Schwechheimer et al. 2013). Although OMV
10 production appears to be a ubiquitous physiological process, the factors triggering OMV
11 formation have not been conclusively identified. Previous research suggested that
12 disturbances in growth, exposure to antibiotics, or simply turnover in cell wall
13 components initiates vesicle formation (Knox et al. 1966; Kadurugamuwa and
14 Beveridge 1995; Zhou et al. 1998).

15 Recent work has also suggested that OMV production is closely related to envelope
16 stress (McBroom and Kuehn 2007; Schwechheimer and Kuehn 2015). The hypothesis
17 that vesiculation plays an important role in disposing envelope “garbage” was first
18 proposed with respect to proteinaceous waste accumulation in an *Escherichia coli*
19 *degP*-deficient strain. DegP is a periplasmic chaperone and protease in *E. coli* that
20 manages envelope stress caused by unfolded and misfolded periplasmic proteins
21 (Lipinska et al. 1990; Schwechheimer and Kuehn 2013). Random mutagenesis of *E. coli*
22 determined that a transposon insertion immediately following the start codon of *degP*
23 resulted in hypervesiculation (McBroom et al. 2006). Misfolded proteins were not
24 digested in the *degP*-deficient strain, and such undigested substrates can cause lethality

1 under conditions of relatively high temperature, at which protein misfolding is more
2 likely to occur (Strauch et al. 1989). Notably, the lumen of OMVs produced by the
3 *degP*-deficient strain contained misfolded outer-membrane proteins, which can be
4 substrates for DegP (Schwechheimer and Kuehn 2013). Thus, *E. coli* strains lacking
5 DegP likely increase OMV production as a survival strategy to eliminate these
6 undesired proteins from the cell envelope.

7 *Shewanella oneidensis* is a Gram-negative facultative anaerobe that can use a wide
8 variety of terminal electron acceptors for anaerobic respiration (Myers and Nealson
9 1988; Myers and Nealson 1990; Myers and Myers 1994; Myers and Myers 2000). It
10 plays an important role in many environmental and biotechnological processes,
11 including removing toxic metal contaminants, such as uranium and chromium, by
12 reductive reactions (Fredrickson et al. 2008; Belchik et al. 2011). The genome sequence
13 of *S. oneidensis* suggested the presence of a putative periplasmic serine protease, DegQ
14 (Dai et al. 2015). Furthermore, deletion of *degQ* resulted in severe growth defects at
15 higher temperatures, indicating that DegQ might act as a major protease for protein
16 quality control in the periplasm (Dai et al. 2015). It is therefore important to clarify the
17 relationship between OMV production and periplasmic protease DegQ in *S. oneidensis*.

18 In this study, we investigated the relationship between DegQ and OMV production
19 in *S. oneidensis* from the perspective of envelope stress. DegQ was confirmed as a
20 functional serine protease by using purified DegQ. In addition, the *degQ*-deficient *S.*
21 *oneidensis* mutant was compared with the wild type to identify differences in amount
22 and level of protease activity in the periplasmic proteins. OMV production was
23 examined, and the proteins associated with the isolated OMVs from *S. oneidensis* were

1 identified by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE)
2 and nano liquid chromatography-tandem mass spectrometry (LC-MS/MS) analyses.

3

4 **Materials and methods**

5 **Bacterial strains and culture conditions**

6 The strains and plasmids used in this study are listed in Table 1. For genetic
7 manipulations, *E. coli* and *S. oneidensis* strains were cultured in lysogeny broth (LB)
8 medium (10 g L⁻¹ Hipolypepton (Wako Pure Chemical Industries), 5 g L⁻¹ Bacto-yeast
9 extract, and 10 g L⁻¹ NaCl) at 37 and 30°C, respectively. The culture media of strains
10 harboring the plasmid were supplemented with 25 mg L⁻¹ chloramphenicol and 1 mM
11 isopropyl thiogalactoside (IPTG) when necessary.

12 All *S. oneidensis* strains were precultured in LB medium for 18 h at 30°C, and then
13 inoculated into 80 mL of fresh LB medium in a flask to give an optical density at 660
14 nm (OD₆₆₀) = 0.01. The cultures were placed on a rotary shaker (NR-20, Taitec) with
15 shaking at 120 strokes per minute. Cell growth was recorded on a basis of dry cell
16 weight (DCW) by measuring changes in OD₆₆₀.

17

18 **In-frame deletion mutagenesis**

19 The *degQ* (SO3942) gene was deleted in-frame from the *S. oneidensis* genome as
20 described previously (Fennessey et al. 2010; Gao et al. 2010; Sundararajan et al. 2011).
21 Briefly, a 592-bp fragment of the region upstream of the open reading frame (ORF), a
22 364-bp fragment of *degQ* (from 989 - 1353 bp), and a 292 bp fragment of the region
23 downstream of the ORF were amplified by PCR, generating fragments F1 and F2,
24 which were fused by overlap extension PCR to generate fragment F3 (see

1 Supplementary Fig. S1). The primers used for construction are listed in Supplementary
2 Table S1. Fragment F3 was cloned into suicide plasmid pRE112 (Edwards et al. 1998)
3 using *SacI* and *XbaI* restriction endonucleases, and then transformed into JM109 λ pir
4 (Penfold and Pemberton 1992) by the calcium chloride method. The resulting plasmid,
5 pRE112-F3, was transferred into recipient *S. oneidensis* MR-1 cells by conjugation from
6 *E. coli* SM10 λ pir (Miller and Mekalanos 1988). Integration of the mutagenesis
7 construct into the chromosome was induced to generate the final deletion strain. Gene
8 deletion was verified by PCR using the F1-F and F2-R primers. The resulting mutation
9 was confirmed as an in-frame deletion that removed approximately 1000 bp of the *degQ*
10 gene. The mutant strain was named $\Delta degQ$.

11

12 **Knock-in complementation analysis**

13 The *degQ* region (SO3942) was amplified from *S. oneidensis* MR-1 genomic DNA
14 using primer pair *degQ*-F and R (see Supplementary Table S1). The amplicon was
15 ligated into pHSG399 at the *XhoI* and *BamHI* restriction sites to generate recombinant
16 plasmid pHSG399-*degQ*. The purified pHSG399-*degQ* (approximately 2 μ g) was
17 transformed into $\Delta degQ$ cells by electroporation using 0.2-cm Gene Pulser/MicroPulser
18 cuvettes (Bio-Rad) and a Bio-Rad XPulser. Immediately after transformation, 0.5 mL of
19 super optimal broth with catabolite repression (SOC) medium devoid of antibiotics was
20 added to the cell suspension. Cells were transferred into a sterile culture tube and
21 incubated at 30°C for 1 h with continuous shaking. Following incubation, the cells were
22 spread onto LB agar plates supplemented with 25 μ g/mL chloramphenicol and
23 incubated at 30°C for 2–3 days. The complementation strain was named $\Delta degQ+degQ$.

24

1 **Characterization of DegQ**

2 Plasmid pHSG399-*degQ*-His was also constructed to express DegQ with a His6-tag
3 sequence at the C-terminus. The plasmid was transformed into wild-type *S. oneidensis*
4 MR-1 cells by electroporation, and the resulting strain was named WT+*degQ*-His. The
5 cells were pre-cultured and then inoculated into 80 mL of LB medium containing 25
6 $\mu\text{g/mL}$ chloramphenicol and incubated with shaking at 30°C. When $\text{OD}_{660} = 0.5$, IPTG
7 was added to the cultures at a final concentration of 1 mM, and the cultures were further
8 incubated for 20 h. The cells were harvested by centrifugation at 4,000 $\times g$ for 10 min at
9 4 °C. The cell pellets were resuspended in phosphate-buffered saline (PBS; pH7.4) and
10 disrupted by ultrasonication. The supernatant was recovered by centrifugation at 2,800
11 $\times g$ for 10 min at 4 °C, followed by additional centrifugation at 10,000 $\times g$ for 10 min at
12 4 °C. The supernatant was purified by using a Bio-Scale Mini Profinity IMAC cartridge
13 as well as a Profinia protein purification system (Bio-Rad Laboratories Inc.) based on
14 the affinity of the His6-tag, according to the manufacturer's protocol. The concentration
15 of purified DegQ was determined by a Pierce BCA Protein Assay kit (Thermo Fisher
16 Scientific Inc.) using bovine serum albumin as a standard. A 5- μL aliquot of purified
17 protein was analyzed by sodium dodecyl sulfate polyacrylamide gel electrophoresis
18 (SDS-PAGE) and visualized by Coomassie Blue staining. For western blotting, protein
19 was transferred from the gel to a membrane sheet of Hybond P (GE Healthcare Ltd.)
20 using the semi dry transfer method. Hybridization was conducted using an anti-His6
21 primary antibody and an ECL Western Blotting Starter kit (GE Healthcare Ltd.)
22 according to the manufacturer's protocol. Hybridization signals were detected using a
23 ChemiDoc imaging system (Bio-Rad Laboratories Inc.).

24 Protease activity was determined as described previously (Secades and Guijarro

1 1999) using azocasein as a substrate, with slight modifications. A 120- μ L solution of
2 purified DegQ containing 0.12 mg protein was mixed with 480 μ L of 0.1% (w/v)
3 azocasein in 25 mM Tris-HCl buffer (pH 7.6) with 5 mM MgCl₂, and the mixture was
4 incubated for 30 min at 30°C. When necessary, diisopropyl fluorophosphate (DFP) was
5 added as a serine protease-specific inhibitor. A 600- μ L volume of 10% (v/w)
6 trichloroacetic acid was added to terminate the reaction, and samples were incubated for
7 30 min on ice, followed by centrifugation at 15,000 $\times g$ for 10 min at 4°C. An aliquot of
8 supernatant (800 μ L) was mixed with 200 μ L of 1.8 M NaOH, and the absorbance at
9 440 nm (A_{440}) was measured using an ultraviolet-visible (UV-Vis) spectrometer
10 (UV-2600 PC, Shimadzu Corp.). One unit of enzyme activity was defined as the
11 amount of enzyme that yielded an increase in A_{440} of 0.01 per minute at 30°C.

12

13 **Gene expression analysis**

14 To examine gene expression, each strain was cultured in 80 mL of LB medium at
15 30°C and then harvested at 4 h post-inoculation by centrifugation at 4°C for 10 min at 5
16 000 $\times g$. Total RNA was extracted from the collected cells as described elsewhere
17 (Nguyen et al. 2014), and then reverse-transcribed into cDNA using a PrimeScript RT
18 reagent kit (Takara Bio Inc.) with random hexamer primers. Gene expression was
19 analyzed by real-time PCR, as described in our previous study (Nguyen et al. 2014).
20 The gene expression level was normalized against that of *rrsA* (16S rRNA). The
21 specific primer pairs are listed in Table S1.

22

23 **Isolation of periplasmic proteins and assay for proteolytic activity**

24 Periplasmic proteins from *S. oneidensis* were extracted according to the previously

1 reported method (Quan et al. 2013) with some modifications. *S. oneidensis* cells were
2 harvested from 80 ml of culture broth at 6 h post-inoculation by centrifugation at 3,000
3 $\times g$ for 20 min at 4°C. Precipitant was gently resuspended in 2 mL of Tris-sucrose-EDTA
4 (TSE) buffer (200 mM Tris-HCl buffer containing 500 mM sucrose and 1 mM EDTA,
5 pH 8.0) using a wire loop. The cells were then incubated on ice for 30 min, and the
6 supernatant containing envelope extract was collected by centrifugation at 16,000 $\times g$ for
7 30 min at 4°C. Finally, the periplasmic fraction was isolated as a supernatant by
8 ultracentrifugation at 100,000 $\times g$ (CS100FNX, Hitachi) for 1 h at 4°C. Protein
9 concentrations were determined by the Bradford assay, with bovine serum albumin as a
10 standard.

11 Protease activity of periplasmic proteins was also determined using azocasein, as
12 described above. The periplasmic fraction from the sample (120 μL) was mixed with
13 480 μL of 0.1% (w/v) azocasein in 25 mM Tris-HCl buffer (pH 7.6) containing 5 mM
14 MgCl_2 .

15

16 **Isolation and transmission electron microscope (TEM) observation of OMVs**

17 OMVs were isolated as previously described (Gujrati et al. 2014) with some
18 modifications. Following incubation for 24 h, *S. oneidensis* culture broth (80 mL) was
19 centrifuged at 3,970 $\times g$ for 10 min at 4°C. The supernatant was then passed through a
20 0.45 μm pore-size filter. The contents were precipitated using ammonium sulfate (final
21 concentration, 400 g L^{-1}) at room temperature for 1 h. The crude OMVs obtained by
22 centrifugation at 12,450 $\times g$ for 30 min at 4°C were dissolved in 500 μL of 15% (v/v)
23 glycerol, and then concentrated by ultracentrifugation (CS100FNX, Hitachi) at 109,000
24 $\times g$ for 1 h. The OMV pellets were resuspended in 100 μL of 15% (v/v) glycerol solution.

1 As a result, the OMV samples were concentrated 800–fold as compared with the
2 original culture broth. For observation of flocs using a TEM (H6000, Hitachi), the OMV
3 samples were dropped onto a mesh copper grid and negatively stained with 4% uranyl
4 acetate.

5

6 **SDS-PAGE and nano LC-MS/MS analyses of OMVs**

7 A 5- μ L sample of isolated OMVs from each strain was analyzed by SDS-PAGE
8 with Coomassie Blue staining. OMV production was quantified according to the
9 previously described method (Schwechheimer and Kuehn 2013) with some
10 modifications. The SDS-PAGE band at \sim 37 kDa was analyzed by densitometry (NIH
11 Image J software) as an index of OMV concentration. The OMV density value was
12 divided by dry cell weight (DCW) in each culture to calculate cell-based OMV
13 production, which was then normalized against OMV production of the wild-type strain.
14 The 37-kDa band was cut from the gel and subjected to in-gel digestion with trypsin.
15 The proteins were analyzed using nano-flow liquid chromatography with online tandem
16 mass spectrometry (nanoLC/ESI MS/MS system) composed of an LTQ Orbitrap Velos
17 (Thermo Fisher Scientific Inc.) coupled with a nanoLC (Advance, Michrom
18 BioResources) and an HTC-PAL autosampler (CTC Analytics), at the Core
19 Instrumentation Facility of the Research Institute for Microbial Diseases, Osaka
20 University (Osaka, Japan). Tandem mass spectra were acquired automatically, and then
21 searched against an *S. oneidensis* database from the National Center for Biotechnology
22 Information using the Mascot Server (Matrix Science).

23

24 **Results and Discussion**

1 **Characterization of DegQ**

2 The His6-tagged DegQ protein expressed in the *S. oneidensis* WT+*degQ*-His strain
3 was purified and then analyzed by SDS-PAGE. DegQ was detected as a major band at a
4 molecular weight of 45 kDa (Fig. 1A), corresponding to the expected size of DegQ
5 protein. This indicated that the His6-tagged DegQ was successfully expressed in the
6 form of soluble molecules in the WT+*degQ*-His cells. In addition, western blot analysis
7 using anti-His primary antibody was conducted to further confirm the presence of DegQ.
8 A clear signal at ~ 45 kDa was observed (Fig. 1B), confirming that the purified protein
9 is DegQ.

10 The protease activity of purified DegQ was confirmed by the azocasein hydrolysis
11 assay. As shown in Fig. 1C, the activity of DegQ was 5.4 U mg-protein⁻¹. As DegQ is
12 thought to be a putative periplasmic serine protease, the effect of DFP, a serine
13 protease-specific inhibitor (Swamy et al. 1983), on the activity of DegQ was examined.
14 As expected, 20 mM DFP inhibited protease activity of DegQ by 85%, suggesting that
15 the purified protein was a serine protease. Thus, these results confirmed that the *S.*
16 *oneidensis degQ* gene product is a serine protease.

17

18 **Deletion and complementation of *degQ* in *S. oneidensis***

19 To confirm the in-frame deletion of *degQ*, genomic DNA was extracted from wild
20 type *S. oneidensis* and the Δ *degQ* mutant. The *degQ* gene (1353 bp) plus 592 bp and
21 292 bp of the upstream and downstream regions, respectively, were amplified by PCR.
22 Electrophoresis results are shown in Supplementary Fig. S1. An approximately 2.2-kbp
23 fragment was amplified from the wild type cells, corresponding to the combined size of
24 *degQ* plus the surrounding regions. In contrast, an ~ 900 bp band was amplified from

1 the putative $\Delta degQ$ mutant. Furthermore, real-time PCR analysis did not detect *degQ*
2 mRNA expression from the mutant cells (Table 2). These results indicate the successful
3 deletion of *degQ* from the chromosome of *S. oneidensis*.

4 The complete *degQ* gene was expressed and re-introduced into the $\Delta degQ$ mutant
5 using the pHSG399-*degQ* plasmid. The relative expression of *degQ* (normalized
6 against that of *rrsA*) in the resultant $\Delta degQ+degQ$ cells at 4 h post-inoculation was
7 1.5×10^{-2} , which was much higher than the expression in wild type cells (6.3×10^{-5})
8 (Table 2). This increased expression compared with the wild type can be attributed to
9 the high copy number of pHSG399 plasmid (Takeshita et al. 1987).

10

11 **Periplasmic protease activity**

12 Figure 2 shows representative growth curves of the wild type *S. oneidensis*,
13 deletion mutant $\Delta degQ$, and complemented mutant $\Delta degQ+degQ$ strains. The wild type
14 and $\Delta degQ$ strains showed similar growth profiles (average growth rate of about 0.6 g-
15 DCW h⁻¹ during 6-12 h). The cell densities of the wild type and $\Delta degQ$ cultures at 24 h
16 post-inoculation reached DCW = 6.9 and 7.8 g L⁻¹, respectively. It can be mentioned
17 that the deletion of *degQ* did not repress cell growth at 30°C. Meanwhile, the growth of
18 $\Delta degQ+degQ$ cells appeared to be slightly slower at 12 h than the other two strains,
19 probably owing to the load of the high-copy number plasmid. The cell density of
20 $\Delta degQ+degQ$ reached DCW = 6.7 g L⁻¹ at 24 h. On the whole, these results suggest that
21 the deletion and complementation of *degQ* did not drastically influence the growth of *S.*
22 *oneidensis*.

23 Next, cells of each strain were harvested at exponential growth phase (culture
24 time of 6 h post-inoculation) for isolation of periplasmic proteins and analysis of

1 protease activity. Table 2 shows protease activity on a DCW basis, as well as the
2 concentration of periplasmic proteins obtained from each strain. The specific protease
3 activity of the periplasmic proteins from the wild type was 19.4 U g-DCW⁻¹, while the
4 activity of the $\Delta degQ$ cells was 7.0 U g-DCW⁻¹, corresponding to only 30% of that of
5 the wild type. Thus, the periplasmic protease activity was significantly lowered by
6 deletion of *degQ*. In contrast, the complemented mutant, $\Delta degQ+degQ$, had the highest
7 protease activity of 61.3 U g-DCW⁻¹. This corresponded with the increased *degQ*
8 mRNA expression observed in $\Delta degQ+degQ$ cells compared with the wild type.

9 Interestingly, the concentration of periplasmic proteins from the $\Delta degQ$ cells
10 was 2.56 mg g-DCW⁻¹, which was approximately 2.5 times greater than that from the
11 wild type cells (1.03 mg g-DCW⁻¹). This result strongly suggests that deletion of *degQ*
12 induced protein accumulation in the periplasmic space of *S. oneidensis* cells. In contrast,
13 the complemented mutant $\Delta degQ+degQ$ cells produced the lowest concentration of
14 periplasmic proteins (0.44 mg g-DCW⁻¹), which corresponds with the increased
15 protease activity of this strain. Thus, our results strongly suggest that DegQ is a
16 dominant periplasmic protease of *S. oneidensis*, and that lack of DegQ induces protein
17 accumulation in the periplasmic space.

18

19 **Comparison of OMV production among strains**

20 As in the case of *E. coli* (McBroom and Kuehn 2007; Schwechheimer and Kuehn
21 2013), it is likely that accumulation of periplasmic proteins in *S. oneidensis* cells would
22 induce OMV production, owing to envelope stress. Therefore, the insoluble fraction of
23 the supernatant containing OMVs was obtained from wild type, $\Delta degQ$, and
24 $\Delta degQ+degQ$ culture broths at a culture time of 24 h by ultracentrifugation. The

1 extracted OMVs from each strain were observed by TEM with negative staining (Fig. 3).
2 All strains produced OMVs, and while OMVs from the wild type culture tended to
3 aggregate, there was no significant difference in the size of OMVs between the wild
4 type and $\Delta degQ$ strains. On the other hand, fewer OMVs were obtained from the
5 $\Delta degQ+degQ$ cells compared with the other strains, indicating reduced OMV
6 production. In addition, co-precipitating appendages such as fimbriae and flagella were
7 confirmed in the sample from the $\Delta degQ+degQ$ cells.

8 To compare OMV production, the fractions containing OMVs from each *S.*
9 *oneidensis* strain were analyzed by SDS-PAGE and qualitatively compared with the
10 wild type. The protein bands in the sedimented fractions from the $\Delta degQ$ cells were
11 much more intense (Fig. 4A), while the bands from the $\Delta degQ+degQ$ cells were much
12 weaker. These results suggested that deletion of *degQ* enhanced OMV production, as
13 was observed in *E. coli* (McBroom et al. 2006). However, it has been reported that a
14 portion of the periplasmic proteins is included inside the OMV structure when OMVs
15 are generated (Lee et al. 2007). Therefore, we cannot exclude the possibility that the
16 more intense bands from the $\Delta degQ$ sample is a consequence of increased amounts of
17 periplasmic proteins inside the OMVs, rather than being indicative of enhanced OMV
18 production.

19 From this perspective, in the case of *E. coli*, the bands observed in the sample at
20 ~37 kDa (OmpF, OmpC, and OmpA) could be used as an index of OMV concentration,
21 as these membrane proteins are expressed specifically and abundantly within the outer
22 membrane of cells, and thus, are also found in abundance in OMVs (Schwechheimer
23 and Kuehn 2013). Therefore, the same type of index could be used for evaluation of
24 OMV production by *S. oneidensis*. As shown in Fig. 4A, a prominent band with a

1 molecular mass of ~37 kDa was confirmed from all strains by SDS-PAGE analysis. The
2 proteins contained within this band were then examined by nano LC-MS/MS analysis.
3 Table 3 summarizes the properties of 10 proteins in descending order according to
4 matching score from the first analysis. Among these proteins, OmpS38 (SO3896) had
5 the highest score. Omp35 is a putative porin on the outer membrane of *S. oneidensis*
6 cells that was confirmed by subcellular fraction analysis (Maier and Myers 2004).
7 OmpS38 is upregulated under anaerobic conditions and is involved in respiration of
8 non-oxygen electron acceptors (Gao et al. 2015). The protein with the next highest score
9 was also identified as a putative outer membrane porin (SO1821) (Heidelberg et al.
10 2002). These two outer membrane porins were confirmed to have the highest scores
11 during the second analysis run (Table 3). MotA, a flagellar motor transmembrane
12 channel protein, had the third highest score, while, the fourth protein (SO3545) was
13 OmpA, which is also outer membrane porin (Gao et al. 2015). Thus, outer membrane
14 proteins dominated the OMV samples, indicating that the bands at ~37 kDa can be used
15 as an index of OMV production in *S. oneidensis*. Taking these findings into account, the
16 OMV production was quantitatively analyzed based on the densitometry of the band at
17 ~37 kDa from each strain. As a result, we determined that OMV production by the
18 $\Delta degQ$ strain was about five times greater than that of the wild type strain (Fig. 4B). In
19 contrast, the complemented $\Delta degQ + degQ$ strain showed about a 60% reduction in
20 OMV production compared with the wild type strain. Together with the data from
21 periplasmic protease activity analysis and periplasmic protein concentration, our results
22 suggest that OMV production was promoted by the deletion of *degQ* in *S. oneidensis*
23 cells as a result of the accumulation of excessive amounts of periplasmic proteins.

1 In conclusion, our results show that *degQ* encodes a serine protease in *S.*
2 *oneidensis*. Deletion of *degQ* resulted in the accumulation of periplasmic proteins, and
3 led to a significant decrease in protease activity within the periplasmic space.
4 SDS-PAGE analysis of isolated OMVs showed the band corresponding to outer
5 membrane porins was more intense in the $\Delta degQ$ cells. Thus, OMV production by *S.*
6 *oneidensis* was promoted by the lack of DegQ, most likely as a result of an increase in
7 envelope stress.

8

9 **Acknowledgement**

10 This study was supported in part by a grant from the Noda Institute for Scientific
11 Research. We thank Dr. Muranaka at the Research Center for Ultra-High Voltage
12 Electron Microscopy, Osaka University, Japan, for assistance with TEM imaging.

13

14 **References**

- 15 Belchik SM, Kennedy DW, Dohnalkova AC, Wang Y, Sevinc PC, Wu H, Lin Y, Lu HP,
16 Fredrickson JK, Shi L (2011) Extracellular reduction of hexavalent chromium by
17 cytochromes MtrC and OmcA of *Shewanella oneidensis* MR-1. *Appl Environ*
18 *Microbiol* 77:4035-4041 doi: 10.1128/aem.02463-10
- 19 Beveridge TJ (1999) Structures of gram-negative cell walls and their derived membrane
20 vesicles. *J Bacteriol* 181:4725-4733
- 21 Dai J, Wei H, Tian C, Damron FH, Zhou J, Qiu D (2015) An extracytoplasmic function
22 sigma factor-dependent periplasmic glutathione peroxidase is involved in
23 oxidative stress response of *Shewanella oneidensis*. *BMC Microbiol* 15:34 doi:
24 10.1186/s12866-015-0357-0

1 Edwards RA, Keller LH, Schifferli DM (1998) Improved allelic exchange vectors and
2 their use to analyze 987P fimbria gene expression. *Gene* 207:149-157

3 Fennessey CM, Jones ME, Taillefert M, DiChristina TJ (2010) Siderophores are not
4 involved in Fe(III) solubilization during anaerobic Fe(III) respiration by
5 *Shewanella oneidensis* MR-1. *Appl Environ Microbiol* 76:2425-2432 doi:
6 10.1128/aem.03066-09

7 Fredrickson JK, Romine MF, Beliaev AS, Auchtung JM, Driscoll ME, Gardner TS,
8 Nealson KH, Osterman AL, Pinchuk G, Reed JL, Rodionov DA, Rodrigues JL,
9 Saffarini DA, Serres MH, Spormann AM, Zhulin IB, Tiedje JM (2008) Towards
10 environmental systems biology of *Shewanella*. *Nat Rev Microbiol* 6:592-603 doi:
11 10.1038/nrmicro1947

12 Gao H, Barua S, Liang Y, Wu L, Dong Y, Reed S, Chen J, Culley D, Kennedy D, Yang Y,
13 He Z, Nealson KH, Fredrickson JK, Tiedje JM, Romine M, Zhou J (2010) Impacts
14 of *Shewanella oneidensis* c-type cytochromes on aerobic and anaerobic respiration.
15 *Microb Biotechnol* 3:455-466 doi: 10.1111/j.1751-7915.2010.00181.x

16 Gao T, Ju L, Yin J, Gao H (2015) Positive regulation of the *Shewanella oneidensis*
17 OmpS38, a major porin facilitating anaerobic respiration, by Crp and Fur.
18 *Scientific Reports* 5:14263 doi: 10.1038/srep14263

19 Gujrati V, Kim S, Kim SH, Min JJ, Choy HE, Kim SC, Jon S (2014) Bioengineered
20 bacterial outer membrane vesicles as cell-specific drug-delivery vehicles for
21 cancer therapy. *ACS Nano* 8:1525-1537 doi: 10.1021/nn405724x

22 Heidelberg JF, Paulsen IT, Nelson KE, Gaidos EJ, Nelson WC, Read TD, Eisen JA,
23 Seshadri R, Ward N, Methe B, Clayton RA, Meyer T, Tsapin A, Scott J, Beanan M,
24 Brinkac L, Daugherty S, DeBoy RT, Dodson RJ, Durkin AS, Haft DH, Kolonay JF,

1 Madupu R, Peterson JD, Umayam LA, White O, Wolf AM, Vamathevan J,
2 Weidman J, Impraim M, Lee K, Berry K, Lee C, Mueller J, Khouri H, Gill J,
3 Utterback TR, McDonald LA, Feldblyum TV, Smith HO, Venter JC, Neilson KH,
4 Fraser CM (2002) Genome sequence of the dissimilatory metal ion-reducing
5 bacterium *Shewanella oneidensis*. Nat Biotechnol 20:1118-1123 doi:
6 10.1038/nbt749

7 Kadurugamuwa JL, Beveridge TJ (1995) Virulence factors are released from
8 *Pseudomonas aeruginosa* in association with membrane vesicles during normal
9 growth and exposure to gentamicin: a novel mechanism of enzyme secretion. J
10 Bacteriol 177:3998-4008

11 Kadurugamuwa JL, Beveridge TJ (1996) Bacteriolytic effect of membrane vesicles
12 from *Pseudomonas aeruginosa* on other bacteria including pathogens:
13 conceptually new antibiotics. J Bacteriol 178:2767-2774

14 Knox KW, Vesik M, Work E (1966) Relation between excreted lipopolysaccharide
15 complexes and surface structures of a lysine-limited culture of *Escherichia coli*. J
16 Bacteriol 92:1206-1217

17 Lee EY, Bang JY, Park GW, Choi DS, Kang JS, Kim HJ, Park KS, Lee JO, Kim YK,
18 Kwon KH, Kim KP, Gho YS (2007) Global proteomic profiling of native outer
19 membrane vesicles derived from *Escherichia coli*. Proteomics 7:3143-3153 doi:
20 10.1002/pmic.200700196

21 Lipinska B, Zylicz M, Georgopoulos C (1990) The HtrA (DegP) protein, essential for
22 *Escherichia coli* survival at high temperatures, is an endopeptidase. J Bacteriol
23 172:1791-1797

24 Maier TM, Myers CR (2004) The outer membrane protein Omp35 affects the reduction

1 of Fe(III), nitrate, and fumarate by *Shewanella oneidensis* MR-1. BMC Microbiol
2 4:23 doi: 10.1186/1471-2180-4-23

3 McBroom AJ, Johnson AP, Vemulapalli S, Kuehn MJ (2006) Outer membrane vesicle
4 production by *Escherichia coli* is independent of membrane instability. J Bacteriol
5 188:5385-5392 doi: 10.1128/JB.00498-06

6 McBroom AJ, Kuehn MJ (2007) Release of outer membrane vesicles by Gram-negative
7 bacteria is a novel envelope stress response. Mol Microbiol 63:545-558 doi:
8 10.1111/j.1365-2958.2006.05522.x

9 Miller VL, Mekalanos JJ (1988) A novel suicide vector and its use in construction of
10 insertion mutations: osmoregulation of outer membrane proteins and virulence
11 determinants in *Vibrio cholerae* requires *toxR*. J Bacteriol 170:2575-2583

12 Myers CR, Myers JM (1994) Ferric iron reduction-linked growth yields of *Shewanella*
13 *putrefaciens* MR-1. J Appl Bacteriol 76:253-258

14 Myers CR, Nealson KH (1988) Bacterial manganese reduction and growth with
15 manganese oxide as the sole electron acceptor. Science 240:1319-1321 doi:
16 10.1126/science.240.4857.1319

17 Myers CR, Nealson KH (1990) Respiration-linked proton translocation coupled to
18 anaerobic reduction of manganese(IV) and iron(III) in *Shewanella putrefaciens*
19 MR-1. J Bacteriol 172:6232-6238

20 Myers JM, Myers CR (2000) Role of the tetraheme cytochrome CymA in anaerobic
21 electron transport in cells of *Shewanella putrefaciens* MR-1 with normal levels of
22 menaquinone. J Bacteriol 182:67-75

23 Nguyen MH, Ojima Y, Sakka M, Sakka K, Taya M (2014) Probing of
24 exopolysaccharides with green fluorescence protein-labeled carbohydrate-binding

1 module in *Escherichia coli* biofilms and flocs induced by *bcsB* overexpression. J
2 Biosci Bioeng 118:400-405 doi: 10.1016/j.jbiosc.2014.03.005

3 Nowotny A, Behling UH, Hammond B, Lai CH, Listgarten M, Pham PH, Sanavi F
4 (1982) Release of toxic microvesicles by *Actinobacillus actinomycetemcomitans*.
5 Infect Immun 37:151-154

6 Penfold RJ, Pemberton JM (1992) An improved suicide vector for construction of
7 chromosomal insertion mutations in bacteria. Gene 118:145-146

8 Quan S, Hiniker A, Collet J-F, Bardwell CAJ (2013) Isolation of Bacteria Envelope
9 Proteins. In: Delcour AH (ed) *Bacterial Cell Surfaces : Methods and Protocols*.
10 Springer, pp 359-366

11 Rothfield L, Pearlman-Kothencz M (1969) Synthesis and assembly of bacterial
12 membrane components. A lipopolysaccharide-phospholipid-protein complex
13 excreted by living bacteria. J Mol Biol 44:477-492

14 Schwechheimer C, Kuehn MJ (2013) Synthetic effect between envelope stress and lack
15 of outer membrane vesicle production in *Escherichia coli*. J Bacteriol
16 195:4161-4173 doi: 10.1128/jb.02192-12

17 Schwechheimer C, Kuehn MJ (2015) Outer-membrane vesicles from Gram-negative
18 bacteria: biogenesis and functions. Nat Rev Microbiol 13:605-619 doi:
19 10.1038/nrmicro3525

20 Schwechheimer C, Sullivan CJ, Kuehn MJ (2013) Envelope control of outer membrane
21 vesicle production in Gram-negative bacteria. Biochem. 52:3031-3040 doi:
22 10.1021/bi400164t

23 Secades P, Guijarro JA (1999) Purification and characterization of an extracellular
24 protease from the fish pathogen *Yersinia ruckeri* and effect of culture conditions

1 on production. Appl Environ Microbiol 65:3969-3975

2 Strauch KL, Johnson K, Beckwith J (1989) Characterization of *degP*, a gene required
3 for proteolysis in the cell envelope and essential for growth of *Escherichia coli* at
4 high temperature. J Bacteriol 171:2689-2696

5 Sundararajan A, Kurowski J, Yan T, Klingeman DM, Joachimiak MP, Zhou J, Naranjo B,
6 Gralnick JA, Fields MW (2011) *Shewanella oneidensis* MR-1 sensory box protein
7 involved in aerobic and anoxic growth. Appl Environ Microbiol 77:4647-4656
8 doi: 10.1128/aem.03003-10

9 Swamy KH, Chung CH, Goldberg AL (1983) Isolation and characterization of protease
10 do from *Escherichia coli*, a large serine protease containing multiple subunits.
11 Arch Biochem Biophys 224:543-554


12 Takeshita S, Sato M, Toba M, Masahashi W, Hashimoto-Gotoh T (1987)
13 High-copy-number and low-copy-number plasmid vectors for lacZ
14 alpha-complementation and chloramphenicol- or kanamycin-resistance selection.
15 Gene 61:63-74

16 Wensink J, Witholt B (1981) Identification of different forms of the murein-bound
17 lipoprotein found in isolated outer membranes of *Escherichia coli*. Eur J Biochem
18 113:349-357

19 Zhou L, Srisatjaluk R, Justus DE, Doyle RJ (1998) On the origin of membrane vesicles
20 in Gram-negative bacteria. FEMS Microbiol Lett 163:223-228 doi:
21 10.1111/j.1574-6968.1998.tb13049.x

1 **FIGURE LEGENDS**


2 Fig. 1 Purification and characterization of DegQ protein. (A) SDS-PAGE analysis of
3 purified DegQ protein. DegQ was expressed in the *S. oneidensis* WT+*degQ*-His
4 strain. (B) Western blot analysis of purified DegQ protein. Hybridization was
5 conducted using anti-His6 primary antibody. (C) Protease activity of purified
6 DegQ protein with or without serine protease inhibitor, diisopropyl
7 fluorophosphates (DFP). Data was determined from three independent
8 experiments. Vertical bars indicate standard deviation. An asterisk indicates
9 statistical significance as determined by a Student's t-test ($p < 0.05$).



10

11


1 Fig. 2 Growth profiles of *S. oneidensis* wild type, $\Delta degQ$, and $\Delta degQ+degQ$ strains in
2 LB medium at 30°C. Data were determined from three independent experiments.
3 Vertical bars indicate standard deviation. DCW was calculated by $DCW = \alpha \times$
4 OD_{660} when $\alpha = 0.87$ (wild type), 0.96 ($\Delta degQ$), and 0.93 ($\Delta degQ+degQ$) $g L^{-1}$,
5 respectively.



6

7


1 Fig. 3 TEM images of OMVs isolated from *S. oneidensis* wild type, $\Delta degQ$, and
2 $\Delta degQ+degQ$ strains. The OMVs were stained with uranyl acetate. The scale
3 bars indicate 200 nm.



4

5

1 Fig. 4 Correlation between *degQ* and OMV production. (A) SDS-PAGE analysis of
2 OMVs isolated from the *S. oneidensis* wild type, $\Delta degQ$, and $\Delta degQ+degQ$
3 strains. The closed arrow indicates the protein bands of interest that were
4 analyzed by nano LC-MS/MS. (B) Comparison of OMV production among *S.*
5 *oneidensis* wild type, $\Delta degQ$, and $\Delta degQ+degQ$ strains. OMV production was
6 determined as relative values by normalizing against the value of the wild type
7 strain. Data was determined from three independent experiments. Vertical bars
8 indicate standard deviation. An asterisk indicates the statistical significance as
9 determined by ANOVA with Tukey's test ($p < 0.05$).




10

1 **SUPPORTING INFORMATION**

2 Table S1 Primers used in this study.

3 Fig. S1 Electrophoresis of the *degQ* fragment amplified from genomic DNA isolated
4 from *S. oneidensis* wild type (lane 1) and $\Delta degQ$ (lane 2) strains. Lane M;
5 *HindIII*-digested λ DNA marker.



6

Table 1

Table 1 Strains and vectors used in this study.

| Strain and vector | Description | Reference |
|------------------------------------|---|------------------------------|
| Strains | | |
| <i>S. oneidensis</i> MR-1 | Wild type strain | (Myers and Neilson 1988) |
| WT+ <i>degQ</i> -His | Wild type strain with pHSG399- <i>degQ</i> -His | This study |
| Δ <i>degQ</i> | In-flame deletion of <i>degQ</i> | This study |
| Δ <i>degQ</i> + <i>degQ</i> | Cm ^r , complemented strain with pHSG399- <i>degQ</i> | This study |
| <i>Escherichia coli</i> | | |
| SM10 λ pir | Mating strain | (Miller and Mekalanos 1988) |
| JM109 λ pir | Cloning strain | (Penfold and Pemberton 1992) |
| Plasmids | | |
| pRE112 | Cm ^r , pir-dependent suicide vector | (Edwards et al. 1998) |
| pRE112-F3 | Cm ^r , pRE112 caring F3 fragment | This study |
| pHSG399 | Cm ^r , expression vector | (Takeshita et al. 1987) |
| pHSG399- <i>degQ</i> | Cm ^r , expression vector carrying <i>degQ</i> | This study |
| pHSG399- <i>degQ</i> -His | Cm ^r , expression vector carrying <i>degQ</i> with the His6-Tag sequence at C-terminus | This study |

Table 2

Table 2 mRNA expressions of *degQ* gene, protease activities of DegQ protein, and amounts of periplasmic proteins from *S. oneidensis* wild type, $\Delta degQ$, and *degQ+degQ* strains.

| Strain | mRNA expression [–] | Protease activity [U g–DCW ⁻¹] | Periplasmic proteins [mg g–DCW ⁻¹] |
|------------------|--------------------------------|---|---|
| Wild type | $(6.3 \pm 4.2) \times 10^{-5}$ | 19.4 ± 2.9 | 1.03 ± 0.12 |
| $\Delta degQ$ | N.D. | 7.0 ± 2.3 | 2.56 ± 0.37 |
| <i>degQ+degQ</i> | $(1.5 \pm 1.2) \times 10^{-2}$ | 61.3 ± 14.6 | 0.44 ± 0.03 |

The data were determined using the cells harvested at culture times of 4 h for mRNA expression analysis and 6 h for protease activity assay. The data were determined from three independent experiments.

N.D.: Not detected

Table 3

Table 3 Proteins identified in OMVs from *S. oneidensis* wild type strain.

| Protein | Gene | Mass | Score | Peptide no. | Accession | Note |
|---|---------------|-------|-------|-------------|--------------|--------|
| First analysis | | | | | | |
| Outer membrane porin | <i>ompS38</i> | 39874 | 34463 | 1166 | gi 499386175 | SO3896 |
| Outer membrane porin | - | 39654 | 6912 | 327 | gi 499385907 | SO1821 |
| Flagellar motor protein MotA | <i>motA</i> | 48740 | 1097 | 36 | gi 499384378 | SO4287 |
| Outer membrane porin | <i>ompA</i> | 43970 | 1033 | 28 | gi 499384375 | SO3545 |
| iron ABC transporter substrate-binding protein | <i>fbpA</i> | 37394 | 953 | 30 | gi 499383481 | SO0744 |
| TonB-dependent ferric putrebactin siderophore receptor PutA | <i>putA</i> | 81373 | 877 | 20 | gi 499385445 | SO3033 |
| ATP synthase subunit beta | <i>atpD</i> | 49856 | 784 | 21 | gi 499386863 | SO4747 |
| alcohol dehydrogenase | <i>adhB</i> | 40387 | 783 | 23 | gi 499384108 | SO1490 |
| cytochrome C | <i>omcA</i> | 79992 | 725 | 27 | gi 499384335 | SO1779 |
| 5'-nucleotidase | <i>ushA</i> | 61353 | 671 | 29 | gi 499384520 | SO2001 |
| Second analysis | | | | | | |
| Outer membrane porin | <i>ompS38</i> | 39874 | 11280 | 488 | gi 499386175 | SO3896 |
| Outer membrane porin | - | 39654 | 1960 | 109 | gi 499385907 | SO1821 |
| alcohol dehydrogenase | <i>adhB</i> | 40387 | 709 | 28 | gi 499384108 | SO1490 |
| TonB-dependent receptor | <i>hmuA</i> | 81373 | 527 | 22 | gi 499385445 | SO3669 |
| Outer membrane porin | <i>ompA</i> | 43970 | 384 | 22 | gi 499384375 | SO3545 |
| Isocitrate dehydrogenase | - | 36173 | 370 | 13 | gi 499384148 | SO1538 |
| ABC transporter substrate-binding protein | <i>potF</i> | 40834 | 368 | 16 | gi 499383922 | SO1270 |
| Phage capsid protein | - | 43497 | 319 | 18 | gi 499385376 | SO2963 |
| Cytochrome C | <i>omcA</i> | 79992 | 316 | 12 | gi 499384335 | SO1779 |
| Iron ABC transporter substrate-binding protein | <i>fbpA</i> | 37394 | 299 | 16 | gi 499383481 | SO0744 |

Supplementary Table S1

Supplementary Table S1 Primers used in this study

| Primers | Sequence |
|----------------------------------|---|
| Mutagenesis | |
| F1-F | 5'- ATCGGAGCTCCA AAGAGTTAGGCTCTTCGGCT -3' |
| F1-R | 5'- GACTGGCTTAGGTCGTCTCTATTTTCGTTTTTCATCTATTCATA -3' |
| F2-F | 5'- AGAGACGACCTAAGCCAGTCCCAAGAGCTGCGTGCTAAAGTC -3' |
| F2-R | 5'- ATCGTCTAGAGCCTTGGCAAAGGAGAGTTCA -3' |
| Complementation | |
| <i>degQ</i> -F | 5'- ATCGAAGCTTGAAAACGAAATTATCT -3' |
| <i>degQ</i> -R | 5'- ATCGGAATTCTTAACGAAGCACTAA -3' |
| Purification | |
| <i>degQ</i> -His-R | 5'- ATCGGAATTCTCAGTGGTGGTGGTGGTGGTGACGAAGCACTAAGTAAAG -3' |
| Realtime PCR | |
| <i>rrsA</i> (house keeping)-RT-F | 5'- GCAACGCGAAGAACCTTACC -3' |
| <i>rrsA</i> (house keeping)-RT-R | 5'- CAGCACCTGTCTCACGGTTC-3' |
| <i>degQ</i> (SO3942)-RT-F | 5'- AACAAAGTGCAAGAGCGTCCA -3' |
| <i>degQ</i> (SO3942)-RT-R | 5'- CTTCACGGCCATCGTGTAAG -3' |