

Asymptotic behavior of least energy solutions to the Finsler Lane-Emden problem with large exponents

メタデータ	言語: English 出版者: American Institute of Mathematical Sciences 公開日: 2023-02-17 キーワード (Ja): キーワード (En): Finsler Lane-Emden problem, Finsler Laplacian, least energy solution, large exponent, blow up analysis 作成者: Habibi, Sadaf, 高橋, 太 メールアドレス: 所属: Osaka City University, Osaka Metropolitan University
URL	https://ocu-omu.repo.nii.ac.jp/records/2019811

Asymptotic behavior of least energy solutions to the Finsler Lane-Emden problem with large exponents

Habibi Sadaf, Futoshi Takahashi

Citation	Discrete and Continuous Dynamical Systems. 42(10); 5063-5086
Issue date	2022-10
Type	Journal Article
Textversion	Author
Rights	This article has been published in a revised form in Discrete and Continuous Dynamical Systems, https://doi.org/10.3934/dcds.2022086 . This version is free to download for private research and study only. Not for redistribution, re-sale or use in derivative works.
DOI	10.3934/dcds.2022086

Self-Archiving by Author(s)
Placed on: Osaka City University

ASYMPTOTIC BEHAVIOR OF LEAST ENERGY SOLUTIONS TO THE FINSLER LANE-EMDEN PROBLEM WITH LARGE EXPONENTS

SADAF HABIBI¹ AND FUTOSHI TAKAHASHI²

ABSTRACT. In this paper we are concerned with the least energy solutions to the Lane-Emden problem driven by an anisotropic operator, so-called the Finsler N -Laplacian, on a bounded domain in \mathbb{R}^N . We prove several asymptotic formulae as the nonlinear exponent gets large.

Key words: Finsler Lane-Emden problem, Finsler Laplacian, Least energy solution

2010 Mathematics Subject Classification: 35J60, 35J20

1. INTRODUCTION

Let $N \geq 2$ be an integer. In this paper, we study the following Lane-Emden problem driven by an anisotropic operator Q_N :

$$(1.1) \quad \begin{cases} -Q_N u = u^p & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where Ω is a smooth bounded domain in \mathbb{R}^N , $p > 1$ is any positive number, and Q_N is a quasilinear operator, so-called the *Finsler N -Laplacian*, defined by

$$Q_N u = \sum_{i=1}^N \frac{\partial}{\partial x_i} (H(\nabla u)^{N-1} H_{\xi_i}(\nabla u)).$$

Here $H \in C^2(\mathbb{R}^N \setminus \{0\})$ is any norm on \mathbb{R}^N and $H_{\xi_i}(\xi) = \frac{\partial H(\xi)}{\partial \xi_i}$. We assume that $H^N \in C^1(\mathbb{R}^N)$ and $\text{Hess}(H^N(\xi))$ is positive definite for

¹Department of Mathematics, Osaka City University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan.
e-mail:habibzaisadaf@gmail.com

²Department of Mathematics, Osaka City University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan.
e-mail:futoshi@sci.osaka-cu.ac.jp

Date: October 18, 2021.

1 any $\xi \in \mathbb{R}^N$, $\xi \neq 0$. Note that $Q_N u$ can be written as

$$Q_N u = \operatorname{div} \left(\nabla_\xi \left(\frac{1}{N} H(\xi)^N \right) \Big|_{\xi=\nabla u} \right) = \sum_{i,j=1}^N a_{ij}(\nabla u) \frac{\partial^2 u}{\partial x_i \partial x_j},$$

where $a_{ij}(\nabla u) = \operatorname{Hess}(\frac{1}{N} H^N(\xi))_{i,j} \Big|_{\xi=\nabla u}$. If $H(\xi) = |\xi|$ (the Euclidean norm), then $Q_N u$ coincides with the N -Laplacian $\Delta_N u = \operatorname{div}(|\nabla u|^{N-2} \nabla u)$ of a function u . In this case, the problem (1.1) was treated by Ren and Wei [21] [22] when $N = 2$, and in [23] for general $N \geq 2$. Ren and Wei [23] considered the least energy solution u_p of the following quasilinear problem

$$\begin{cases} -\Delta_N u = u^p & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

2 where Ω is a smooth bounded domain in \mathbb{R}^N . They studied the as-
 3 ymptotic behavior of u_p as the nonlinear exponent $p \rightarrow \infty$, and proved
 4 that the least energy solutions remain bounded in L^∞ -norm regardless
 5 of p . When the dimension $N = 2$, they showed that the least energy
 6 solutions must develop one “peak” in the interior of $\Omega \subset \mathbb{R}^2$, that is,
 7 the shape of graph of u_p looks like a single spike as $p \rightarrow \infty$. Moreover
 8 they showed that this peak point must be a critical point of the Robin
 9 function of the domain. For other generalizations of this problem to
 10 various situations, see for example, [28], [29], [30], [24], [25].

11 Now, main aim of the paper is to extend the results of Ren and Wei
 12 [21], [22], [23] to the anisotropic problem (1.1).

13 As in [21], [22], [23], we restrict our attention to the least energy
 14 solutions to (1.1) constructed as follows:

15 Consider the constrained minimization problem:

$$(1.2) \quad C_p = \inf \left\{ \int_\Omega H(\nabla u)^N dx : u \in W_0^{1,N}(\Omega), \int_\Omega |u|^{L^{p+1}(\Omega)} dx = 1 \right\}.$$

16 Since the Sobolev imbedding $W_0^{1,N}(\Omega) \hookrightarrow L^{p+1}(\Omega)$ is compact for any
 17 $p > 1$, we have at least one minimizer \underline{u}_p for the problem (1.2), where
 18 $\underline{u}_p \in W_0^{1,N}(\Omega)$, $\|\underline{u}_p\|_{p+1} = 1$. As $|\underline{u}_p| \in W_0^{1,N}(\Omega)$ also achieves C_p , we
 19 may assume $\underline{u}_p > 0$. Note that $Q_N(cu) = c^{N-1}Q_N(u)$ for a constant
 20 $c > 0$. Thus if we define

$$u_p = C_p^{\frac{1}{p+1-N}} \underline{u}_p,$$

21 then u_p solves (1.1) and $C_p = \int_\Omega H(\nabla u_p)^N dx / (\int_\Omega |u_p|^{p+1} dx)^{\frac{N}{p+1}}$. Stan-
 22 dard regularity argument implies that any weak solution $u \in W_0^{1,N}(\Omega)$

1 satisfies $u \in C^{1,\alpha}(\overline{\Omega})$ for some $\alpha \in (0, 1)$. We call u_p the *least energy*
 2 *solution* to (1.1).

3 Our first result is the following L^∞ -bound of least energy solutions.

4 **Theorem 1.1.** *Let u_p be a least energy solution to (1.1). Then there*
 5 *exist C_1, C_2 (independent of p), such that*

$$0 < C_1 \leq \|u_p\|_{L^\infty(\Omega)} \leq C_2 < \infty$$

6 *for p large enough.*

Furthermore, we have

$$\lim_{p \rightarrow \infty} p^{N-1} \int_{\Omega} H(\nabla u_p)^N dx = \lim_{p \rightarrow \infty} p^{N-1} \int_{\Omega} u_p^{p+1} dx = \left(\frac{Ne\beta_N}{N-1} \right)^{N-1}$$

7 *where $\beta_N = N(N\kappa_N)^{\frac{1}{N-1}}$, $\kappa_N = |\mathcal{W}|$ is the volume (with respect to the*
 8 *N -dimensional Hausdorff measure) of the unit Wulff ball associated*
 9 *with the dual norm H^0 of H :*

$$\mathcal{W} = \{x \in \mathbb{R}^N : H^0(x) < 1\}.$$

10 On the asymptotic behavior of the L^∞ -norm of u_p , we have

11 **Theorem 1.2.** *Let u_p be a least energy solution to (1.1). Then it holds*
 12 *that*

$$1 \leq \limsup_{p \rightarrow \infty} \|u_p\|_{L^\infty(\Omega)} \leq e^{\frac{N-1}{N}}.$$

13 To state further results, we need some definitions. Set

$$(1.3) \quad v_p = \frac{u_p}{\left(\int_{\Omega} u_p^p dx \right)^{\frac{1}{N-1}}}.$$

14 Then v_p is a weak solution of

$$(1.4) \quad \begin{cases} -Q_N v_p = f_p(x) = \frac{u_p^p}{\int_{\Omega} u_p^p dx} & \text{in } \Omega, \\ v_p > 0 & \text{in } \Omega, \\ v_p = 0 & \text{on } \partial\Omega. \end{cases}$$

15 For $X, Y \in \mathbb{R}^N$, $X, Y \neq 0$, $X \neq Y$, we put

$$d(X, Y) = \frac{(H^{N-1}(X)(\nabla_{\xi} H)(X) - H^{N-1}(Y)(\nabla_{\xi} H)(Y)) \cdot (X - Y)}{H^N(X - Y)},$$

16 and

$$(1.5) \quad d_N = \inf\{d(X, Y) \mid X, Y \in \mathbb{R}^N, X, Y \neq 0, X \neq Y\},$$

- 1 where $X \cdot Y = \sum_{j=1}^N X_j Y_j$ denotes the usual inner product for $X, Y \in$
 2 \mathbb{R}^N . As in [32] Lemma 5.1, we can obtain the estimate

$$\min\left\{\frac{\lambda}{\beta_N}, 1\right\} \leq d_N \leq 1$$

where λ is the least eigenvalue of $\text{Hess}\left(\frac{1}{N}H^N(\xi)\right)$, which is positive by the assumption (2.2) and β is as in (2.3), see §2. Also define

$$(1.6) \quad L_0 = \limsup_{p \rightarrow \infty} \frac{p \left(\int_{\Omega} u_p^p dx\right)^{\frac{1}{N-1}}}{\left(\frac{N}{N-1} e^{\frac{N-1}{N}}\right)}, \quad L_1 = d_N^{-\left(\frac{1}{N-1}\right)} L_0.$$

For a sequence v_{p_n} of v_p , we define the *blow-up set* S of $\{v_{p_n}\}$ as usual:

$$S = \{x \in \bar{\Omega} : \exists \text{ a subsequence } v_{p'_n}, \exists \{x_n\} \subset \Omega \text{ s.t. } x_n \rightarrow x \text{ and } v_{p'_n}(x_n) \rightarrow \infty\}.$$

- 3 In the following, $\sharp A$ denotes the cardinality of a set A and $[\cdot]$ denotes
 4 the Gauss symbol.

- 5 **Theorem 1.3.** *Let $\Omega \subset \mathbb{R}^N$ be a smooth bounded domain. Then for*
 6 *any sequence v_{p_n} of v_p with $p_n \rightarrow \infty$, the blow-up set S of v_{p_n} is non-*
 7 *empty. Also there exists a subsequence (still denoted by v_{p_n}) such that*
 8 *the estimate*

$$\sharp(S \cap \Omega) \leq \left\lceil \frac{e^{\frac{N-1}{N}}}{d_N} \right\rceil$$

- 9 *holds true for this subsequence.*

- 10 *Assume $S \cap \Omega = \{x_1, \dots, x_k\} \subset \Omega$. Then we have*

(i)

$$f_n = \frac{u_{p_n}^{p_n}}{\int_{\Omega} u_{p_n}^{p_n} dx} \overset{*}{\rightharpoonup} \sum_{i=1}^k \gamma_i \delta_{x_i}$$

- 11 *in the sense of Radon measures of Ω , where*

$$\gamma_i \geq \left(\frac{\beta_N}{L_1}\right)^{N-1}$$

- 12 *and $\sum_{i=1}^k \gamma_i \leq 1$.*

- 13 (ii) *$v_{p_n} \rightarrow G$ in $C_{loc}^1(\Omega \setminus (S \cap \Omega))$ for some function G satisfying*

$$\begin{cases} -Q_N G = 0 & \text{in } \Omega \setminus (S \cap \Omega), \\ G = +\infty & \text{on } S \cap \Omega, \\ G = 0 & \text{on } \partial\Omega \setminus (\partial\Omega \cap S). \end{cases}$$

- 14 (iii) *$\|u_{p_n}\|_{L^\infty(K)} \rightarrow 0$ as $n \rightarrow \infty$ for any compact set $K \subset \Omega \setminus (S \cap \Omega)$.*

1 In [23], Ren and Wei obtained an estimate of the number of interior
2 blow-up set

$$\sharp(S \cap \Omega) \leq \left[\frac{1}{d_N} \left(\frac{N}{N-1} \right)^{N-1} \right]$$

3 when $H(\xi) = |\xi|$ case. Since $e^x < \frac{1}{1-x}$ for $x \in (0, 1)$, we check that
4 $e^{\frac{N-1}{N}} < \left(\frac{N}{N-1} \right)^{N-1}$ for all $N \geq 2$. Thus the estimate in Theorem 1.3 is
5 better than that in [23] even when $H(\xi)$ coincides with the Euclidean
6 norm $|\xi|$. Also, Theorem 1.2 seems new even for $H(\xi) = |\xi|$ and $N > 2$
7 case.

8 Finally, we prove that if the blow-up set consists of one point, it
9 must be an interior point of Ω .

10 **Theorem 1.4.** *Assume $\sharp S = 1$ and $S = \{x_0\}$, $x_0 \in \overline{\Omega}$. Then $x_0 \in \Omega$
11 must hold.*

12 The organization of the paper is as follows: In §2, we recall basic
13 properties of the Finsler norm and collect useful lemmas about the
14 Finsler N -Laplacian. In §3, we obtain asymptotic formula for C_p as
15 $p \rightarrow \infty$, and prove the latter half part of Theorem 1.1. In §4, we prove
16 the L^∞ -bound of least energy solutions in Theorem 1.1. In §5, we prove
17 Theorem 1.2 using an argument by Adimurthi and Grossi [1]. In §6,
18 we prove Theorem 1.3. We use a notion of (L, δ) -regular, or irregular
19 points, which was originally introduced by Brezis and Merle [6]. Finally
20 in §7, we prove Theorem 1.4 by using a local Pohozaev identity and an
21 idea by Santra and Wei [25].

22 2. NOTATIONS AND BASIC PROPERTIES

23 Let H be any *norm* on \mathbb{R}^N , i.e., H is convex, $H(\xi) \geq 0$ and $H(\xi) = 0$
24 if and only if $\xi = 0$, and H satisfies

$$(2.1) \quad H(t\xi) = |t|H(\xi), \quad \forall \xi \in \mathbb{R}^N, \forall t \in \mathbb{R}.$$

25 By (2.1), H must be even: $H(-\xi) = H(\xi)$ for all $\xi \in \mathbb{R}^N$. Throughout
26 of the paper, we also assume that $H \in C^2(\mathbb{R}^N \setminus \{0\})$, $H^N \in C^1(\mathbb{R}^N)$,
27 and

$$(2.2) \quad \text{Hess} \left(H^N(\xi) \right) \text{ is positive definite for any } \xi \in \mathbb{R}^N, \xi \neq 0.$$

28 Since all norms on \mathbb{R}^N are equivalent to each other, we see the existence
29 of positive constants α and β such that

$$(2.3) \quad \alpha|\xi| \leq H(\xi) \leq \beta|\xi|, \quad \xi \in \mathbb{R}^N.$$

1 The dual norm of H is the function $H^0 : \mathbb{R}^N \rightarrow \mathbb{R}$ defined by

$$H^0(x) = \sup_{\xi \in \mathbb{R}^N \setminus \{0\}} \frac{\xi \cdot x}{H(\xi)}.$$

2 It is well-known that H^0 is also a norm on \mathbb{R}^N and satisfies the inequality
3

$$\frac{1}{\beta}|x| \leq H^0(x) \leq \frac{1}{\alpha}|x|, \quad \forall x \in \mathbb{R}^N.$$

4 The set

$$\mathcal{W} = \{x \in \mathbb{R}^N : H^0(x) < 1\}$$

5 is called the *Wulff ball*, or the H^0 -unit ball, and we denote $\kappa_N =$
6 $\mathcal{H}^N(\mathcal{W})$, where \mathcal{H}^N denotes the N -dimensional Hausdorff measure on
7 \mathbb{R}^N . We also denote $\mathcal{W}_r = \{x \in \mathbb{R}^N : H^0(x) < r\}$ for any $r > 0$.

8 For a domain $\Omega \subset \mathbb{R}^N$ and a Borel set $E \subset \mathbb{R}^N$, the *anisotropic*
9 *H -perimeter* of a set E with respect to Ω is defined as

$$P_H(E, \Omega) = \sup \left\{ \int_{E \cap \Omega} \operatorname{div} \sigma dx : \sigma \in C_0^\infty(\Omega, \mathbb{R}^N), H^0(\sigma(x)) \leq 1 \right\}.$$

10 If E is Lipschitz, then it holds $P_H(E, \Omega) = \int_{\Omega \cap \partial^* E} H(\nu) d\mathcal{H}^{N-1}$, where
11 $\partial^* E$ denotes the reduced boundary of the set E and $\nu(x)$ is the mea-
12 sure theoretic outer unit normal of $\partial^* E$ (see [16]). Also we have
13 $P_H(\mathcal{W}, \mathbb{R}^N) = N\kappa_N$. For more explanation about the anisotropic
14 perimeter, see [3] and [5].

15 Here we just recall some properties of H and H^0 . These will be
16 proven by using the homogeneity property of H and H^0 , see [4] Lemma
17 2.1, and Lemma 2.2.

18 **Proposition 2.1.** *Let H be a Finsler norm on \mathbb{R}^N . Then the following*
19 *properties hold true:*

- 20 (1) $|\nabla_\xi H(\xi)| \leq C$ for any $\xi \neq 0$.
- 21 (2) $\nabla_\xi H(\xi) \cdot \xi = H(\xi)$, $\nabla_x H(x) \cdot x = H(x)$ for any $\xi \neq 0$, $x \neq 0$.
- 22 (3) $(\nabla_\xi H)(t\xi) = \frac{t}{|t|} (\nabla_\xi H)(\xi)$ for any $\xi \neq 0$, $t \neq 0$.
- 23 (4) $H(\nabla H^0(x)) = 1$. $H^0(\nabla_\xi H(\xi)) = 1$.
- 24 (5) $H^0(x)(\nabla_\xi H)(\nabla_x H^0(x)) = x$.

Finally, given a smooth function u on \mathbb{R}^N , the *Finsler Laplace operator* of u (associated with H) is defined by

$$\begin{aligned} Qu(x) &= \operatorname{div} (H(\nabla u(x)) (\nabla_\xi H)(\nabla u(x))) \\ &= \sum_{j=1}^N \frac{\partial}{\partial x_j} \left(H(\xi) H_{\xi_j}(\xi) \Big|_{\xi=\nabla u(x)} \right) \end{aligned}$$

1 and, more generally, for any $1 < q < \infty$, the Finsler q -Laplace operator
 2 Q_q by

$$Q_q u(x) = \operatorname{div} \left(H^{q-1}(\nabla u(x)) (\nabla_\xi H)(\nabla u(x)) \right).$$

3 If we assume that $\operatorname{Hess}(H^q(\xi))$ is positive definite on $\mathbb{R}^N \setminus \{0\}$, Q_q
 4 becomes a uniformly elliptic operator locally. The Finsler q -Laplacian
 5 has been widely investigated in literature by many authors in different
 6 settings, see [2], [5], [8], [9], [10], [12], [13], [14], [17], [20] [34] and the
 7 references therein.

8 We collect here several useful facts.

9 **Theorem 2.2.** (Finsler Trudinger-Moser inequality [32]) *Let Ω be a*
 10 *bounded domain in \mathbb{R}^N , $N \geq 2$. Let $u \in W_0^{1,N}(\Omega)$ satisfy $\int_\Omega H(\nabla u)^N dx \leq$*
 11 *1. Then there exists a constant C depending only on the dimension N*
 12 *such that*

$$\int_\Omega \exp \left(\beta |u|^{\frac{N}{N-1}} \right) dx \leq C |\Omega|$$

13 *holds for any $\beta \leq \beta_N = N(N\kappa_N)^{\frac{1}{N-1}}$. Furthermore, β_N is opti-*
 14 *mal in the sense that there exists a sequence $\{u_n\} \subset W_0^{1,N}(\Omega)$ with*
 15 *$\int_\Omega H(\nabla u_n)^N dx \leq 1$, such that $\int_\Omega \exp \left(\beta |u_n|^{\frac{N}{N-1}} \right) dx \rightarrow +\infty$ as $n \rightarrow \infty$*
 16 *for $\beta > \beta_N$.*

17 Next is the unique existence of the Green function for the Finsler
 18 p -Laplacian.

19 **Theorem 2.3.** ([32]) *Let $\Omega \subset \mathbb{R}^N$ be a bounded domain containing the*
 20 *origin. Define $\Omega^* = \Omega \setminus \{0\}$ and*

$$\Gamma(x) = \begin{cases} C(p, N) (H^0(x))^{\frac{p-N}{p-1}} & \text{for } 1 < p < N, \\ C(N) \log \frac{1}{H^0(x)} & \text{for } p = N, \end{cases}$$

where $C(p, N) = \frac{p-N}{p-1} (N\kappa_N)^{-\frac{1}{p-1}}$ and $C(N) = (N\kappa_N)^{-\frac{1}{N-1}}$. Then
 there exists a unique function $G(\cdot, 0) \in C^{1,\alpha}(\Omega^*)$ with $|\nabla G| \in L^{p-1}(\Omega)$,
 $G/\Gamma \in L^\infty(\Omega)$, satisfying

$$\begin{cases} -Q_p G(\cdot, 0) = \delta_0 & \text{in } \Omega, \\ G(\cdot, 0) = 0 & \text{on } \partial\Omega. \end{cases}$$

21 Moreover, $g = G - \Gamma$ satisfies $g \in C(\Omega)$ and $\lim_{x \rightarrow 0} H^0(x) \nabla g(x) = 0$.

22 We recall here useful regularity estimates which are valid for the
 23 Finsler N -Laplacian equations, under the assumption (2.2); see Serrin
 24 [26], Tolksdorf [31], DiBenedetto [15] and Lieberman [19].

Theorem 2.4. *Let $\Omega \subset \mathbb{R}^N$ be a smooth bounded domain. Then the following statements are true.*

- (1) *Let $u \in W^{1,N}(\Omega)$ be a weak solution of $-Q_N u = f$ in Ω , where $f \in L^q(\Omega)$ for some $q > 1$. Then for any subdomain $\Omega' \subset\subset \Omega$, there exists a constant $C = C(\Omega, \Omega', q, N) > 0$ such that*

$$\|u\|_{L^\infty(\Omega')} \leq C (\|f\|_{L^q(\Omega)} + \|u\|_{L^N(\Omega)})$$

holds.

- (2) *Let $u \in W^{1,N}(\Omega)$ be a weak solution of $-Q_N u = f$ in Ω . Suppose $\|u\|_{L^\infty(\Omega)} \leq a$ and $\|f\|_{L^\infty(\Omega)} \leq b$ for some $a, b < \infty$. Then $u \in C_{loc}^{1,\alpha}(\Omega)$ for some $\alpha \in (0, 1)$ and for any subdomain $\Omega' \subset\subset \Omega$, there exists a constant $C = C(\Omega, \Omega', a, b, \alpha) > 0$ such that*

$$\|u\|_{C^{1,\alpha}(\Omega')} \leq C$$

holds. If, in addition, u satisfies the Dirichlet boundary condition $u = \phi$ on $\partial\Omega$ where $\phi \in C^{1,\beta}(\partial\Omega)$, $\beta \in (0, 1)$, then $u \in C^{1,\alpha}(\overline{\Omega})$ for some $\alpha \in (0, 1)$ holds.

- (3) (Harnack inequality) *Let $u \in W^{1,N}(\Omega)$ be a nonnegative weak solution of $-Q_N u = f$ in Ω . Suppose $\|f\|_{L^q(\Omega)} \leq b$ for some $q > 1$. Then for any subdomain $\Omega' \subset\subset \Omega$, there exists a constant $C = C(\Omega, \Omega', q, b) > 0$ such that*

$$\sup_{x \in \Omega'} u(x) \leq C \left(1 + \inf_{x \in \Omega'} u(x) \right)$$

holds.

Next is the result from [33] (Theorem 1.1 and Theorem 1.2).

Theorem 2.5. (Finsler Brezis-Merle type inequality [33]) *Let Ω be a bounded domain in \mathbb{R}^N , $N \geq 2$.*

- (1) *Suppose u is a weak solution to*

$$\begin{cases} -Q_N u = f(x) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $f \in L^1(\Omega)$. Then for any $\varepsilon \in (0, \beta_N)$ where $\beta_N = N(N\kappa_N)^{\frac{1}{N-1}}$, it holds that

$$\int_{\Omega} \exp \left(\frac{(\beta_N - \varepsilon)|u(x)|}{\|f\|_{L^1(\Omega)}^{\frac{1}{N-1}}} \right) dx \leq \frac{\beta_N}{\varepsilon} |\Omega|.$$

1 (2) Suppose u and v are weak solutions to

$$-Q_N u = f(x) > 0 \quad \text{in } \Omega$$

2 and

$$-Q_N v = 0 \quad \text{in } \Omega \quad v = u \quad \text{on } \partial\Omega,$$

3 respectively. Then for any $\varepsilon \in (0, \beta_N)$, we have

$$\int_{\Omega} \exp \left(\frac{(\beta_N - \varepsilon) d_N^{\frac{1}{N-1}} |u(x) - v(x)|}{\|f\|_{L^1(\Omega)}^{\frac{1}{N-1}}} \right) dx \leq \frac{|\Omega|}{\varepsilon},$$

4 where d_N is defined in (1.5).

5 Next is the Pohozaev identity for the Finsler q -Laplacian problem
 6 without the boundary condition. This is a special case of much more
 7 general identity proved in [11]. The identity below is known to hold
 8 for solutions in $C^1(\overline{\Omega}) \cap C^2(\Omega)$. The important point is that we can
 9 remove the condition $u \in C^2(\Omega)$ with the cost of the convexity and the
 10 $C^1(\mathbb{R}^N)$ -regularity of the map $\mathbb{R}^N \ni \xi \mapsto H^q(\xi)$. This improvement is
 11 crucial for the application to the Finsler Laplacian problem, since the
 12 best possible regularity result of solutions is $C^{1,\alpha}$, not C^2 , see Theorem
 13 2.4.

Theorem 2.6. ([11]) *Let $1 < q < \infty$. Let $u \in C^1(\overline{\Omega})$ be a weak solution of $-Q_q u = f(u)$ in Ω , where $\Omega \subset \mathbb{R}^N$ is a domain with the boundary of class C^1 , and $f \in C(\mathbb{R}, \mathbb{R})$. Assume the map $\mathbb{R}^N \ni \xi \mapsto H^q(\xi)$ is convex and belongs to $C^1(\mathbb{R}^N)$. Then the identity*

$$\begin{aligned} N \int_{\Omega} F(u) dx - \left(\frac{N-q}{q} \right) \int_{\Omega} H^q(\nabla u) dx \\ = \int_{\partial\Omega} F(u)(x-y) \cdot \nu(x) ds_x \\ - \frac{1}{q} \int_{\partial\Omega} H^q(\nabla u)(x-y) \cdot \nu(x) ds_x \\ + \int_{\partial\Omega} (H^{q-1}(\nabla u)(\nabla_{\xi} H)(\nabla u) \cdot \nu(x)) ((x-y) \cdot \nu(x)) ds_x \end{aligned}$$

14 holds true for any $y \in \mathbb{R}^N$. Here ν is the outer unit normal of $\partial\Omega$ and
 15 $F(s) = \int_0^s f(t) dt$.

Proof. Indeed, since $\mathcal{L}(x, s, \xi) = \frac{1}{q} H^q(\xi) - F(s)$ is of the “splitting” form, $F \in C^1(\mathbb{R})$, and $\xi \mapsto H^q(\xi)$ is convex and in $C^1(\mathbb{R}^N)$, Lemma 5, thus the equation (3) in [11] holds as it is. Also, if we do not impose

the boundary condition $u = 0$ on $\partial\Omega$ (and put $f = 0$ there) in Lemma 2 in [11], we obtain the identity

$$\begin{aligned} & \int_{\partial\Omega} \mathcal{L}(x, u, \nabla u)(h \cdot \nu) ds_x - \sum_{i,j=1}^N \int_{\partial\Omega} h_j D_{\xi_i} \mathcal{L}(x, u, \nabla u) D_{x_j} u \nu_i ds_x \\ &= \int_{\Omega} (\operatorname{div} h) \mathcal{L}(x, u, \nabla u) dx - \sum_{i,j=1}^N \int_{\Omega} D_i h_j D_{\xi_i} \mathcal{L}(x, u, \nabla u) D_{x_j} u dx \end{aligned}$$

1 for every $h \in C^1(\overline{\Omega}, \mathbb{R}^N)$. Inserting $h(x) = x$ leads to the claim. \square

2 Finally, we prove the following simple lemma.

3 **Lemma 2.7.** *Let $u \in W_0^{1,N}(\Omega)$ be a weak solution to $-Q_N u = f(u)$*
 4 *in $\Omega \subset \mathbb{R}^N$, where $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous. Let $a, c > 0$, $d \in \mathbb{R}$ and*
 5 *$b \in \mathbb{R}^N$. Then $v(x) = cu(ax + b) + d$, $x \in \Omega_{a,b} = \frac{\Omega - b}{a}$ is a weak solution*
 6 *to*

$$-Q_N v = a^N c^{N-1} f\left(\frac{v-d}{c}\right) \quad \text{in } \Omega_{a,b}, \quad v = 0 \quad \text{on } \partial\Omega_{a,b}.$$

Proof. For $x \in \Omega_{a,b}$, put $y = ax + b \in \Omega$. Then for any $\phi \in C_0^\infty(\Omega_{a,b})$, $\tilde{\phi}(y) = \phi(x)$ belongs to $C_0^\infty(\Omega)$. Therefore we have

$$\begin{aligned} & \int_{\Omega_{a,b}} H^{N-1}(\nabla v(x))(\nabla_\xi H)(\nabla v(x)) \cdot \nabla \phi(x) dx \\ &= \int_{\Omega_{a,b}} H^{N-1}(ca(\nabla u)(ax + b))(\nabla_\xi H)(ca(\nabla u)(ax + b)) \cdot \nabla \phi(x) dx \\ &= \int_{\Omega} c^{N-1} a^{N-1} H^{N-1}(\nabla u(y))(\nabla_\xi H)((\nabla u(y)) \cdot a \nabla \tilde{\phi}(y) a^{-N} dy \\ &= c^{N-1} \int_{\Omega} H^{N-1}(\nabla u(y))(\nabla_\xi H)(\nabla u(y)) \cdot \nabla \tilde{\phi}(y) dy \\ &= c^{N-1} \int_{\Omega} f(u(y)) \tilde{\phi}(y) dy \\ &= c^{N-1} \int_{\Omega_{a,b}} f\left(\frac{v(x)-d}{c}\right) \phi(x) a^N dx, \end{aligned}$$

where we have used (2.1) and Proposition 2.1 (3). Thus we see

$$\int_{\Omega_{a,b}} H^{N-1}(\nabla v(x))(\nabla_\xi H)(\nabla v(x)) \cdot \nabla \phi(x) dx = a^N c^{N-1} \int_{\Omega_{a,b}} f\left(\frac{v(x)-d}{c}\right) \phi(x) dx.$$

7 This holds true for any $\phi \in C_0^\infty(\Omega_{a,b})$, which implies Lemma. \square

3. ASYMPTOTIC ESTIMATE FOR C_p

In this section, first by using the Finsler Trudinger-Moser inequality Theorem 2.2, we establish the refined Sobolev embedding.

Lemma 3.1. *Let $\Omega \subset \mathbb{R}^N$ be a bounded domain. For any $t \geq 2$, there exists $D_t > 0$ such that for any $u \in W_0^{1,N}(\Omega)$,*

$$\|u\|_{L^t(\Omega)} \leq D_t t^{\frac{N-1}{N}} \|H(\nabla u)\|_{L^N(\Omega)}$$

holds true. Furthermore, we have

$$\lim_{t \rightarrow \infty} D_t = \left(\frac{1}{N \kappa_N^{1/N}} \right) \left(\frac{N-1}{Ne} \right)^{\frac{N-1}{N}}.$$

Proof. Let $u \in W_0^{1,N}(\Omega)$. By the elementary inequality $\frac{x^s}{\Gamma(s+1)} \leq e^x$ for $x \geq 0$ and $s \geq 0$, where $\Gamma(s)$ is the Gamma function, and the Finsler Trudinger-Moser inequality, we have

$$\begin{aligned} & \frac{1}{\Gamma(\frac{N-1}{N}t + 1)} \int_{\Omega} |u|^t dx \\ &= \frac{1}{\Gamma(\frac{N-1}{N}t + 1)} \int_{\Omega} \left(\beta_N \left(\frac{|u|}{\|H(\nabla u)\|_{L^N(\Omega)}} \right)^{\frac{N}{N-1}} \right)^{\frac{N-1}{N}t} dx \beta_N^{-\frac{N-1}{N}t} \|H(\nabla u)\|_{L^N(\Omega)}^t \\ &\leq \int_{\Omega} \exp \left(\left(\beta_N \frac{|u(x)|}{\|\nabla u\|_{L^N(\Omega)}} \right)^{\frac{N}{N-1}} \right) dx \beta_N^{-\frac{N-1}{N}t} \|H(\nabla u)\|_{L^N(\Omega)}^t \\ &\leq C |\Omega| \beta_N^{-\frac{N-1}{N}t} \|H(\nabla u)\|_{L^N(\Omega)}^t. \end{aligned}$$

Put

$$D_t = \Gamma \left(\frac{N-1}{N}t + 1 \right)^{1/t} C^{1/t} |\Omega|^{1/t} \beta_N^{-\frac{N-1}{N}t} t^{-\frac{N-1}{N}}.$$

Then we have

$$\|u\|_{L^t(\Omega)} \leq D_t t^{\frac{N-1}{N}} \|H(\nabla u)\|_{L^N(\Omega)}.$$

Stirling's formula implies that

$$\left(\Gamma \left(\frac{(N-1)t}{N} + 1 \right) \right)^{\frac{1}{t}} \sim \left(\frac{N-1}{Ne} \right)^{\frac{N-1}{N}} t^{\frac{N-1}{N}}$$

as $t \rightarrow \infty$. So we have

$$\lim_{t \rightarrow \infty} D_t = \beta_N^{-\frac{N-1}{N}} \left(\frac{N-1}{Ne} \right)^{\frac{N-1}{N}} = \left(\frac{1}{N \kappa_N^{1/N}} \right) \left(\frac{N-1}{Ne} \right)^{\frac{N-1}{N}},$$

which is a desired result. \square

1 Recall that C_p is defined in (1.2). Using the above Lemma and energy
2 comparison, we get the following.

3 **Proposition 3.2.** *We have*

$$\lim_{p \rightarrow \infty} p^{N-1} C_p = \left(\frac{Ne}{N-1} \beta_N \right)^{N-1}.$$

4 where $\beta_N = N(N\kappa_N)^{\frac{1}{N-1}}$.

5 *Proof.* Lower bound $\liminf_{p \rightarrow \infty} (p+1)^{N-1} C_p \geq \left(\frac{Ne}{N-1} \beta_N \right)^{N-1}$ is a direct
6 consequence of Lemma 3.1 and the fact

$$(3.1) \quad C_p = \frac{\|H(\nabla u_p)\|_{L^N(\Omega)}^N}{\|u_p\|_{L^{p+1}(\Omega)}^N}$$

7 for least energy solutions u_p .

8 Therefore we must prove only the upper bound. We will do this by
9 constructing a suitable test function for the value C_p .

10 We may assume that $0 \in \Omega$ and $\mathcal{W}_L \subset \Omega$ where $\mathcal{W}_L = \{x \in \mathbb{R}^N : H^0(x) < L\}$. For $0 < l < L$, consider the Finsler Moser function

$$m_l(x) = \frac{1}{(N\kappa_N)^{1/N}} \begin{cases} \left(\log \frac{L}{l} \right)^{\frac{N-1}{N}}, & 0 \leq H^0(x) \leq l, \\ \frac{\log \frac{L}{H^0(x)}}{\left(\log \frac{L}{l} \right)^{\frac{1}{N}}}, & l \leq H^0(x) \leq L, \\ 0, & L \leq H^0(x). \end{cases}$$

12 We check that the Moser function $m_l \in W_0^{1,N}(\Omega)$ and $\|H(\nabla m_l)\|_{L^N(\Omega)} =$
13 1. Also it is easily checked that

$$\left(\int_{\Omega} m_l^{p+1} dx \right)^{\frac{1}{p+1}} \geq \left(\int_{\mathcal{W}_l} m_l^{p+1} dx \right)^{\frac{1}{p+1}} \geq \frac{1}{(N\kappa_N)^{1/N}} \left(\log \frac{L}{l} \right)^{\frac{N-1}{N}} (l^N \kappa_N)^{\frac{1}{p+1}}.$$

14 Choosing $l = L \exp \left(-\left(\frac{N-1}{N^2} \right) (p+1) \right)$, we have

$$\|m_l\|_{L^{p+1}(\Omega)} \geq \frac{1}{(N\kappa_N)^{1/N}} \left(\frac{N-1}{N^2} \right)^{\frac{N-1}{N}} e^{-\frac{N-1}{N}} (p+1)^{\frac{N-1}{N}} (L^N \kappa_N)^{\frac{1}{p+1}}.$$

15 and

$$C_p \leq \frac{\|H(\nabla m_l)\|_{L^N(\Omega)}^N}{\|m_l\|_{L^{p+1}(\Omega)}^N} \leq N\kappa_N \left(\frac{N^2 e}{N-1} \right)^{N-1} (p+1)^{-(N-1)} (L^N \kappa_N)^{-\frac{N}{p+1}},$$

16 which implies $\limsup_{p \rightarrow \infty} (p+1)^{N-1} C_p \leq \left(\frac{Ne}{N-1} \beta_N \right)^{N-1}$. \square

1 Since

$$\int_{\Omega} H(\nabla u_p)^N dx = \int_{\Omega} u_p^{p+1} dx$$

2 and (3.1), we have the following lemma.

Lemma 3.3.

$$\lim_{p \rightarrow \infty} p^{N-1} \int_{\Omega} H(\nabla u_p)^N dx = \lim_{p \rightarrow \infty} p^{N-1} \int_{\Omega} u_p^{p+1} dx = \left(\frac{Ne}{N-1} \beta_N \right)^{N-1}.$$

3 4. PROOF OF THEOREM 1.1

4 To obtain a lower bound for $\|u_p\|_{L^\infty(\Omega)}$, define the first eigenvalue of
5 the Finsler N -Laplacian Q_N :

$$\lambda_1(\Omega) = \inf \left\{ \int_{\Omega} H(\nabla u)^N dx : u \in W_0^{1,N}(\Omega), \int_{\Omega} |u|^N dx = 1 \right\}.$$

6 It is known that $0 < \lambda_1(\Omega) < \infty$ and

$$\int_{\Omega} u_p^{p+1} dx = \int_{\Omega} H(\nabla u_p)^N dx \geq \lambda_1(\Omega) \int_{\Omega} u_p^N dx.$$

7 Thus

$$\int_{\Omega} (u_p^{p+1} - \lambda_1(\Omega) u_p^N) dx \geq 0,$$

8 which implies

$$(4.1) \quad \|u_p\|_{L^\infty(\Omega)}^{p+1-N} \geq \lambda_1(\Omega).$$

To obtain a uniform upper bound of $\|u_p\|_{L^\infty(\Omega)}$, we use an argument with the coarea formula and the Finsler isoperimetric inequality in \mathbb{R}^N . Set

$$\begin{aligned} \gamma_p &= \max_{x \in \Omega} u_p(x), \\ \Omega_t &= \{x \in \Omega : u_p(x) > t\}, \\ \mathcal{A} &= \{x \in \Omega : u_p(x) > \frac{\gamma_p}{2}\}. \end{aligned}$$

9 By Lemma 3.1 with $t = \frac{Np}{N-1}$ and by Lemma 3.3, we have

$$\left(\int_{\Omega} u_p^{\frac{Np}{N-1}} dx \right)^{\frac{N-1}{Np}} \leq D^{\frac{Np}{N-1}} \left(\frac{Np}{N-1} \right)^{\frac{N-1}{N}} \|H(\nabla u_p)\|_{L^N(\Omega)} \leq M$$

10 where M is independent of p if p large. From this and Chebyshev's
11 inequality, we have

$$(4.2) \quad \left(\frac{\gamma_p}{2} \right)^{\frac{Np}{N-1}} |\mathcal{A}| \leq M^{\frac{Np}{N-1}}.$$

1 On the other hand, by approximating the constant 1 on Ω_t by C_0^∞ -
 2 functions, we have

$$-\int_{\Omega_t} \operatorname{div} (H(\nabla u_p)^{N-1}(\nabla_\xi H)(\nabla u_p)) dx = \int_{\Omega_t} u_p^p dx.$$

Thus integration by parts leads to

$$\begin{aligned} \int_{\Omega_t} u_p^p dx &= - \int_{\partial\Omega_t} H(\nabla u_p)^{N-1}(\nabla_\xi H)(\nabla u_p) \cdot \nu ds \\ (4.3) \quad &= \int_{\partial\Omega_t} \frac{H(\nabla u_p)^{N-1}(\nabla_\xi H)(\nabla u_p) \cdot \nabla u_p}{|\nabla u_p|} ds \\ &= \int_{\partial\Omega_t} \frac{H(\nabla u_p)^N}{|\nabla u_p|} ds, \end{aligned}$$

3 since the outer unit normal ν to $\partial\Omega_t$ is $\nu = -\frac{\nabla u_p}{|\nabla u_p|}$. Here we used
 4 Proposition 2.1 (3) in the last equality. Coarea formula implies

$$|\Omega_t| = \int_{\Omega_t} 1 dx = \int_t^\infty \int_{\{u_p=s\}} \frac{ds}{|\nabla u_p|}.$$

5 Thus

$$(4.4) \quad -\frac{d}{dt}|\Omega_t| = \int_{\partial\Omega_t} \frac{ds}{|\nabla u_p|}.$$

By (4.3), (4.4), and the Schwartz inequality, we have

$$\begin{aligned} \left(-\frac{d}{dt}|\Omega_t|\right)^{N-1} \int_{\Omega_t} u_p^p dx &= \left(\int_{\partial\Omega_t} \frac{1}{|\nabla u_p|} ds\right)^{N-1} \left(\int_{\partial\Omega_t} \frac{H^N(\nabla u_p)}{|\nabla u_p|} ds\right) \\ (4.5) \quad &\geq \left(\int_{\partial\Omega_t} \frac{H(\nabla u_p)}{|\nabla u_p|} ds\right)^N \\ &= \left(\int_{\partial\Omega_t} H(\nu) ds\right)^N \\ &= P_H(\Omega_t, \mathbb{R}^N)^N \geq N^N \kappa_N |\Omega_t|^{N-1}. \end{aligned}$$

6 In the last inequality of (4.5), we used the Finsler isoperimetric in-
 7 equality in \mathbb{R}^N [3], [27], [18]:

$$(4.6) \quad P_H(E, \mathbb{R}^N) \geq N \kappa_N^{\frac{1}{N}} |E|^{\frac{N-1}{N}}$$

8 for any set of finite perimeter $E \subset \mathbb{R}^N$ with respect to H .

9 Now, define $r(t) > 0$ such that

$$|\Omega_t| = \kappa_N r^N(t).$$

1 Then

$$\frac{d}{dt}|\Omega_t| = N\kappa_N r^{N-1}(t)r'(t).$$

Note that $r'(t) < 0$. Putting this in (4.5), we have

$$\begin{aligned} \left(-N\kappa_N r^{N-1}(t)\frac{dr}{dt}(t)\right)^{N-1} \int_{\Omega_t} u_p^p dx &\geq N^N \kappa_N |\Omega_t|^{N-1}, \\ \left(-\frac{dr}{dt}\right)^{N-1} \int_{\Omega_t} u_p^p dx &\geq (N\kappa_N) r^{N-1}, \\ -\frac{dt}{dr} &\leq \left(\int_{\Omega_t} u_p^p dx\right)^{\frac{1}{N-1}} (N\kappa_N)^{-\frac{1}{N-1}} r^{-1} \\ &\leq C r^{-1} \gamma_p^{\frac{p}{N-1}} |\Omega_t|^{\frac{1}{N-1}} = C \gamma_p^{\frac{p}{N-1}} r^{\frac{1}{N-1}}, \end{aligned}$$

2 where C is a constant dependent only on N and varies from line to
3 line. Integrating the last inequality from $r = 0$ to $r = r_0$, we have

$$t(0) - t(r_0) \leq C \gamma_p^{\frac{p}{N-1}} r_0^{\frac{N}{N-1}}.$$

Choose r_0 such that $t(r_0) = \frac{\gamma_p}{2}$. Then the above inequality implies

$$\gamma_p \leq C \gamma_p^{\frac{p}{N-1}} r_0^{\frac{N}{N-1}}, \quad \text{i.e.,} \quad \gamma_p \leq C \gamma_p^{\frac{p}{N-1}} |\mathcal{A}|^{\frac{1}{N-1}}.$$

Combining this with (4.2), we have

$$\begin{aligned} \gamma_p &\leq C \gamma_p^{\frac{p}{N-1}} \left(\frac{M^{\frac{Np}{N-1}}}{\left(\frac{\gamma_p}{2}\right)^{\frac{Np}{N-1}}}\right)^{\frac{1}{N-1}} = C \gamma_p^{-\frac{p}{(N-1)^2}} M^{\frac{Np}{(N-1)^2}}, \\ \gamma_p^{1+\frac{p}{(N-1)^2}} &\leq C M^{\frac{Np}{(N-1)^2}}, \\ \gamma_p &\leq C^{\frac{(N-1)^2}{(N-1)^2+p}} M^{\frac{Np}{(N-1)^2+p}}. \end{aligned}$$

4 From this, we conclude that there exists $C > 0$ (independent of p) such
5 that $\gamma_p \leq C$ for p large.

6 The latter half part of Theorem 1.1 is already proven in Lemma 3.3.

7 Thus we have completed the proof of Theorem 1.1. \square

8 From Theorem 1.1, we have the following consequence.

9 **Corollary 4.1.** *There exist $C, C' > 0$ independent of p large such that*

$$C \leq p^{N-1} \int_{\Omega} u_p^p dx \leq C'$$

10 *holds true.*

1 *Proof.* By Theorem 1.1, we have

$$\frac{1}{C_2} p^{N-1} \int_{\Omega} u_p^{p+1} dx \leq \frac{\|u_p\|_{L^\infty(\Omega)}}{C_2} p^{N-1} \int_{\Omega} u_p^p dx \leq p^{N-1} \int_{\Omega} u_p^p dx$$

2 where C_2 is as in Theorem 1.1. The left-hand side of the above inequality is bounded from below by a positive constant by Lemma 3.3. On
3 the other hand, Hölder's inequality implies

$$p^{N-1} \int_{\Omega} u_p^p dx \leq \left(p^{N-1} \int_{\Omega} u_p^{p+1} dx \right)^{\frac{p}{p+1}} p^{\frac{1}{p+1}} |\Omega|^{\frac{1}{p+1}}$$

5 and the right-hand side of the above inequality is bounded from above
6 by Lemma 3.3. This proves the conclusion. \square

7 5. PROOF OF THEOREM 1.2

8 In this section, we prove Theorem 1.2. Since $\limsup_{p \rightarrow \infty} \|u_p\|_{L^\infty(\Omega)} \geq$
9 1 immediately follows from (4.1) (this is true for any solution sequence, not necessary least energy solutions), we just need to prove
10 $\limsup_{p \rightarrow \infty} \|u_p\|_{L^\infty(\Omega)} \leq e^{\frac{N-1}{N}}$. For this purpose, we follow the argument by Adimurthi and Grossi [1].

12 Let $x_p \in \Omega$ be a point so that the least energy solution to (1.1) takes
13 its maximum: $u_p(x_p) = \|u_p\|_{L^\infty(\Omega)}$. As in [1], We make a change of
14 variable

$$(5.1) \quad z_p(x) = \frac{p}{u_p(x_p)} (u_p(\varepsilon_p x + x_p) - u_p(x_p)), \quad x \in \Omega_p = \frac{\Omega - x_p}{\varepsilon_p},$$

16 where $\varepsilon_p > 0$ is defined so that

$$(5.2) \quad \varepsilon_p^N p^{N-1} u_p(x_p)^{p+1-N} \equiv 1.$$

By Theorem 1.1, we see $\varepsilon_p \rightarrow 0$ as $p \rightarrow \infty$. Since u_p is a weak solution to (1.1), z_p is a weak solution to

$$(5.3) \quad \begin{cases} -Q_N z_p = \left(1 + \frac{z_p}{p}\right)^p & \text{in } \Omega_p, \\ z_p|_{\partial\Omega_p} = -p, \\ \max_{x \in \bar{\Omega}_n} z_n(x) = z_n(0) = 0, \\ -p < z_p \leq 0 & \text{in } \Omega_p \end{cases}$$

by Lemma 2.7. We want to pass to the limit as $p \rightarrow \infty$ in (5.3). For this purpose, take any ball $B_R(0) \subset \Omega_p$ centered at the origin and radius R . Consider

$$\begin{cases} -Q_N w_p = \left(1 + \frac{z_p}{p}\right)^p & \text{in } B_R(0), \\ w_p|_{\partial B_R(0)} = 0. \end{cases}$$

1 Comparison principle for $-Q_N$ (see for example, [33] Theorem 3.1) and
 2 Serrin's elliptic estimate Theorem 2.4 yield that $0 \leq w_p \leq C$ on $B_R(0)$
 3 where C is a constant independent of p . Set $\psi_p(x) = w_p(x) - z_p(x)$, $x \in$
 4 $B_R(0)$. Then ψ_p is a nonnegative in $B_R(0)$ and $\psi_p(0) = w_p(0) - z_p(0) =$
 5 $w_p(0) \leq C$ uniformly in p . Moreover, we have

$$0 = -(Q_N w_p - Q_N z_p) = -\tilde{Q}_N(w_p - z_p) = -\tilde{Q}_N \psi_p$$

where

$$\begin{aligned} & \tilde{Q}_N(w_p - z_p) \\ &= \sum_{i,j=1}^N \frac{\partial}{\partial x_i} \left[\int_0^1 \frac{1}{N} \frac{\partial^2 H^N}{\partial \xi_i \partial \xi_j} (t \nabla w_p + (1-t) \nabla z_p) dt \frac{\partial}{\partial x_j} (w_p(x) - z_p(x)) \right]. \end{aligned}$$

6 Thanks to the assumption that $\text{Hess} H^N(\xi)$ is positive definite, \tilde{Q}_N is
 7 a quasilinear elliptic differential operator. Thus we can apply Serrin's
 8 Harnack inequality (Theorem 2.4 (3)) to ψ_p , which implies that there
 9 exists $C = C(R, r) > 0$ for any $0 < r < R$ such that

$$\sup_{B_r(0)} \psi_p(x) \leq C \left(1 + \inf_{x \in B_r(0)} \psi_p(x) \right) \leq C(1 + \psi_p(0)) = C(1 + w_p(0)) \leq C.$$

10 Thus we have

$$0 \geq z_p(x) = w_p(x) - \psi_p(x) \geq -C$$

11 for $x \in B_r(0)$. Since $0 < r < R$ is arbitrary, we have $\{|z_p|\} \subset$
 12 $L_{loc}^\infty(B_R(0))$ is uniformly bounded in p . Again Serrin's regularity es-
 13 timate implies that $\{z_p\}$ is bounded in $C_{loc}^{1,\alpha}(B_R(0))$ for any $R > 0$
 14 uniformly in p .

15 Now, we consider two cases:

16 **Case (i):** $\frac{\text{dist}(x_p, \partial \Omega_p)}{\varepsilon_p} \rightarrow +\infty$

17 **Case (ii):** $\frac{\text{dist}(x_p, \partial \Omega_p)}{\varepsilon_p}$ is bounded and

$$\Omega_p \rightarrow \mathbb{R}_+^N(s_0) = \{x = (x', x_N) \in \mathbb{R}^N : x_N > s_0\} \quad (p \rightarrow \infty)$$

18 for some s_0 .

19 In the case (i), note that $\Omega_p \rightarrow \mathbb{R}^N$ as $p \rightarrow \infty$. Hence by the Ascoli-
 20 Arzelá theorem, we know that (up to a subsequence), $\{z_p\}$ converges
 21 to some function $z \in C^1(\mathbb{R}^N)$ and z satisfies

$$-Q_N z = e^z \quad \text{in } \mathbb{R}^N.$$

22 Now we claim that $\int_{\mathbb{R}^N} e^z dx < +\infty$. In fact, since $z_p \rightarrow z$ in
 23 $C_{loc}^1(\mathbb{R}^N)$, we obtain

$$1_{\Omega_p}(x) \left(1 + \frac{z_p(x)}{p} \right)^p \rightarrow e^{z(x)}$$

pointwisely for $x \in \mathbb{R}^N$, where 1_{Ω_p} is the characteristic function of Ω_p . By using Fatou's lemma and Hölder's inequality, we deduce

$$\begin{aligned} \int_{\mathbb{R}^N} e^z dx &\leq \liminf_{p \rightarrow \infty} \int_{\Omega_p} \left(1 + \frac{z_p(x)}{p}\right)^p dx \\ &\leq \lim_{p \rightarrow \infty} \frac{p^{N-1}}{(u_p(x_p))^{N-1}} \int_{\Omega} (u_p(y))^p dy \\ &\leq \lim_{p \rightarrow \infty} \frac{p^{N-1}}{(u_p(x_p))^{N-1}} \left(\int_{\Omega} (u_p(y))^{p+1} dy \right)^{p/(p+1)} |\Omega|^{1/(p+1)} \\ &\leq C < \infty \end{aligned}$$

- 1 where we have used the facts that $\int_{\Omega} u_p^{p+1} dy = \frac{O(1)}{p^{N-1}}$ by Lemma 3.3
2 and $u_p(x_p) \geq C_1 > 0$ by Theorem 1.1. Hence, we check that the limit
3 function satisfies

$$(5.4) \quad \begin{cases} -Q_N z = e^z & \text{in } \mathbb{R}^N, \\ z \leq 0, & \text{in } \mathbb{R}^N, \\ \int_{\mathbb{R}^N} e^z dx < \infty. \end{cases}$$

- 4 In the case (ii), almost the same proof works, and we see that the
5 limit function z is a solution of

$$(5.5) \quad \begin{cases} -Q_N z = e^z & \text{in } \mathbb{R}_+^N(s_0), \\ z \leq 0, & \text{in } \mathbb{R}_+^N(s_0), \\ z = -\infty, & \text{on } \partial \mathbb{R}_+^N(s_0), \\ \int_{\mathbb{R}_+^N(s_0)} e^z dx < \infty. \end{cases}$$

- 6 Now we prove the following lemma. The case $N = 2$ was treated by
7 Ding (see [7]) when $H(\xi) = |\xi|$, and by Wang and Xia [32] for general
8 $H(\xi)$.

- 9 **Lemma 5.1.** *If z is a C^1 weak solution of (5.4), then we have*

$$\int_{\mathbb{R}^N} e^z dx \geq \left(\frac{N}{N-1} \right)^{N-1} N^N \kappa_N.$$

- 10 *If z is a C^1 weak solution of (5.5), then we have*

$$\int_{\mathbb{R}_+^N(s_0)} e^z dx \geq \left(\frac{N}{N-1} \right)^{N-1} N^N \kappa_N.$$

- 11 *Proof.* As in the proof of Theorem 1.1, we use a level set argument.
12 First, we assume z is a solution of (5.4). Put

$$\Omega_t = \{x \in \mathbb{R}^N : z(x) > t\}, \quad \mu(t) = |\Omega_t|.$$

Integration by parts on Ω_t leads to

$$\begin{aligned} \int_{\Omega_t} e^z dx &= - \int_{\Omega_t} Q_N z dx = \int_{\partial\Omega_t} H^{N-1}(\nabla z)(\nabla_\xi H)(\nabla z) \cdot \frac{\nabla z}{|\nabla z|} ds_x \\ &= \int_{\partial\Omega_t} \frac{H^N(\nabla z)}{|\nabla z|} ds_x. \end{aligned}$$

By the Finsler isoperimetric inequality (4.6) and Hölder's inequality, we see

$$\begin{aligned} N\kappa_N^{1/N} |\Omega_t|^{\frac{N-1}{N}} &\leq P_H(\Omega_t, \mathbb{R}^N) = \int_{\partial\Omega_t} \frac{H(\nabla z)}{|\nabla z|} ds_x \\ &\leq \left(\int_{\partial\Omega_t} \frac{H^N(\nabla z)}{|\nabla z|} ds_x \right)^{\frac{1}{N}} \left(\int_{\partial\Omega_t} \frac{ds_x}{|\nabla z|} \right)^{\frac{N-1}{N}} \\ &= \left(\int_{\Omega_t} e^z dx \right)^{\frac{1}{N}} (-\mu'(t))^{\frac{N-1}{N}}, \end{aligned}$$

1 here we have used coarea formula

$$\mu(t) = \int_t^\infty \int_{\{x: z(x)=s\}} \frac{ds_x}{|\nabla z|} ds.$$

2 Thus we have

$$\mu(t) \leq \left\{ \frac{1}{N\kappa_N^{1/N}} \left(\int_{\Omega_t} e^z dx \right)^{\frac{1}{N}} (-\mu'(t))^{\frac{N-1}{N}} \right\}^{\frac{N}{N-1}}.$$

Therefore, we obtain

$$\begin{aligned} \int_{\mathbb{R}^N} e^z dx &= \int_{-\infty}^{\max z} e^t \mu(t) dt \\ &\leq \left(\frac{1}{N\kappa_N^{1/N}} \right)^{\frac{N}{N-1}} \int_{-\infty}^{\max z} e^t \left(\int_{\Omega_t} e^z dx \right)^{\frac{1}{N-1}} (-\mu'(t)) dt \\ &= \left(\frac{1}{N\kappa_N^{1/N}} \right)^{\frac{N}{N-1}} \left(\frac{N-1}{N} \right) \int_{-\infty}^{\max z} \frac{d}{dt} \left(\int_{\Omega_t} e^z dx \right)^{\frac{N}{N-1}} dt \\ &= \left(\frac{1}{N\kappa_N^{1/N}} \right)^{\frac{N}{N-1}} \left(\frac{N-1}{N} \right) \left(\int_{\mathbb{R}^N} e^z dx \right)^{\frac{N}{N-1}}, \end{aligned}$$

3 which implies

$$\left(\frac{N}{N-1} \right)^{N-1} N^N \kappa_N \leq \int_{\mathbb{R}^N} e^z dx.$$

1 The proof when z is a solution to (5.5) is similar, since the boundary
 2 condition $z = -\infty$ on $\partial\mathbb{R}_+^N(s_0)$ assures that all level sets of z are
 3 confined in $\mathbb{R}_+^N(s_0)$. \square

4 By the change of variables, we have

$$(5.6) \quad p^{N-1} \int_{\Omega} u_p^{p+1}(y) dy = u_p^N(x_p) \int_{\Omega_p} \left(1 + \frac{z_p(x)}{p}\right)^{p+1} dx.$$

5 Let us take $\limsup_{p \rightarrow \infty}$ of both sides of (5.6). Then we see

$$\limsup_{n \rightarrow \infty} \text{LHS of (5.6)} = \left(\frac{Ne}{N-1}\right)^{N-1} N^N \kappa_N$$

by Lemma 3.3. On the other hand, Fatou's lemma and Lemma 5.1 implies

$$\begin{aligned} \limsup_{p \rightarrow \infty} \text{RHS of (5.6)} &\geq (\limsup_{p \rightarrow \infty} u_p(x_p))^N \times \begin{cases} \int_{\mathbb{R}^N} e^z dx & \text{when case (i)} \\ \int_{\mathbb{R}_+^N(s_0)} e^z dx & \text{when case (ii)} \end{cases} \\ &\geq (\limsup_{p \rightarrow \infty} u_p(x_p))^N \left(\frac{N}{N-1}\right)^{N-1} N^N \kappa_N. \end{aligned}$$

6 Hence, we have

$$e^{N-1} \geq (\limsup_{p \rightarrow \infty} \|u_p\|_{L^\infty(\Omega)})^N.$$

7 which implies Theorem 1.2 \square

8 6. PROOF OF THEOREM 1.3

In this section, we prove Theorem 1.3. Given any sequence p_n of p with $p_n \rightarrow \infty$, let us recall (1.3) and (1.6) for $p = p_n$, $u_n = u_{p_n}$.

$$\begin{aligned} v_n &= \frac{u_n}{\lambda_n} = \frac{u_n}{\left(\int_{\Omega} u_n^{p_n} dx\right)^{\frac{1}{N-1}}}, \quad \lambda_n = \left(\int_{\Omega} u_n^{p_n} dx\right)^{\frac{1}{N-1}}, \\ f_n(x) &= \frac{u_n^{p_n}}{\int_{\Omega} u_n^{p_n} dx}, \\ L_0 &= \limsup_{n \rightarrow \infty} \frac{p_n \left(\int_{\Omega} u_n^{p_n} dx\right)^{\frac{1}{N-1}}}{\left(\frac{N}{N-1} e^{\frac{N-1}{N}}\right)}, \quad L_1 = d_N^{-\left(\frac{1}{N-1}\right)} L_0. \end{aligned}$$

9 Then v_n is a weak solution of (1.4) for $p = p_n$. By Hölder's inequality
 10 and Theorem 1.1, we see

$$p_n^{N-1} \int_{\Omega} u_n^{p_n} dx \leq p_n^{N-1} \left(\int_{\Omega} u_n^{p_n+1} dx\right)^{\frac{p_n}{p_n+1}} |\Omega|^{\frac{1}{p_n+1}} \rightarrow \left(\frac{Ne}{N-1} \beta_N\right)^{N-1}$$

1 as $n \rightarrow \infty$. This shows that

$$L_0 \leq e^{\frac{1}{N}} \beta_N, \quad L_1 \leq e^{\frac{1}{N}} \beta_N d_N^{-\left(\frac{1}{N-1}\right)}.$$

2 First, we prove $S \neq \emptyset$ for any sequence $v_n = v_{p_n}$ of v_p with $p_n \rightarrow \infty$.
 3 Indeed, by Theorem 1.1, we have $\|u_n\|_{L^\infty(\Omega)} \geq C_1 > 0$ for any $n \in \mathbb{N}$.
 4 Let $x_n \in \Omega$ be a point such that $u_n(x_n) = \|u_n\|_{L^\infty(\Omega)}$, then

$$v_n(x_n) = \frac{u_n(x_n)}{\left(\int_\Omega u_n^{p_n} dx\right)^{\frac{1}{N-1}}} \geq \frac{C_1}{\left(\int_\Omega u_n^{p_n} dx\right)^{\frac{1}{N-1}}} = \frac{C_1}{O\left(\frac{1}{p_n}\right)} \rightarrow +\infty$$

5 by Lemma 3.3. This implies that any accumulation point of $\{x_n\}$ is
 6 contained in S and hence $S \neq \emptyset$.

7 Next, as in [6], [21], [22], we define (L, δ) -regular set and (L, δ) -
 8 irregular set of a sequence $\{u_n\}$. Since

$$f_n = \frac{u_n^{p_n}}{\int_\Omega u_n^{p_n} dx} \in L^1(\Omega), \quad f_n \geq 0, \quad \int_\Omega f_n dx = 1,$$

9 there exists a subsequence (still denoted by n) such that

$$f_n \xrightarrow{*} \mu, \quad \mu(\Omega) \leq 1$$

10 in the sense of Radon measures of Ω , where μ is a nonnegative Radon
 11 measure.

12 Given $L > 0$ and $\delta > 0$, we call a point $x_0 \in \Omega$ a (L, δ) -regular point
 13 of $\{u_n\}$ if there exists $\varphi \in C_0(\Omega)$, $0 \leq \varphi \leq 1$ with $\varphi \equiv 1$ near x_0 such
 14 that

$$\int_\Omega \varphi d\mu < \left(\frac{\beta_N}{L + 3\delta}\right)^{N-1}$$

where $\beta_N = N(N\kappa_N)^{\frac{1}{N-1}}$ is as in Theorem 2.2. We put

$$R_L(\delta) = \{x_0 \in \Omega : x_0 \text{ is a } (L, \delta)\text{-regular point}\},$$

$$\Sigma_L(\delta) = \Omega \setminus R_L(\delta).$$

15 We call a point in $\Sigma_L(\delta)$ an (L, δ) -irregular point of the sequence $\{u_n\}$.
 16 Note that (L, δ) -regular, or (L, δ) -irregular points are automatically
 17 interior points of Ω . Also note that if $x_0 \in \Sigma_L(\delta)$, then we have

$$(6.1) \quad \mu(\{x_0\}) \geq \left(\frac{\beta_N}{L + 3\delta}\right)^{N-1}.$$

18 Since

$$1 \geq \mu(\Omega) \geq \left(\frac{\beta_N}{L + 3\delta}\right)^{N-1} \# \Sigma_L(\delta)$$

19 by (6.1), we see that $\Sigma_L(\delta)$ is a finite set for any $L > 0$ and $\delta > 0$.

20 Next Lemma is the key to analyze the interior blow-up set $S \cap \Omega$.

Lemma 6.1. (smallness of μ implies boundedness) *Let x_0 be a (L_1, δ) -regular point of a sequence $\{u_n\}$ where L_1 is defined in (1.6). Then $\{v_n\}$ is bounded in $L^\infty(B_{R_0}(x_0))$ for some $R_0 > 0$.*

Proof. Key point in the proof is to get the following pointwise estimate

$$(6.2) \quad f_n(x) < \exp \left((L_1 + \delta/2) d_N^{\frac{1}{N-1}} v_n(x) \right), \quad x \in \Omega.$$

In checking (6.2), we use the elementary inequality

$$(6.3) \quad \frac{\log x}{x} \leq \frac{\log y}{y} \quad \text{for any } 0 < x \leq y \leq e.$$

Let

$$\alpha_n = \frac{\|u_n\|_{L^\infty(\Omega)}}{\left(\int_\Omega u_n^{p_n} dx\right)^{\frac{1}{p_n}}} = \frac{\|u_n\|_{L^\infty(\Omega)}}{\lambda_n^{\frac{N-1}{p_n}}},$$

and recall that $\lambda_n = O\left(\frac{1}{p_n}\right)$ by Corollary 4.1, so $\lambda_n^{\frac{N-1}{p_n}} = O\left(\frac{1}{p_n}\right)^{\frac{N-1}{p_n}} \rightarrow 1$ as $n \rightarrow \infty$. Thus we have

$$\limsup_{n \rightarrow \infty} \alpha_n = \limsup_{n \rightarrow \infty} \|u_n\|_{L^\infty(\Omega)} \leq e^{\frac{N-1}{N}}$$

by Theorem 1.2. From this, we see that for any small $\varepsilon' > 0$,

$$\frac{u_n(x)}{\lambda_n^{\frac{N-1}{p_n}}} \leq \alpha_n \leq e^{\frac{N-1}{N}} + \varepsilon' < e$$

holds for any $x \in \Omega$ and for large n . Therefore by (6.3), we have for fixed small $\varepsilon > 0$

$$\frac{\log \left(\frac{u_n(x)}{\lambda_n^{\frac{N-1}{p_n}}} \right)}{\frac{u_n(x)}{\lambda_n^{\frac{N-1}{p_n}}}} \leq \frac{\log \alpha_n}{\alpha_n} \leq \left(\frac{N-1}{N} \right) \frac{1}{e^{\frac{N-1}{N}}} + \varepsilon$$

for large n . Hence

$$\begin{aligned} \log f_n(x) &= p_n \log \frac{u_n(x)}{\lambda_n^{\frac{N-1}{p_n}}} \leq p_n \left(\frac{u_n(x)}{\lambda_n^{\frac{N-1}{p_n}}} \right) \left(\frac{N-1}{N} \frac{1}{e^{\frac{N-1}{N}}} + \varepsilon \right) \\ &= p_n \lambda_n \left(\frac{N-1}{N} e^{-\frac{N-1}{N}} + \varepsilon \right) \frac{v_n(x)}{\lambda_n^{\frac{N-1}{p_n}}} \\ &\leq \left(\frac{N}{N-1} e^{\frac{N-1}{N}} L_1 d_N^{\frac{1}{N-1}} + \varepsilon \right) \left(\frac{N-1}{N} e^{-\frac{N-1}{N}} + 2\varepsilon \right) v_n(x), \end{aligned}$$

1 here we have used $\lim_{n \rightarrow \infty} \lambda_n^{\frac{N-1}{p_n}} = 1$ and

$$p_n \lambda_n \leq \frac{N}{N-1} e^{\frac{N-1}{N}} L_1 d_N^{\frac{1}{N-1}} + \varepsilon$$

2 for large n by the definition of L_1 . Therefore

$$\log f_n(x) \leq \left((L_1 + \delta/2) d_N^{\frac{1}{N-1}} \right) v_n(x)$$

3 holds if we choose $\varepsilon > 0$ small enough. This proves the pointwise
4 estimate (6.2).

5 Next, by the use of Brezis-Merle theory for the Finsler N -Laplacian,
6 we obtain the integral estimate

$$(6.4) \quad \int_{B_{R_1/2}(x_0)} \exp \left((L_1 + \delta) d_N^{\frac{1}{N-1}} v_n(x) \right) dx \leq C$$

7 for some $R_1 > 0$ small and $C > 0$ independent of n , here x_0 is a
8 (L_1, δ) -regular point.

9 Indeed, by the definition of (L_1, δ) -regular point, we can find $R_1 > 0$
10 such that

$$\int_{B_{R_1}(x_0)} f_n dx \leq \left(\frac{\beta_N}{L_1 + 2\delta} \right)^{N-1}.$$

11 Also by Theorem 2.5 (i) and the fact that $\|f_n\|_{L^1(\Omega)} = 1$, we have

$$\int_{\Omega} \exp((\beta_N - \varepsilon) v_n(x)) dx \leq \frac{\beta_N}{\varepsilon} |\Omega|$$

12 for any $\varepsilon \in (0, \beta_N)$. From this, we obtain

$$(6.5) \quad \|v_n\|_{L^N(\Omega)} \leq C$$

13 where $C > 0$ is independent of n . Next, let ϕ_n be a weak solution of

$$-Q_N \phi_n = 0 \quad \text{in } B_{R_1}(x_0) \quad \phi_n = v_n \quad \text{on } \partial B_{R_1}(x_0).$$

14 Then by Theorem 2.5 (2) and the fact that $\|f_n\|_{L^1(B_{R_1}(x_0))}^{\frac{1}{N-1}} < \frac{\beta_N}{L_1 + 2\delta}$, we
15 have

$$(6.6) \quad \int_{\Omega} \exp \left((L_1 + \delta) d_N^{\frac{1}{N-1}} |v_n(x) - \phi_n(x)| \right) dx \leq C$$

16 if we choose $\varepsilon \in (0, \beta_N)$ sufficiently small. By the comparison prin-
17 ciple for the Finsler N -Laplacian (see [33] Theorem 3.2) and Serrin's
18 estimates Theorem 2.4 (i), we have

$$\|\phi_n\|_{L^\infty(B_{R_1/2}(x_0))} \leq \|v_n\|_{L^\infty(B_{R_1/2}(x_0))} \leq C \|v_n\|_{L^N(B_{R_1}(x_0))} \leq C$$

19 where we have used (6.5). Combining this with (6.6), we obtain the
20 desired integral estimate (6.4).

1 Comparing (6.4) and (6.2), we see that f_n is bounded uniformly in n
 2 in $L^q(B_{R_1/2}(x_0))$ where $q = \frac{L_1+\delta}{L_1+\delta/2} > 1$. Therefore, Serrin's regularity
 3 estimate Theorem 2.4 (i) again implies that

$$\|v_n\|_{L^\infty(B_{R_1/4}(x_0))} \leq C$$

4 independent of n . Taking $R_0 = R_1/4$ ends the proof of Lemma 6.1. \square

5 We know that $\Sigma_{L_1}(\delta)$ is a set of finite points, all of those are interior
 6 of Ω . From Lemma 6.1, we obtain $S \cap \Omega = \Sigma_{L_1}(\delta)$ for any $\delta > 0$ and

$$1 \geq \mu(\Omega) \geq \left(\frac{\beta_N}{L_1 + 3\delta}\right)^{N-1} \#(\Sigma_{L_1}(\delta)) = \left(\frac{\beta_N}{L_1 + 3\delta}\right)^{N-1} \#(S \cap \Omega).$$

7 Hence

$$\#(S \cap \Omega) \leq \left(\frac{L_1 + 3\delta}{\beta_N}\right)^{N-1} \leq \left(\frac{e^{\frac{1}{N}} \beta_N d_N^{-(\frac{1}{N-1})} + 3\delta}{\beta_N}\right)^{N-1}.$$

8 Taking a limit $\delta \rightarrow 0$, we have

$$\#(S \cap \Omega) \leq e^{\frac{N-1}{N}} d_N^{-1}$$

9 This proves the first part of Theorem 1.3.

10 If $x_0 \in S \cap \Omega = \Sigma_{L_1}(\delta)$, then for any $R > 0$, we have

$$(6.7) \quad \lim_{n \rightarrow \infty} \|v_n\|_{L^\infty(B_R(x_0))} = +\infty.$$

11 Indeed, if for some $R > 0$, assume there exists $C > 0$ independent of n
 12 such that $\|v_n\|_{L^\infty(B_R(x_0))} \leq C$ for all large n . Then

$$f_n = \frac{v_n^{p_n}}{\lambda_n^{N-1-p_n}} \leq C^{p_n} O\left(\frac{1}{p_n}\right)^{p_n-(N-1)} \rightarrow 0 \quad (n \rightarrow \infty)$$

13 uniformly on $B_R(x_0)$. This implies x_0 is a (L_1, δ) -regular point, which
 14 is absurd. The same kind of argument leads to that the limit measure
 15 μ is atomic and of the form

$$\mu = \sum_{i=1}^k \gamma_i \delta_{x_i}$$

16 where $S \cap \Omega = \{x_1, \dots, x_k\}$. Since $\mu(\Omega) \leq 1$, we have $\sum_{i=1}^k \gamma_i \leq 1$ and

$$\gamma_i \geq \left(\frac{\beta_N}{L_1}\right)^{N-1}$$

17 for all $i = 1, \dots, k$ by letting $\delta \rightarrow 0$ in (6.1) with $L = L_1$. This proves
 18 Theorem 1.3 (i).

1 On any compact sets in $\Omega \setminus (S \cap \Omega)$, $\{v_n\}$ is uniformly bounded. Then
 2 by Serrin's and Tolksdorf's regularity estimate, $\{v_n\}$ is also bounded in
 3 $C_{loc}^{1,\alpha}(\Omega \setminus (S \cap \Omega))$ for some $\alpha \in (0, 1)$. By Ascoli-Arzelá theorem, we have
 4 a subsequence and a function G such that $v_n \rightarrow G$ in $C_{loc}^1(\Omega \setminus (S \cap \Omega))$.
 5 That this G satisfies Theorem 1.3 (ii) is clear.

6 Finally, since $\lambda_n = O(\frac{1}{p_n})$ as $n \rightarrow \infty$ and $v_n(x) = \frac{u_n(x)}{\lambda_n}$ is uniformly
 7 bounded in $L_{loc}^\infty(\Omega \setminus (S \cap \Omega))$, we easily see that Theorem 1.3 (iii) holds.

8 Thus all the proof of Theorem 1.3 has been completed.

7. PROOF OF THEOREM 1.4

10 In this section, we prove Theorem 1.4.

Proof. Assume the contrary that $x_0 \in \partial\Omega$, where x_0 is the unique blow-up point of a sequence $v_n = v_{p_n}$ with $p_n \rightarrow +\infty$ as $n \rightarrow \infty$. For $R > 0$ small, we may use the Pohozaev identity Theorem 2.6 on $\Omega \cap B_R(x_0)$, with the aid of Theorem 2.4:

$$\begin{aligned}
 (7.1) \quad & \frac{N}{p_n + 1} \int_{\Omega \cap B_R(x_0)} u_n^{p_n+1} dx = \int_{\partial(\Omega \cap B_R(x_0))} \frac{u_n^{p_n+1}}{p_n + 1} (x - y) \cdot \nu(x) ds_x \\
 & - \frac{1}{N} \int_{\partial(\Omega \cap B_R(x_0))} H^N(\nabla u_n)(x - y) \cdot \nu(x) ds_x \\
 & + \int_{\partial(\Omega \cap B_R(x_0))} (H^{N-1}(\nabla u_n)(\nabla_\xi H)(\nabla u_n) \cdot \nu(x)) (x - y) \cdot \nu(x) ds_x.
 \end{aligned}$$

11 In order to remove the integral terms involving $\partial\Omega$, we use a trick in
 12 [25]. Define

$$\rho_n = \frac{\int_{\partial\Omega \cap B_R(x_0)} H^N(\nabla u_n)(x - x_0) \cdot \nu(x) ds_x}{\int_{\partial\Omega \cap B_R(x_0)} H^N(\nabla u_n)\nu(x_0) \cdot \nu(x) ds_x}$$

13 and put $y_n = x_0 + \rho_n \nu(x_0)$. We assume $R > 0$ so small such that
 14 $1/2 \leq \nu(x_0) \cdot \nu(x) \leq 1$ for $x \in \partial\Omega \cap B_R(x_0)$. Then we have that
 15 $\rho_n \leq 2R$. By the definition of y_n and ρ_n , we see that

$$\int_{\partial\Omega \cap B_R(x_0)} H^N(\nabla u_n)(x - y_n) \cdot \nu(x) ds_x \equiv 0$$

for all $n \in \mathbb{N}$. Also since $u_n = 0$ on $\partial\Omega$ and $u_n > 0$ in Ω , we see $\nu(x) = -\frac{\nabla u_n(x)}{|\nabla u_n(x)|}$. By using these, we see (7.1) with $y = y_n$ becomes

$$(7.2) \quad \begin{aligned} & \frac{N}{p_n + 1} \int_{\Omega \cap B_R(x_0)} u_n^{p_n+1} dx = \frac{1}{p_n + 1} \int_{\Omega \cap \partial B_R(x_0)} u_n^{p_n+1} (x - y_n) \cdot \nu(x) ds_x \\ & - \frac{1}{N} \int_{\Omega \cap \partial B_R(x_0)} H^N(\nabla u_n)(x - y_n) \cdot \nu(x) ds_x \\ & + \int_{\Omega \cap \partial B_R(x_0)} (H^{N-1}(\nabla u_n)(\nabla_\xi H)(\nabla u_n) \cdot \nu(x)) (x - y_n) \cdot \nabla u_n(x) ds_x. \end{aligned}$$

Multiplying $(\frac{1}{\lambda_n})^N$ to both sides of (7.2) and recalling $v_n = \frac{u_n}{\lambda_n}$, we have

$$(7.3) \quad \begin{aligned} & \frac{N}{p_n + 1} \left(\frac{1}{\lambda_n} \right)^N \int_{\Omega \cap B_R(x_0)} u_n^{p_n+1} dx \\ & = \frac{1}{p_n + 1} \left(\frac{1}{\lambda_n} \right)^N \int_{\Omega \cap \partial B_R(x_0)} u_n^{p_n+1} (x - y_n) \cdot \nu(x) ds_x \\ & - \frac{1}{N} \int_{\Omega \cap \partial B_R(x_0)} H^N(\nabla v_n)(x - y_n) \cdot \nu(x) ds_x \\ & + \int_{\Omega \cap \partial B_R(x_0)} (H^{N-1}(\nabla v_n)(\nabla_\xi H)(\nabla v_n) \cdot \nu(x)) (x - y_n) \cdot \nabla v_n(x) ds_x \\ & = I + II + III. \end{aligned}$$

We estimate the terms I, II, III on the right-hand side of (7.3) as follows:

$$\begin{aligned} |I| &= \frac{1}{p_n + 1} \left(\frac{1}{\lambda_n} \right)^N \left| \int_{\Omega \cap \partial B_R(x_0)} u_n^{p_n+1} (x - y_n) \cdot \nu(x) ds_x \right| \\ &\leq \frac{O(p_n^N)}{p_n^{N-1}(p_n + 1)} \|p_n^{N-1} u_n^{p_n+1}\|_{L^\infty(\Omega \cap \partial B_R(x_0))} \int_{\Omega \cap \partial B_R(x_0)} |(x - y_n) \cdot \nu(x)| ds_x \\ &= \frac{O(p_n^N)}{p_n^{N-1}(p_n + 1)} \|p_n^{N-1} u_n^{p_n+1}\|_{L^\infty(\Omega \cap \partial B_R(x_0))} O(R^{N-1}). \end{aligned}$$

- 1 We note that since $S \cap \Omega = \emptyset$ by assumption,

$$f_n = \frac{u_n^{p_n}}{\lambda_n^{N-1}} \rightarrow 0$$

- 2 uniformly on compact sets in Ω and

$$p_n^{N-1} u_n^{p_n+1}(x) \leq \|u_n\|_{L^\infty(\Omega)} p_n^{N-1} u_n^{p_n}(x) \leq C \frac{u_n^{p_n}(x)}{\lambda_n^{N-1}} \leq C f_n(x)$$

- 1 by Theorem 1.1 and the fact that $\lambda_n = O(\frac{1}{p_n})$ as $n \rightarrow \infty$. Thus we
 2 have

$$\|p_n^{N-1}u_n^{p_n+1}\|_{L^\infty(\Omega \cap \partial B_R(x_0))} \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

- 3 and thus

$$\lim_{R \rightarrow 0} \lim_{n \rightarrow \infty} |I| = 0.$$

Also, by Theorem 1.3 (ii), we have $v_n \rightarrow G$ in $C_{loc}^{1,\alpha}(\Omega \setminus (S \cap \Omega))$. Thus we have $H^N(\nabla v_n) = O(1)$ on $\Omega \cap \partial B_R(x_0)$, which implies

$$\begin{aligned} |II| &= \frac{1}{N} \left| \int_{\Omega \cap \partial B_R(x_0)} H^N(\nabla v_n)(x - y_n) \cdot \nu(x) ds_x \right| \\ &\leq O(1) \int_{\Omega \cap \partial B_R(x_0)} |(x - y_n) \cdot \nu(x)| ds_x \leq O(1)O(R^{N-1}), \\ |III| &= \left| \int_{\Omega \cap \partial B_R(x_0)} (H^{N-1}(\nabla v_n)(\nabla_\xi H)(\nabla v_n) \cdot \nu(x)) (x - y_n) \cdot \nabla v_n(x) ds_x \right| \\ &\leq O(1) \int_{\Omega \cap \partial B_R(x_0)} |(x - y_n) \cdot \nu(x)| ds_x \leq O(1)O(R^{N-1}). \end{aligned}$$

- 4 Therefore we have

$$\lim_{R \rightarrow 0} \lim_{n \rightarrow \infty} |II| = \lim_{R \rightarrow 0} \lim_{n \rightarrow \infty} |III| = 0.$$

- 5 From these, we obtain

$$(7.4) \quad \lim_{R \rightarrow 0} \lim_{n \rightarrow \infty} (\text{RHS of (7.3)}) = 0.$$

On the other hand, recall

$$\begin{aligned} z_n(x) &= \frac{p_n}{u_n(x_n)} (u_n(\varepsilon_n x + x_n) - u_n(x_n)), \\ x \in \Omega_{R,n} &= \frac{(\Omega \cap B_R(x_0)) - x_n}{\varepsilon_n}, \end{aligned}$$

where $\varepsilon_n^N p_n^{N-1} u_n(x_n)^{p_n+1-N} \equiv 1$. Then we see from Fatou's lemma, Theorem 1.1, and Lemma 5.1, that

$$\begin{aligned} &\lim_{R \rightarrow 0} \lim_{n \rightarrow \infty} \int_{\Omega \cap B_R(x_0)} p_n^{N-1} u_n^{p_n+1}(y) dy \\ &= \lim_{R \rightarrow 0} \lim_{n \rightarrow \infty} u_n(x_n)^N \int_{\Omega_{R,n}} \left(1 + \frac{z_n(x)}{p_n}\right)^{p_n+1} dx \\ &\geq C_1^N \int_U e^z dx \geq C_1^N \left(\frac{N}{N-1}\right)^{N-1} N^N \kappa_N \end{aligned}$$

- 6 where $u = \mathbb{R}^N$ or $\mathbb{R}_+^N(s_0)$ for some $s_0 > 0$ according to the cases
 7 $\frac{\text{dist}(x_n, \partial \Omega_{R,n})}{\varepsilon_n} \rightarrow +\infty$ or $\frac{\text{dist}(x_n, \partial \Omega_{R,n})}{\varepsilon_n} \rightarrow s_0$. Note that our assumption

- 1 $\#S = 1$ assures that we can choose x_n as a maximum points of u_n .
 2 From this and the fact that $\lambda_n = O(\frac{1}{p_n})$ as $n \rightarrow \infty$, we have

$$(7.5) \quad \lim_{R \rightarrow 0} \lim_{n \rightarrow \infty} (\text{LHS of (7.3)}) \geq C > 0$$

- 3 for some positive constant $C > 0$ independent of n .
 4 Clearly (7.5) contradicts to (7.4), and we conclude that $x_0 \notin \partial\Omega$. \square

5 Finally, as a corollary, we prove the following.

6 **Corollary 7.1.** *Let $R > 0$ and let $\{u_p\}$ be a sequence of least energy*
 7 *solutions to*

$$(7.6) \quad \begin{cases} -Q_N u_p = u_p^p & \text{in } \mathcal{W}_R, \\ u_p > 0 & \text{in } \mathcal{W}_R, \\ u_p = 0 & \text{on } \partial\mathcal{W}_R \end{cases}$$

where $\mathcal{W}_R = \{x \in \mathbb{R}^N : H^0(x) < R\}$. Then the blow-up set S of v_p satisfies $S \cap \mathcal{W}_R = \{0\}$, and

$$u_p \rightarrow G(\cdot, 0) \quad \text{in } C_{loc}^1(\mathcal{W}_R \setminus \{0\})$$

where G is the unique Green function on \mathcal{W}_R obtained in Theorem 2.3, and

$$f_p = \frac{u_p^p}{\int_{\mathcal{W}_R} u_p^p dx} \xrightarrow{*} \delta_0$$

- 8 in the sense of Radon measures on \mathcal{W}_R , along the full sequence.

Proof. The usual method of moving plane to prove the symmetry of solutions is not applicable in the anisotropic situation. However, we can use Theorem 4.1 in [5] under the convexity and C^1 -assumption of the map $\xi \mapsto H^N(\xi)$. (Note that the key point of the proof of Theorem 4.1 in [5] is the Pohozaev identity Theorem 4.2 in [5] for $C^1(\bar{\Omega})$ -weak solutions, which is valid by the above assumptions). Thus we assure that any positive solution u_p to (7.6) is *Finsler-radial*, that is, all level sets of u_p are homothetic to \mathcal{W}_R for any $p > 1$. Let S be the blow-up set of v_p . Then we see that $S \cap \mathcal{W}_R = \{0\}$. Indeed, if there were a point $x_0 \in S \cap \mathcal{W}_R$, then all points on the level set of u_p passing through x_0 must be blow-up points of v_p , which contradicts to the fact that $\#(S \cap \mathcal{W}_R)$ is finite. Thus by Theorem 1.3, we see

$$v_p \rightarrow G(\cdot, 0) \quad \text{in } C_{loc}^1(\mathcal{W}_R \setminus \{0\})$$

- 9 for some function G along a subsequence. The limit function must
 10 be the unique Green function constructed in Theorem 2.3, and by the
 11 uniqueness, the convergence is true for the full sequence. \square

1 Acknowledgments.

2 This work was partly supported by Osaka City University Advanced
 3 Mathematical Institute MEXT Joint Usage / Research Center on Math-
 4 ematics and Theoretical Physics JPMXP0619217849. The second au-
 5 thor (F.T.) was supported by JSPS KAKENHI Grant-in-Aid for Scien-
 6 tific Research (B), JP19H01800, and JSPS Grant-in-Aid for Scientific
 7 Research (S), JP19H05597.

8

REFERENCES

- 9 [1] Adimurthi, and M. Grossi: *Asymptotic estimates for a two-dimensional prob-*
 10 *lem with polynomial nonlinearity*, Proc. Amer. Math. Soc., **132**, no. 4, (2003),
 11 1013–1019.
- 12 [2] A. Alvino, V. Ferone, A. Mercaldo, F. Takahashi, R. Volpicelli: *Finsler Hardy-*
 13 *Kato's inequality*, J. Math. Anal. Appl., **470**, no. 1, (2019), 360–374.
- 14 [3] A. Alvino, V. Ferone, G. Trombetti, and P. L. Lions: *Convex symmetrization*
 15 *and applications*, Ann. Inst. H. Poincaré Anal. Non Linéaire, **14** (1997), no. 2,
 16 275–293.
- 17 [4] G. Bellettini, and M. Paolini: *Anisotropic motion by mean curvature in the*
 18 *context of Finsler geometry*, Hokkaido Math. J., **25** (1996), 537–566.
- 19 [5] M. Belloni, V. Ferone, and B. Kawohl: *Isoperimetric inequalities, Wulff shape*
 20 *and related questions for strongly nonlinear elliptic operators*, ZAMP. **54**
 21 (2003), 771–783.
- 22 [6] H. Brezis, and F. Merle: *Uniform estimates and blow-up behavior for solutions*
 23 *of $-\Delta u = V(x)e^u$ in two dimensions*, Comm. Partial Differential Equations,
 24 **16** (1991) 1223–1253.
- 25 [7] E. Chen, and C. Li: *Classification of solutions of some nonlinear elliptic equa-*
 26 *tions*, Duke Math. J., **63**, no.3, (1991) 615–622.
- 27 [8] A. Cianchi and P. Salani: *Overdetermined anisotropic elliptic problems*, Math.
 28 Ann., **345** (2009), 859–881.
- 29 [9] G. Ciruolo, A. Figalli and A. Roncoloni: *Symmetry results for critical*
 30 *anisotropic p -Laplacian equations in convex cones*, Geom. Funct. Anal., **30**
 31 (2020), 770–803.
- 32 [10] M. Cozzi, A. Farina and E. Valdinoci: *Gradient bounds and rigidity results for*
 33 *singular, degenerate, anisotropic partial differential equations*, Comm. Math.
 34 Phys., **331** (2014), 189–214.
- 35 [11] M. Degiovanni, A. Musesti, and M. Squassina: *On the regularity of solutions in*
 36 *the Pucci-Serrin identity*, Calc. Var. Partial Differential Equations, **18** (2003),
 37 318–334.
- 38 [12] F. Della Pietra, and G. di Blasio: *Blow-up solutions for some nonlinear elliptic*
 39 *equations involving a Finsler-Laplacian*, Publ. Mat., **61** (2017), 213–238
- 40 [13] F. Della Pietra, G. di Blasio, and N. Gavitone: *Anisotropic Hardy inequalities*,
 41 Proc. Roy. Soc. Edinburgh Sect. A, 148A, (2018), no.3, 483–498.
- 42 [14] F. Della Pietra, and N. Gavitone: *Symmetrization with respect to the*
 43 *anisotropic perimeter and applications*, Math. Ann., **363** (2015), 953–971
- 44 [15] E. DiBenedetto: *$C^{1,\alpha}$ local regularity of weak solutions of degenerate elliptic*
 45 *equations*, Nonlinear Anal., **7** (1983), 827–850

- 1 [16] L. C. Evans, and R. F. Gariepy: *Measure Theory and Fine Properties of Functions*, CRC press (1992), iv + 268 pages.
- 2
- 3 [17] V. Ferone, and B. Kawohl: *Remarks on a Finsler-Laplacian*, Proc. Amer. Math. Soc., **137** (2009), no.1, 247–253.
- 4
- 5 [18] I. Fonseca, and S. Müller: *A uniqueness proof for the Wulff theorem*, Proc. Roy. Soc. Edinburgh Sect. A., **119** (1991) 125–136.
- 6
- 7 [19] G. Lieberman. *Boundary regularity for solutions of degenerate elliptic equations*, Nonlinear Anal., **12** (1983), no.11, 1203–1219.
- 8
- 9 [20] A. Mercaldo, M. Sano, and F. Takahashi: *Finsler Hardy inequalities*, Math. Nachr., **293** (2020), no.12, 2370–2398.
- 10
- 11 [21] X. Ren, and J. Wei: *On a two-dimensional elliptic problem with large exponent in nonlinearity*, Trans. Amer. Math. Soc., **343** (1994) 749–763.
- 12
- 13 [22] X. Ren, and J. Wei: *Single-point condensation and least-energy solutions*, Proc. Amer. Math. Soc., **124** (1996) 111–120.
- 14
- 15 [23] X. Ren, and J. Wei: *Counting peaks of solutions to some quasilinear elliptic equations with large exponents*, J. Differential Equations, **117** (1995) 28–55.
- 16
- 17 [24] S. Santra: *Existence and shape of the least energy solutions of a fractional Laplacian*, Calc. Var. and Partial Differential Equations, **58** no.2, (2019) page no. 48, 25 pages.
- 18
- 19
- 20 [25] S. Santra, and J. Wei: *Asymptotic behavior of solutions of a biharmonic Dirichlet problem with large exponents*, J. Anal. Math., **115** (2011) 1–31.
- 21
- 22 [26] J. Serrin: *Local behavior of solutions of quasi-linear equations*, Acta. Math., **111** (1964) 248–302.
- 23
- 24 [27] J. Taylor: *Crystalline variational problems*, Bull. Amer. Math. Soc., **84** (1978) 568–588.
- 25
- 26 [28] F. Takahashi: *Asymptotic behavior of least energy solutions to a four-dimensional biharmonic semilinear problem*, Osaka. J. Math., **42** (2005) 633–651.
- 27
- 28
- 29 [29] F. Takahashi: *Single-point condensation phenomena for a four-dimensional biharmonic Ren-Wei problem*, Calc. Var. and Partial Differential Equations, **29** (2007) 509–520.
- 30
- 31
- 32 [30] F. Takahashi: *Asymptotic behavior of least energy solutions for a 2D nonlinear Neumann problem with large exponent*, J. Math. Anal. Appl., **411** (2014) no.1, 95–106.
- 33
- 34
- 35 [31] P. Tolksdorf: *Regularity for a more general class of quasilinear elliptic equations*, J. Differential Equations, **51** (1984) 126–150.
- 36
- 37 [32] G. Wang, and C. Xia: *Blow-up analysis of a Finsler-Liouville equation in two dimensions*, J. Differential Equations, **252** (2012), 1668–1700.
- 38
- 39 [33] R. L. Xie, and H. J. Gong: *A priori estimates and blow-up behavior for solutions of $-Q_N u = V e^u$ in bounded domain in \mathbb{R}^N* , Sci. China Math., **59** (2016), 479–492.
- 40
- 41
- 42 [34] Changliang Zhou, and Chunqin Zhou: *Moser-Trudinger inequality involving the anisotropic Dirichlet norm $(\int_{\Omega} F^N(\nabla u) dx)^{\frac{1}{N}}$ on $W_0^{1,N}(\Omega)$* , J. Funct. Anal., **276** (2019), no.9, 2901–2935.
- 43
- 44