

# Host selection and ovipositor length in eight sympatric species of sculpins that deposit their eggs into tunicates or sponges

メタデータ	言語: English 出版者: Springer 公開日: 2020-06-01 キーワード (Ja): 産卵管, カジカ科, ホヤ, カイメン, 卵寄託 キーワード (En): ovipositor, sculpins, tunicates, sponges, marine fish, ostracophils 作成者: 安房田, 智司, 佐々木, 晴香, 後藤, 智仁, 古屋, 康則, 武島, 弘彦, 山崎, 彩, 宗原, 弘幸 メールアドレス: 所属: Nigata University, Osaka City University, Nigata University, Nigata University, Gifu University, Research Institute for Humanity and Nature, Tokai University, Hokkaido University, Hokkaido University
URL	<a href="https://ocu-omu.repo.nii.ac.jp/records/2020094">https://ocu-omu.repo.nii.ac.jp/records/2020094</a>

# Host selection and ovipositor length in eight sympatric species of sculpins that deposit their eggs into tunicates or sponges

Satoshi Awata, Haruka Sasaki, Tomohito Goto, Yasunori Koya,  
Hirohiko Takeshima, Aya Yamazaki, Hiroyuki Munehara

<b>Citation</b>	Marine Biology, 166(5); 59
<b>Issue Date</b>	2019-04-06
<b>Type</b>	Journal Article
<b>Textversion</b>	Author
<b>概要</b>	<p>研究グループは、他の生物に卵を預ける海産のカジカ科魚類 8 種について産卵宿主種（ホヤやカイメン）の特定に成功し、さらにはカジカ科魚類が産卵管を宿主種の種類や大きさに応じて進化させていることを世界で初めて発見しました。</p> <p>今回の発見は、生態研究がほとんど進んでいない冷たい海域の動物の生態を知るうえで、非常に重要な成果と言えます。</p> <p>‘Host selection and ovipositor length in eight sympatric species of sculpins that deposit their eggs into tunicates or sponges’. 大阪市立大学. <a href="https://www.osaka-cu.ac.jp/ja/news/2019/190418">https://www.osaka-cu.ac.jp/ja/news/2019/190418</a>. (参照 2019-04-18)</p>
<b>Rights</b>	<p>This is a post-peer-review, pre-copyedit version of an article published in Marine Biology. The final authenticated version is available online at:</p> <p><a href="https://doi.org/10.1007/s00227-019-3506-4">https://doi.org/10.1007/s00227-019-3506-4</a>.</p> <p>See Springer Nature terms of use.</p> <p><a href="https://www.springer.com/gp/open-access/publication-policies/aam-terms-of-use">https://www.springer.com/gp/open-access/publication-policies/aam-terms-of-use</a>.</p>
<b>Electronic supplementary material</b>	<p>Electronic supplementary material is available online at:</p> <p><a href="https://doi.org/10.1007/s00227-019-3506-4">https://doi.org/10.1007/s00227-019-3506-4</a>.</p>
<b>DOI</b>	10.1007/s00227-019-3506-4

Self-Archiving by Author(s)  
Placed on: Osaka City University Repository

**Original paper**

**Host selection and ovipositor length in eight sympatric species of sculpins that deposit their eggs into tunicates or sponges**

Satoshi Awata<sup>1,2</sup> · Haruka Sasaki<sup>1</sup> · Tomohito Goto<sup>1</sup> · Yasunori Koya<sup>3</sup> · Hirohiko Takeshima<sup>4,5</sup> · Aya Yamazaki<sup>6</sup> · Hiroyuki Munehara<sup>6</sup>

<sup>1</sup>Sado Marine Biological Station, Faculty of Science, Niigata University, Sado, Japan

<sup>2</sup>Laboratory of Animal Sociology, Graduate School of Science, Osaka City University, Osaka, Japan

<sup>3</sup>Department of Biology, Faculty of Education, Gifu University, Gifu, Japan

<sup>4</sup>Research Institute for Humanity and Nature, Kyoto, Japan

<sup>5</sup>Department of Marine Biology, School of Marine Science and Technology, Tokai University, Shizuoka, Japan

<sup>6</sup>Usujiri Fisheries Station, Field Science Center for Northern Biosphere, Hokkaido University, Hakodate, Japan

Short title: Host selection and ovipositor length in sculpins

Correspondence

Satoshi Awata

Laboratory of Animal Sociology, Graduate School of Science, Osaka City University, 3-3-138 Sugimoto, Sumiyoshi, Osaka 558-8585, Japan.

Tel.: +81 6 6605 2607

e-mail: sa-awata@sci.osaka-cu.ac.jp

ORCID ID: 0000-0003-3254-7943

## Abstract

Interspecific interactions between parasites and hosts can influence the evolution of behavioural and morphological adaptations of both parasites and their hosts. There is, however, little empirical evidence available regarding the evolution of reproductive traits driven by these interactions. In this paper, we investigated host selection and ovipositor length in nine sympatric marine sculpins that oviposit into tunicates or sponges. Field and genetic studies have revealed host use for eight out of nine species of sculpins investigated here: five species of *Pseudoblennius*, two species of *Furcina* and one species of *Vellitor*. For one species studied (*V. minutus*), no egg masses could be found. Ovipositor length reflects morphology of host species utilised: six sculpin species had extremely long ovipositors allowing females to attach eggs to the deep atrium of solitary tunicates, whereas the two species that attached their eggs to the small space of atrial siphon of colonial tunicates and the spongocoel of sponges had short ovipositors. Ovipositor length varied between solitary-tunicate spawners and species with longer ovipositors selected larger tunicates. Since the ancestral form is non-parasitic, the ovipositor evolved as an adaptation to utilise sponges and tunicates as hosts. Sculpins found sympatrically may show host specificity to avoid interspecific competition for spawning niches and ovipositors may have evolved depending on the species and size of host invertebrates.

Keywords: ovipositor; sculpins; tunicates; sponges; marine fish; ostracophils.

## Introduction

Sexual selection, arising from interaction between sexes, has considerable influence on evolutionary adaptations in both sexes, for example, the morphological adaptation of reproductive traits (Andersson 1994; Birkhead and Møller 1998). However, as yet only a few studies have examined the evolution of reproductive traits in parasites through the interactions between parasites and hosts. In parasitic animals that utilise other organisms as oviposition hosts, female selection and oviposition success can strongly affect the survival of offspring and thereby parental fitness (e.g. Smith et al. 2001; Mills and Reynolds 2002, 2003; Kitamura 2005; Reichard et al. 2007), whereas hosts often incur costs due to egg deposition (e.g. Peden and Corbett 1973; Poltev and Mukhametov 2009; Reichard et al. 2006). Therefore, such interspecifically antagonistic interactions can influence the evolution of behavioural and morphological adaptations of either or both parasites and their hosts.

Some of the parasitic animals that inhabit aquatic environments preferentially deposit their eggs in live invertebrates (known as ‘ostracophils’: Balon 1975; Leung 2014). For example, bitterlings (Acheilognathinae) deposit their eggs in the gill chambers of freshwater mussels through their atrial siphon (e.g. Smith et al. 2004; Kitamura 2007), the Japanese tubesnout (*Aulichthys japonicus*) conceals their eggs in tunicates (Akagawa et al. 2008) and snailfishes (Liparidae) deposit eggs within the branchial chambers of crabs (e.g. Hunter 1969; Somerton and Donaldson 1998; Poltev and Mukhametov 2009; Gardner et al. 2016). All of these species display some degree of host specificity, and various degrees of adaptation of reproductive traits, for the deposition of their eggs inside invertebrates (reviewed in Leung 2014); for example, the extremely long ovipositors observed in both bitterlings (Smith et al. 2004; Kitamura 2007; Kitamura et al. 2012) and snailfishes (Hunter 1969). Similar adaptations of reproductive traits through interactions between parasites and hosts have been

reported for various parasitic wasps including parasitoids that target free-living insects (Leralec et al. 1996), gall-makers (Ghara et al. 2011) and figs (Elias et al. 2018). These ostracophilous fishes and parasitic wasps have shown that sexual traits can be inter- and intraspecifically variable, due to interactions between parasites and hosts. However, empirical evidence regarding these variations in reproductive traits and host selection remain scarce.

Sculpins (Teleostei: Cottidae), which belong to the Cottoidea superfamily, are a large group of fish that have 70 genera and 282 species worldwide, and primarily inhabit boreal and cold temperate regions (Nelson et al. 2016). Although evidence is scarce, several species of Cottoidea are known to deposit their eggs inside alive invertebrates (reviewed in Abe and Munehara 2009; Munehara et al. 2011). For example, the little dragon sculpin *Blepsias cirrhosus* and the bigmouth sculpin *Hemitripterus bolini* utilise sponges as their spawning host (Munehara 1991; Busby et al. 2012). It has also been reported that four sculpin species in the genus *Pseudoblennius* that inhabit temperate waters seem to be ostracophils (Uchida 1932, 1979; Shinomiya and Ikemoto 1987; Okamura and Amaoka 1997; Nishida et al. 2008; Uryu 2011). According to these descriptive studies and to underwater photographs taken by divers, these species are all copulating species in which females deposit their eggs exclusively inside tunicates or sponges using their ovipositors (Shiogaki and Dotsu 1974; Shinomiya 1985; Abe and Munehara 2009; Nishida et al. 2008; Uryu 2011). In addition, during a preliminary study of marine sculpins in the nearshore reef of Sado Island, located in the Sea of Japan, it was found that there may be nine species of potentially ostracophilous sculpins (five species of *Pseudoblennius*, two species of *Furcina* and two species of *Vellitor*). They all develop ovipositors and occur sympatrically in shallow water of the island (Awata 2015, 2017). Therefore, these sculpins may be ideal for an examination of the evolution of interspecific variation in reproductive traits that are driven by interactions between parasites and hosts. However, no studies have demonstrated host selection and ovipositor morphology in sculpins,

with the exception of one study that showed host tunicate selection by *Pseudoblennius percooides* (Nishida et al. 2008).

In this study, we investigated interspecific variation in host selection and ovipositor length in nine sympatric species of marine sculpins that deposit their eggs into tunicates or sponges on the coast of Sado Island, northern Japan. We hypothesise that species-specific host selection by sculpin females drives the evolution of ovipositor morphology. To confirm our hypothesis, we tested three predictions. Firstly, that the nine ostracophilous sculpin species would utilise different species and sizes of host tunicates or sponges to reduce interspecific competition for hosts (Reichard et al. 2007). This is because these sculpins co-occurred at the study sites. Secondly, that only ostracophilous sculpins, and not species that lay eggs on rock surfaces, would have elongated ovipositors. The morphological and molecular phylogeny of the sculpins suggests that ostracophilous sculpins have evolved from non-ostracophilous sculpins (Yabe 1985; Abe and Munehara 2009; Koya et al. 2011; Knope 2013). Therefore, the development of the ovipositor is likely related to such unique spawning behaviour. Finally, that ovipositor length would reflect the species and size of the hosts utilised.

## **Materials and methods**

### **Study sites**

Fieldwork using self-contained underwater breathing apparatus (SCUBA) was conducted on the coast of Sado Island in the Sea of Japan. The island is located about 45 km off the Japanese mainland coastal city of Niigata (Fig. 1a). The amplitude of the tides around the island is relatively low, ranging from only 0.2 to 0.3 m. Four sites (Fig. 1b) were selected for fieldwork. All of the host invertebrates and ostracophilous sculpin species were abundant at the study sites, with the exception of *V. minutus* that was only found at Tassyia (Awata 2017).

The four study sites were located on rocky shores where seaweed grows abundantly.

### **Fish study species**

There were nine species of potentially ostracophilous sculpins in the nearshore reef of Sado Island (Nakabo and Kai 2013; Awata 2015, 2017). Microhabitat preferences differ between species (Iwata 1983; Awata 2017), but all species occurred sympatrically in rocky shallow water ranging in depth from 1 to 20 m. They are copulating species and their fertilisation mode is internal gametic association (IGA: Munehara et al. 1989), where fertilisation occurs only when eggs are released into the seawater (Shiogaki and Dotsu 1974; Kimura et al. 1987, 1988; Koya et al. 2015). Eggs that are released into seawater are spherical of 1.3–2.0 mm in diameter, demersal and adhesive, forming egg masses of up to 500 eggs (e.g. Kimura et al. 1987, 1988; Nishida et al. 2008). There are nine species in three genera, *Furcina*, *Pseudoblennius* and *Vellitor*, which form a closely related and monophyletic group (Yabe 1985; Munehara et al. 2011). The reproductive ecology is similar among species, but their body size and shape are notably different: *Pseudoblennius* sp. (“Kirin-anahaze” in Japanese, see Nakabo and Kai 2013) and *Ps. percoides* are large sized (130–170 mm in standard length, SL), *Ps. cottoides*, *Ps. marmoratus* and *Ps. zonostigma* are medium sized (80–100 mm SL) and *F. ishikawae*, *F. osimae*, *V. centropomus* and *V. minutus* are small sized sculpins (40–70 mm SL; Okamura and Amaoka 1997; Munehara et al. 2011; Nakabo and Kai 2013; Awata 2017).

### **Fish collection and measurement of ovipositor length**

To examine the ovipositor length of the nine species of ostracophilous sculpins, adult females were captured using hand-nets while SCUBA diving during the spawning season, from late November to late April in 2014, 2015 and 2016, in depth of 2 to 15 m at all four study sites



(Fig. 1b). The fish were transported in aerated coolers to the laboratory at the Sado Marine Biological Station (SMBS), Niigata University, Sado, Japan. Species identification of sculpins was based on that of Nakabo and Kai (2013). Fish were anaesthetised with a solution of MS222 or 2-phenoxyethanol buffered with seawater, and their SL (to 1 mm) was measured using a scale. The ovipositor was induced to its maximum extrusion by gently pressing the belly of the fish between ventral fins and anus. The ovipositor was photographed alongside a scale, and ovipositor length (OL) was measured to the nearest 1 mm from the apex to the ventral edge of the ovipositor using image analysis software ImageJ 1.50i. A part of the right pectoral fin was clipped from each individual and preserved in 99 % ethanol for genetic analysis. Of the 124 females analysed, 40 were released after the operation and 84 were used for other studies after sacrifice under deep anaesthesia (taxonomy and sperm analyses: Awata 2015, 2017; histology: Koya et al. 2015).

To determine whether only ostracophilous sculpins have evolved elongated ovipositors, we compared the OL of ostracophilous sculpins and sculpin species that oviposit on the rock surface. We captured 13 females of four non-ostracophilous sculpin species (*Alcichthys alcicornis*, *Bero elegans*, *Icelinus pietschi* and *Radulinopsis derjavini*) using SCUBA at Usujiri (Fig. 1a), southern Hokkaido, Japan and at Kotoura, Sado Island, Japan, and measured their SL and OL.

### **Collection of egg masses deposited in host invertebrates**

Egg masses deposited in the spongocoels of sponges and in the common atrial siphons of colonial tunicates were collected in April and May 2012, from March to May 2013 and from February to April 2014 at Kotoura and Kitakoura (Fig. 1b). Egg masses were externally visible, and therefore only whole colonies of sponges and colonial tunicates that contained egg masses were collected using scrapers. The colonies that contained eggs were distributed

across rock wall habitats at depths of 1–12 m. The sea water temperature was 9–14 °C.

Because the egg masses deposited in the atria of solitary tunicates were rarely visible from outside, we haphazardly collected solitary tunicates ( $n = 1212$ ) in December 2011, December 2012 and from December 2013 to May 2014 (approximately once a month) at Kitakoura, Mushizaki and Tassya (Fig. 1b). Of these, 1129 (93 %) tunicates were captured from the wall of Kitakoura Port at a depth of 0.5–4.0 m (Fig. 1c), where the wall was covered with in excess of a hundred thousand solitary tunicates. The remaining samples were collected at 8–18 m depth of off Kitakoura Port, at 8–30 m depth of Mushizaki and at 2–5 m of Tassya. The sea water temperature ranged from 9 to 16 °C (the coldest being in February).

All tunicates, sponges and egg masses were transferred to the laboratory in aerated coolers. Sponges and colonial tunicates were cut from the osculum and common atrial siphon, respectively, to the spongocoel and common atrium. The length between the open top and the top of the egg mass was measured using a scale (defined as “egg mass position”). Solitary tunicates were photographed with a scale and their long diameter (LD) and short diameter (SD) were later measured using ImageJ 1.50i software. After being photographed, tunicates were carefully cut in half vertically using a kitchen knife, in order to avoid damaging the egg masses. When egg masses were found inside tunicates, opened tunicates were photographed again with a scale to measure the length from entrance of the atrial siphon to the nearest edge of the egg masses (also defined as “egg mass position”). It is natural to expect that utilisation of larger tunicates will require longer ovipositors, and thus egg mass position will be deeper when larger tunicates are used as hosts. The LD was used as an indicator of the body size of tunicates, given that their LD and SD are strongly correlated (Pearson’s  $r = 0.92$ ,  $p < 0.0001$ ,  $n = 1212$ ).

Each egg mass removed from the host was counted and transferred to a box-shaped net cage (12.5 × 13.0 × 8.0 cm, mesh size: 0.5 mm, s-5330, Sudo & Company, Inc., Nagoya,

Japan) that was fixed to the surface of a 90 × 45 × 45 cm tank at the SMBS. Each egg mass was separately incubated for 1–20 days in the cage at 15 °C. Hatched larvae or late stage eggs (just before hatching) were preserved in 99 % ethanol after being killed by an overdose of anaesthesia.

Note that egg masses of the Japanese tubenout *Aulichthys japonicas* were also collected during this study, but the details of this will be published in a later paper.

### **Genetic identification of parent species of egg masses**

To infer the parental species of the egg masses, 1–3 larvae per egg mass and fin-clip samples of nine potentially ostracophilous sculpins, with the addition of *Jordania zonope* as an outgroup, were chosen for use in genetic analyses. Fin-clip samples were obtained from 2011–2017 (Table S1, Supplementary Information). Whole genomic DNA was extracted from 148 adults and from 223 larvae (including late stage eggs) of 120 egg masses, using a Genra Puregene Tissue Kit (Qiagen, Hilden, Germany). Samples were PCR amplified for the mitochondrial cytochrome *b* gene (*cytb*): GLUDG-L: 5'-YGA CTTGAARAACCA YCGTTG-3' and CB3-H: 5'-GGCAAATAGGAARTATCATTC-3' (Palumbi 1996; Knope 2013). PCR amplifications were performed at a 15 µL volume using 7.5 µL of 2 × Go Taq Green Master Mix (Promega, WI, USA), 0.6 µL (5 µM) of each primer, 1.5 µL ca. 50 ng/µL of template DNA and 4.8 µL of nuclease free water. PCR thermal cycling was performed using the following protocol for *cytb*: 2 min of initial denaturation at 95 °C, followed by 35 cycles of 94 °C for 30 sec, 50 °C for 30 sec and 72 °C for 1 min. PCR amplifications were successful except for some *Ps. cottoides* and *Ps. zonostigma* and many of the egg masses, for which sequences corresponding to the primers were permutated. Therefore, these samples were PCR amplified using another primer set for *cytb*: H15915: 5'-CAACGATCTCCGGTTT-3' and L14724: 5'-GTGACTTGAAAAACCA-3' (Schmidt and Gold 1993). The 800 bp sequences

amplified by this primer set overlapped with those amplified by the former primer set. PCR products were electrophoresed on 1.5 % agarose gel stained with GelRed™ (Biotium, Inc., CA, USA). Positive amplicons (evidenced by a clear single band of the correct size) were enzymatically cleaned with illustra™ ExoStar™ (GE Healthcare UK Limited, Little Chalfont, Buckinghamshire, UK). Double stranded PCR products were sequenced directly with the same primers used for the PCR amplifications at Macrogen Japan Corp. (Kyoto, Japan). All sequences were aligned using the ClustalW algorithm implemented in MEGA ver. 7.0.14. (Kumar et al. 2016).

The phylogenetic relationships of nine potential parental species ( $n = 146$ ) and *Jordania zonope* ( $n = 2$ , as the root for the tree) were reconstructed using 800 bp *cytb* sequences with MEGA ver. 7.0.14, and the Maximum Likelihood method was employed using the Tamura Nei model settings (Tamura and Nei 1993) and 1000 bootstrap replicates. After assessing the usefulness of the sequences for species identification (Fig. S1, Supplementary Information; all these sequences available on GenBank, Table S1), the parental species of egg masses were estimated using the reconstructed phylogeny of the parental species and egg masses. Of 223 eggs analysed, 205 were identified using 800 bp *cytb* sequences, and 18 were identified using 628 bp, due to the low quality of the sequencing data.

### **Data analyses**

Statistical analyses were performed using SPSS ver. 23.0 (IBM Corp., Armonk, NY, USA). Data were analysed using parametric statistics throughout. All reported  $p$ -values were two tailed, and results were considered statistically significant at  $p$ -values of  $< 0.05$ . As we did not obtain all data from all sponges and tunicates, sample size varied between analyses. To determine whether the relative OL differed between ostracophilous and non-ostracophilous sculpins, we used an analysis of covariance (ANCOVA) with log-transformed OL as the

dependent variable and deposition site (sponge, colonial tunicate, solitary tunicate and rock surface) as the independent variable and log-transformed SL as the covariate, including their interaction (full model). If the interaction term (deposition site  $\times$  log SL) was not statistically significant, then a reduced model was constructed by reducing the interaction term.

## Results

### Egg masses deposited in sponges and colonial tunicates

Nine egg masses were obtained from sponges (Demospongiae) of nine colonies. Since each colony was small and formed mostly only one spongocoel, multiple egg masses were not found in sponges sampled (Fig. 2a). In contrast, the colony size and shape of colonial tunicates (Didemnidae) were variable, ranging from 20 to 100 mm in maximum width, each of which had 1–38 exhalant siphons and contained 1–5 sculpin egg masses at similar or different developmental stages (Fig. 2b, c). In total, 34 egg masses were obtained from the colonial tunicates of 22 colonies.

### Egg masses deposited in solitary tunicates

Of 1212 solitary tunicates collected, seven species were identified. The most commonly occurring was *Halocynthia ritteri* (55.9 %,  $n = 678$ ), followed by *Polycarpa cryptocarpa kroboja* (25.3 %,  $n = 307$ ), *Cnemidocarpa irene* (11.6 %,  $n = 141$ ), *Pyura sacciformis* (3.5 %,  $n = 42$ ), *Microcosmus hartmeyeri* (2.0 %,  $n = 24$ ), *H. roretzi* (1.6 %,  $n = 19$ ) and *Herdmania japonica* (0.1 %,  $n = 1$ ). Although a considerable number of tunicates were collected, sculpins only utilised three species (Fig. 2d–h). The two largest and the smallest species of tunicates were not used by the sculpins (Fig. 3). The second most abundant tunicate *Po. c. kroboja* was never used by sculpins, although notably overlapped in size with the most abundant and most

frequently used tunicate *H. ritteri* (Fig. 3).

Of the 89 sculpin egg masses collected, > 90 % were found in *H. ritteri* and the remainder were found in *Py. sacciformis* and *H. roretzi* (Fig. 2f–h; Fig. 3). Although the abundance of *H. ritteri* was more than 16 times greater than that of *Py. sacciformis* at the study sites, the proportion of parasitized tunicates was equal ( $n_{with\ eggs}/n_{without\ eggs} = 75/593$  for *H. ritteri*;  $6/30$  for *Py. sacciformis*: Fisher's exact test,  $p = 0.29$ ). Note that sculpin eggs were only found in the atria of tunicates, indicating that the ovipositors were inserted through the atrial siphon of the tunicates.

### Parental species of egg mass

The reconstructed phylogenetic relationships of the potentially parental species and 120 egg masses showed that sponges were utilised by *Ps. marmoratus* ( $n = 4$  egg masses, Fig. 2a) and *V. centropomus* ( $n = 4$ ) as spawning hosts (details in Table S2). All 26 egg masses deposited in colonial tunicates were assigned to *V. centropomus* (Fig. 2b). Solitary tunicates were used by six sculpin species (Table 1; Fig. 2d–h; details in Table S3): egg masses of *Ps. cottoides* were the most frequent ( $n = 52$  egg masses), followed by those of *Ps. zonostigma* ( $n = 13$ ), *Pseudoblennius* sp. “Kirin-anahaze” ( $n = 9$ ), *Ps. percoides* ( $n = 8$ ), *F. osimae* ( $n = 3$ ) and *F. ishikawae* ( $n = 1$ ). Five of six species mainly employed *H. ritteri* as their hosts (Table 1; Fig. 2f). *Pseudoblennius cottoides* and *Ps. percoides* also utilised *Py. sacciformis*. Egg masses deposited in *H. roretzi* were identified as belonging to *Pseudoblennius* sp. “Kirin-anahaze” (Fig. 2g). *Furcina osimae* only utilised *Py. sacciformis* as the tunicate host (Fig. 2h). No egg masses were genetically assigned to *V. minutus* and so further analyses excluded this species.

### Egg mass position in different hosts

Since egg masses were visible from the outside of sponges and colonial tunicates, but were

not for solitary tunicates, the egg mass position would be expected to differ between the hosts. As expected, the egg masses were deposited significantly deeper in solitary tunicates ( $16.44 \pm 6.39$  mm,  $n = 73$ ) than in sponges ( $2.70 \pm 1.69$  mm,  $n = 6$ ) and colonial tunicates ( $2.02 \pm 1.79$  mm,  $n = 11$ ) (one-way ANOVA,  $F_{2,87} = 40.46$ ,  $p < 0.0001$ ; Tukey's HSD,  $p < 0.0001$  for both). The egg mass position did not differ between sponges and colonial tunicates (Tukey's HSD,  $p = 0.97$ ). This indicates that ovipositor length likely differs between sculpin species using solitary tunicates and those using sponges and colonial tunicates.

### **Comparison of ovipositor length between fish using different deposition sites**

An ANCOVA analysis showed that the interaction between deposition site and log SL did not have a significant effect on log OL (full model,  $F_{3,129} = 2.09$ ,  $p = 0.11$ ), and overall, larger species/individuals had longer ovipositors (reduced model,  $F_{1,132} = 176.77$ ,  $p < 0.0001$ ; Fig. 4a). As predicted, the relative OL was significantly related to deposition site (reduced model,  $F_{3,132} = 557.23$ ,  $p < 0.0001$ ; Fig. 4a). The relative OL of the ostracophilous sculpins was significantly longer than that of the non-ostracophilous sculpins (Fig. 4b–e). Moreover, the sculpin species that use solitary tunicates as hosts had much longer OLs (ca. 20 % of SL) than that of those using colonial tunicates and sponges. Although the relative OL differed significantly between *V. centropomus* (ca. 5 % of SL) and *Ps. marmoratus* (ca. 3 % of SL), the absolute OL did not differ between them ( $t$ -test,  $t = 0.99$ ,  $p = 0.33$ ; Fig. 4a–c).

### **Ovipositor length and the size of solitary tunicates used by sculpins**

The body size of the six parental species that utilised solitary tunicates as hosts differed significantly among species (one-way ANOVA,  $F_{5,89} = 166.84$ ,  $p < 0.0001$ ; Fig. 5a). Female *Pseudoblennius* sp. “Kirin-anahaze” were the largest, and two species in the *Furcina* genus were the smallest of all six species. Overall, larger species had longer ovipositors ( $F_{5,89} =$

28.05,  $p < 0.0001$ ; Fig. 5b), but the OL of *Ps. percooides* was significantly shorter than that of *Ps. cottoides* despite its larger body size (also see Fig. 4a).

As expected, egg mass position was deeper when larger tunicates were used as hosts (Pearson's  $r = 0.49$ ,  $p < 0.0001$ ,  $n = 72$ ). The LD of tunicates used as hosts and the egg mass position were significantly different among sculpin species (LD: Fig. 5c; one-way ANOVA,  $F_{4,79} = 8.81$   $p < 0.0001$ ; egg mass position: Fig. 5d;  $F_{4,68} = 7.05$ ,  $p < 0.0001$ ). Consistent with the expectation that utilisation of larger tunicates will require longer ovipositors, the LD of tunicates used as hosts and the egg mass position were closely related to the sculpin OL (Fig 5); *Pseudoblennius* sp. “Kirin-anahaze”, *Ps. cottoides* and *Ps. zonostigma* that have longer ovipositors spawned in a deeper position in larger tunicates, and the genus *Furcina* with a smaller body size and a shorter ovipositor consequently utilised smaller tunicates.

*Pseudoblennius percooides*, which has a larger body size but a shorter ovipositor, selected smaller tunicates as spawning hosts.

*Pseudoblennius* sp. “Kirin-anahaze”, *Ps. cottoides* and *Ps. zonostigma* used similar-sized *H. ritteri* tunicates as hosts (Fig. 5c), indicating severe competitions for hosts among the three species. However, if host *H. ritteri* is abundant and sculpins use the most common size of tunicates, competition for hosts would be weakened. Indeed, *H. ritteri* was used by only 11 % of all individuals (Fig. 3d). Additionally, the average and variance of host size were similar between unused *H. ritteri* (84.20 LD  $\pm$  20.31,  $n = 595$ ) and that used by *Pseudoblennius* sp. “Kirin-anahaze” (92.84 LD  $\pm$  10.37,  $n = 7$ ,  $t$ -test,  $t = 1.12$ ,  $p = 0.26$ ;  $F$  test to compare two variances,  $F = 3.84$ ,  $p = 0.09$ ), *Ps. cottoides* (87.16 LD  $\pm$  18.38,  $n = 46$ ,  $t = 0.96$ ,  $p = 0.34$ ;  $F = 1.22$ ,  $p = 0.41$ ) or *Ps. zonostigma* (although slightly larger tunicates were used: 99.12 LD  $\pm$  17.37,  $n = 11$ ,  $t = 2.42$ ,  $p = 0.02$ ;  $F = 1.37$ ,  $p = 0.61$ ). These show that these three sculpins used the most common size of *H. ritteri* (Fig. 3d).



## Discussion

### Host specificity

In this study, we identified oviposition hosts in eight out of the nine ostracophilous sculpins examined. While a number of researchers and divers have observed spawning behaviours of some of these sculpins, such as *Ps. marmoratus*, *Ps. percoides* and *Ps. zonostigma* (Shinomiya 1985; Okamura and Amaoka 1997; Abe and Munehara 2009; Uryu 2011), so far only one study has reported host tunicate selection by *Ps. percoides* (Nishida et al. 2008). Thus, to our knowledge, this is the first study to confirm the host species used by these eight sculpin species, including *Ps. percoides*. There are three patterns of host use by sculpins, such as species that mainly use colonial tunicates, those that only use sponges and those that only use solitary tunicates for oviposition. This result mostly confirms our first prediction, that ostracophilous sculpins would show host specificity. However, as found in ostracophilous bitterlings (Reichard et al. 2007), sculpins also showed variation in their level of specificity. *Pseudoblennius marmoratus* oviposited only in the spongocoel of sponges, but *V. centropomus* utilised colonial tunicates and sponges for oviposition. Eggs of six sculpin species were found in solitary tunicates, but three of these fishes used only a single tunicate species and the others used two tunicate species.

Egg mass numbers for each sculpin species collected in this study mostly corresponded to the abundance of parental species at the study sites (see Fig. 4, Table 1). Although we captured most tunicates on the wall of Kitakoura Port, the consistency of appearance frequencies between eggs and parental species implies that the six sculpin species do not avoid the artificial habitat for spawning. Since eggs of *V. minutus* were not found and only one egg mass of *F. ishikawae* was collected, further field sampling including different sites and seasons are needed to clarify their host use.

Sponges are only distantly related to colonial tunicates (e.g. Dunn et al. 2014). Their body structures are therefore notably different. However, they share morphological similarities: both have myriad brachial apertures and shared, common cloacal apertures. These similarities, especially the common structure of the cloacal apertures, may allow female *V. centropomus* to utilise both colonial tunicates and sponges as hosts. To our knowledge, it is unlikely that any other fishes utilise such distantly related taxa. *Pseudoblennius marmoratus* only oviposits in sponges, which is consistent with previous underwater observations (Uryu 2011). It remains unclear whether *Ps. marmoratus* also utilises colonial tunicates. To confirm this, more information on its spawning behaviour in the field needs to be gathered. These two sculpin species have short and similarly sized ovipositors, suggesting that they are specialised in ovipositing in colonial tunicates and sponges.

We found that sculpins used only three out of seven species of solitary tunicates as hosts, even though many species of tunicates co-occurred at the study sites. There are three possibilities to explain sculpin host selection: 1) the size of the tunicates, 2) the internal structure of the tunicates and 3) chemical substances in the atria of the tunicates. First, sculpins may select for tunicates of optimal size. Solitary tunicate spawners have developed long ovipositors, which may allow the placement of their eggs in the atrium of tunicates via the atrial siphon. The elongation of the ovipositor is limited by the cost associated with growing and maintaining a long ovipositor. Sculpin females did not use the large tunicates *M. hartmeyeri* and *Herdmania japonica*, probably due to the excessively long distance from the entrance of atrial siphon to the atrium of the tunicate. In contrast, the small atrium and narrow entrance of the atrial siphon of the smallest tunicate, *C. irene*, was too small to allow sculpins to use them as hosts.

Second, the internal structure of tunicates may be related to host selection by sculpins. Our data showed that the size range of unused tunicate *Po. c. kroboja* greatly

overlapped with that of the commonly used tunicate *H. Ritteri*. The unused *Po. c. kroboja* tunicates are more compressed compared with the *H. Ritteri* tunicates (Awata S. pers. obs.), suggesting that sculpins do not utilise *Po. c. kroboja* due to the small inner space of their atria. Finally, substances in the atria of tunicates might also affect host use by female sculpins. A number of tunicates produce chemical defences, such as high sulfuric acid and high vanadium content, which may reduce their predation risks (e.g. Stoecker 1980; Paul and Puglisi 2004; Paul et al. 2006). *Polycarpa cryptocarpa kroboja* has a uniquely strong smell (Awata S. pers. obs.) compared to the other tunicate species dissected. Although the chemical components responsible for this smell were not identified, they may have adverse effects on the fish eggs. Our aquarium experiments have demonstrated that when *Ps. cottoides* females with ripe eggs were allowed to choose *H. Ritteri* and *Po. c. kroboja* of the similar size, they spent more time inspecting *H. Ritteri*, and eggs were deposited only in *H. Ritteri* (Awata S, Igarashi N, unpub. data). This indicates that the female sculpins likely discriminate between *H. Ritteri* and *Po. c. kroboja* prior to spawning, and clearly prefer the former over the latter as oviposition hosts. It remains to be experimentally determined which combinations of these proximate factors such as tunicate size, structure and chemical substances affect host selection by female sculpins.

### **Ovipositor length**

In accordance with our second prediction, only ostracophilous sculpins had elongated ovipositors, while non-ostracophilous sculpins lacked this feature. Since non-ostracophilous sculpins have been suggested to be the ancestral state (Yabe 1985; Koya et al. 2011; Munehara et al. 2011; Knope 2013), we can conclude that the elongated ovipositor has evolved as an adaptation to the ostracophilous reproduction strategy. The ovipositor in species using solitary tunicates as hosts was much longer than that in species using colonial tunicates and sponges. The position of eggs oviposited in colonial tunicates and sponges were as short

as they were observed to be from the outside, and perfectly matched the short ovipositor length of *V. centropomus* and *Ps. marmoratus*. In contrast, sculpins that oviposited in solitary tunicates had extremely long ovipositors, which in turn matched the distance from the entrance of the atrial siphon to the edge of the egg masses. These results are consistent with the third prediction, that ovipositor length would reflect the species and size of hosts utilised.

Both the branchial and atrial siphons of solitary tunicates close immediately by ejecting the water from their body when they detect external stimuli such as vibrations in the water. In contrast, sponges and colonial tunicates shrink a little after the detection of such stimuli (Awata S. pers. obs.). In response to such host behaviour, female sculpins that use solitary tunicates as hosts insert their ovipositors rapidly into the atrium (Movie S1, S2, Supplementary Information). In contrast, species that use sponges as hosts oviposit slowly (Movie S3). Based on these observations, we suggest that spawning behaviour and ovipositor length may have adapted to suit the different hosts.

### **Competition among sculpin species over hosts**

In this study, six species of sculpins were determined to use solitary tunicates as hosts.

*Furcina osimae* only used the tunicate *Py. sacciformis* as a spawning host, but the other five species mainly deposited their eggs into *H. Ritteri*. This suggests these five species did not prefer specific host tunicates. However, their host use may be also related to the size and species of the host tunicates. Indeed, two smallest species of genus *Furcina*, which have shorter ovipositors, used smaller tunicates; the largest *Pseudoblennius* sp. “Kirin-anahaze”, which has longer ovipositors, spawned in larger tunicates. In contrast, the ovipositor of *Ps. percoides* was shorter than that of *Ps. cottoides*, despite its larger body, and matches the size of the host tunicates. Medium-sized sculpins such as *Ps. cottoides* and *Ps. zonostigma* had medium-sized ovipositors but used larger tunicates as hosts. Thus, as expected from our first

prediction, sculpins likely choose different sized hosts that fit their ovipositor size, probably to reduce interspecific competition for hosts (Reichard et al. 2007).

However, three sculpins (*Pseudoblennius* sp. “Kirin-anahaze”, *Ps. cottoides* and *Ps. zonostigma*) showed similar host choice in terms of species and size, indicating that interspecific competition for hosts may be unavoidable (Kitamura 2007; Reichard et al. 2007). Previous studies have shown that host utilisation by female bitterlings may be determined by host availability, as well as by host preference and interspecific competition among bitterlings (Kitamura 2007; Reichard et al. 2007). In this study, the most abundant host, *H. ritteri*, was used by only 11 % of all individuals. Furthermore, these three sculpins used the most common size of tunicates (65–105 mm in LD of host *H. ritteri*; see Fig. 3d), although *Ps. zonostigma* oviposited into somewhat larger tunicates. Therefore, their host preferences are strongly related to host availability, and the ovipositor length of those three species may have been shaped by the size of the most abundantly available tunicates. Further experimental studies are needed to identify the “real” host size and species preference by controlling for host availability and interspecific competition among ostracophilous sculpins (Reichard et al. 2007). Since most *Pseudoblennius* species are piscivores (Okamura and Amaoka 1997; Uryu 2011), it would be necessary to take the risk of predation of smaller species and individuals by larger ones into account when determining interspecific competition at the spawning sites.

### **Costs and benefits for ostracophilous sculpins**

There are at least three potential benefits for female sculpins that use tunicates and sponges as oviposition hosts. First, the hosts may provide suitable environments for embryonic development (Hunter 1969; Munehara 1991; Spence and Smith 2013). Both tunicates and sponges are filter feeders that draw water and food particles through incurrent siphon or canals and excrete the filtered water through atria and excurrent siphons, or through

spongocoels and oscula. Parental care by fishes that exclusively exhibit care for their eggs mainly consists of fanning and mouthing, which are assumed to increase the oxygen supply and to help to remove dead eggs and dust particles (e.g. Keenleyside 1991). Since filtered water constantly flows through the atria of tunicates and the spongocoels of sponges where the eggs are deposited, the hosts may inadvertently serve a similar function to fanning and mouthing for the eggs. Furthermore, sponges may provide another benefit to egg development in sculpins. Many studies have reported that sponges contain compounds with antibacterial and antifungal properties (e.g. reviewed in Mehbub et al. 2014). Considering the absence of dead eggs in sponges, the antibacterial and antifungal activities of sponges may increase the survival rate of sculpin eggs. Although dead, un-decayed eggs of the little dragon sculpin *B. cirrhosus* have been found to be present in sponge tissues (Munehara 1991), such eggs were not found in our study, perhaps due to the difference in embryonic periods (< 1 month in our study species and > 8 months in *B. cirrhosus*; Munehara 1991). Rose bitterling embryos developing on the gills of freshwater mussels are known to gain nutrients and oxygen from the water taken in by the mussels (Spence and Smith 2013). However, this may not be the case in sculpins, because their eggs are placed where few or no nutrients may be available due to the filtered water, and larvae are likely to leave their host soon after hatching since they are capable of active swimming (Kimura et al. 1987, 1988; Awata S. pers. obs.).

Second, eggs are likely protected from predators by being hidden inside the hosts (Hunter 1969; Munehara 1991). Since eggs are completely hidden in solitary tunicates that have rigid bodies, predation avoidance may be highly effective. Furthermore, many species of sponges are chemically defended from predation by marine animals, such as fishes, sea urchins, sea stars and hermit crabs (e.g. reviewed in Paul and Puglisi 2004; Paul et al. 2006). Thus, oviposition in sponges may also be effective for predation avoidance due to their chemical defences. Third, females are not required to provide parental care, which may allow

them to utilise their energy reserves for further reproduction, such as increasing egg numbers and size.

However, it is costly to be ostracophilous. First, the development of elongated ovipositors and unique spawning behaviour are essential. Second, during the spawning season, sculpin females must seek out suitable, limited oviposition hosts, which might increase the probability of predation and consume energy reserves. Despite such costs, the benefits for female sculpins specialising in the use of tunicates and sponges outweigh the costs, which may be the evolutionary forces that produce and maintain the unique spawning behaviours observed in sculpins. This also may be the case for other ostracophilous fishes such as bitterlings, snailfishes and Japanese tubesnouts (Leung 2014).

### **Costs and benefits for hosts**

In contrast to the potential benefits obtained by the sculpins, it is unlikely that sculpins and their eggs have positive impacts on the host tunicates and sponges. Conversely, the eggs deposited by sculpins may adversely affect their hosts. One possible cost for the hosts harbouring sculpin egg masses is the reduction of sea water exchange, as observed in mussels (Smith et al. 2001; Mills and Reynolds 2002, 2003; Kitamura 2005). In many cases, one egg mass was found in a solitary tunicate, and occupied only a small space in the atrium.

However, the eggs, especially those oviposited by larger sculpin species such as *Pseudoblennius* sp. “Kirin-anahaze”, sometimes entirely fill up the tunicate atrium due to their large clutch size (see Table S3). In this case, the hosts cannot effectively exchange seawater and consequently might incur some costs such as lowered growth and survival from the overload of fish eggs. Such an overload of fish eggs has been reported for the snailfish-crab system, where the load of the egg masses in the gill cavity of crabs deposited by snail fish caused severe damage to their gills (Love and Shirley 1993; Somerton and Donaldson

1998; Poltev and Mukhametov 2009). Spence and Smith (2013) have also found that some host mussels with many bitterling eggs died, probably due to severe competition for oxygen between the host mussels and bitterling embryos. Since sculpin eggs are not oviposited on the gills of tunicates through the incurrent siphon, sculpin eggs unlikely do cause severe damage to the respiratory system of sponges and tunicates. Experimental work is needed to ascertain the cost to the hosts of sculpin eggs.

## **Conclusions**

In summary, we have shown host selection and variation of ovipositor length in eight out of nine sympatric marine sculpins found off the shore of an island in the Sea of Japan. To date, ostracophilous behaviour has only been described for a small number of fish species, and therefore only limited data on the behavioural and morphological adaptations of parasites and hosts are available, with the exception of the bitterlings. A more recent study, however, has found evidence of ostracophilous behaviour in the tubenose poacher *Pallasina barbata* (Momota and Munehara 2017). Sculpins and tubenose poachers are different fish groups from other ostracophilous fishes (Leung 2014), suggesting that ostracophilous reproductive strategy has likely evolved in parallel across different taxa (Leung 2014). This study has shown the morphological adaptations of sculpin females at the interspecific level. However, ongoing research will need to examine the intraspecific variation in ovipositor size and morphology, depending on the host species used and their size. Studying ostracophils at the population level might be particularly worthwhile to identify the (co-)evolution of behavioural and morphological adaptations of parasites and their hosts.

## **Ethical statement**



The research reported in this study was carried out in accordance with Animal Care and Use Committees at Niigata University and Osaka City University. All of the procedures described above meet the ABS/ASAB guidelines for the ethical treatment of animals. Research permission was obtained from Fisheries Cooperative Association of Sado, Japan.

## **Acknowledgements**

We thank Teruaki Nishikawa and Yuji Ise for help with the classification of tunicates and sponges. We are also grateful to Mitsuo Homma (Diving Service F. Wave), Ryo Honma (Sado Diving Centre), Yoshihisa Sato (Senkakuwan Ageshima Aquarium), Akihiro Yamada, Hiromitsu Takashima (Ogi Diving Centre), Tadashi Shoji (Diving Service S. World), Sadogashima SCUBA Diving Association and Kitakoura Recreational Fishing Cooperative for support in the field. We would like to thank Tomonobu Uryu for creating videos for us and Editage (<http://www.editage.jp>) for English language editing. The anonymous reviewers provided helpful comments on the manuscript. The project was financially supported in part by JSPS KAKENHI Grant Numbers JP24770016 and JP16H04841 to S.A., JP26450259 to Y.K. and JP25304011 to H.M. and by the Sasaki Environment Technology Foundation (H26) to S.A.

## **References**

- Akagawa I, Hara M, Iwamoto T (2008) Egg concealment in tunicates by females of the Japanese tubenout, *Aulichthys japonicus* (Gasterosteiformes), and its subsequent copulation. *Ichthyol Res* 55:85–89
- Abe T, Munehara H (2009) Adaptation and evolution of reproductive mode in copulating cottoid species. In: Jamieson BGM (Ed.) *Reproductive Biology and Phylogeny of Fishes (Agnathans and Bony Fishes)*. Science Publishers, Enfield, pp 221–246
- Andersson M (1994) *Sexual selection*. Princeton Univ. Press, Princeton
- Awata S (2015) Diversity and evolution of reproductive strategies in marine sculpins.

Aquabiol 37:614–621

- Awata S (2017) Taxonomic and ecological notes on marine sculpins on the coast of Sado Island in the Sea of Japan. *Nat Hist Sado Isl* 5:9–20
- Balon EK (1975) Reproductive guilds of fishes – proposal and definition. *J Fish Res Board Can* 32:821–864
- Birkhead TR, Møller AP (1998) Sperm competition and sexual selection. Academic Press, London
- Busby MS, Blood DM, Fleischer AJ, Nichol DG (2012) Egg deposition and development of eggs and larvae of bigmouth sculpin (*Hemitripterus bolini*). *Northwest Nat* 93:1–16
- Dunn CW, Giribet G, Edgecombe GD, Hejnol A (2014) Animal phylogeny and its evolutionary implications. *Annu Rev Ecol Evol Syst* 45:371–395
- Elias LG, Kjellberg F, Farache FHA, Almeida EAB, Rasplus J-Y, Cruaud A, Peng Y-Q, Yang D-R, Pereira RAS (2018) Ovipositor morphology correlates with life history evolution in agaonid fig wasps. *Acta Oecol* 90:109–116
- Gardner JR, Orr JW, Stevenson DE, Spies I, Somerton DA (2016) Reproductive parasitism between distant phyla: molecular identification of snailfish (Liparidae) egg masses in the gill cavities of king crabs (Lithodidae). *Copeia* 104:645–657
- Ghara M, Kundanati L, Borges RM (2011) Nature’s Swiss Army knives: ovipositor structure mirrors ecology in a multitrophic fig wasp community. *PLoS ONE* 6:e23642
- Hunter CJ (1969) Confirmation of symbiotic relationship between liparid fishes (*Careproctus* spp.) and male king crab (*Paralithodes camtschatica*). *Pac Sci* 23:546–547
- Iwata A (1983) A revision of the cottid fish genus *Vellitor*. *Jap J Ichthyol* 30:1–9
- Keenleyside MHA (1991) Parental care. In: Keenleyside MHA (Ed.) *Cichlid fishes: behaviour, ecology and evolution*. Chapman and Hall, London, pp 191–208
- Kimura S, Tsumoto K, Mori K (1987) Development of eggs, larvae and juveniles of the cottid fish, *Pseudoblennius cottoides*, reared in the laboratory. *Jap J Ichthyol* 34: 346–350
- Kimura S, Tsumoto K, Mori K (1988) Development of the cottid fish, *Pseudoblennius percoides*, reared in the laboratory, with brief descriptions of juvenile *P. marmoratus* and *P. zonostigma*. *Jap J Ichthyol* 35:19–24
- Kitamura J (2005) Factors affecting seasonal mortality of rosy bitterling (*Rhodeus ocellatus kurumeus*) embryos on the gills of their host mussel. *Popul Ecol* 47:41–51
- Kitamura J (2007) Reproductive ecology and host utilization of four sympatric bitterling (Acheilognathinae, Cyprinidae) in a lowland reach of the Harai River in Mie, Japan. *Environ Biol Fish* 78:37–55
- Kitamura J, Nagata N, Nakajima J, Sota T (2012) Divergence of ovipositor length and egg shape in a brood parasitic bitterling fish through the use of different mussel hosts. *J Evol Biol* 25:566–573
- Knobe ML (2013) Phylogenetics of the marine sculpins (Teleostei: Cottidae) of the North American Pacific Coast. *Mol Phylogenet Evol* 66:341–349
- Koya Y, Hayakawa Y, Markevich A, Munehara H (2011) Comparative studies of testicular

- structure and sperm morphology among copulatory and non-copulatory sculpins (Cottidae: Scorpaeniformes: Teleostei). *Ichthyol Res* 58:109–125
- Koya Y, Mitsunashi N, Awata S, Ito T, Munehara H (2015) Identification of the reproductive mode for internal gamete association in *Vellitor centropomus* (Cottidae): gonadal histological analysis. *Japan J Ichthyol* 62:121–131
- Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Mol Biol Evol* 33:1870–1874
- Le Ralec A, Rabasse JM, Wajnberg E (1996) Comparative morphology of the ovipositor of some parasitic Hymenoptera in relation to characteristics of their hosts. *Can Entomol* 128:413–433
- Leung TLF (2014) Fish as parasites: an insight into evolutionary convergence in adaptations for parasitism. *J Zool* 294:1–12
- Love DC, Shirley TC (1993) Parasitism of the golden king crab, *Lithodes aequispinus* Benedict, 1895, by a liparid fish. *Crustaceana* 65: 97–104
- Mehbub MF, Lei J, Franco C, Zhang W (2014) Marine sponge derived natural products between 2001 and 2010: trends and opportunities for discovery of bioactives. *Mar Drugs* 12:4539–4577
- Momota K, Munehara H (2017) Reproductive ecology and morphological changes during the early life stages of *Pallasina barbata* (Steindachner, 1876). *Bull Fish Sci Hokkaido Univ* 67:7–12
- Mills SC, Reynolds JD (2002) Mussel ventilation rates as approximate cue for host selection by bitterling, *Rhodeus sericeus*. *Oecologia* 131:473–478
- Mills SC, Reynolds JD (2003) The bitterling-mussel interaction as a test case for co-evolution. *J Fish Biol* 63:84–104
- Munehara H (1991) Utilization and ecological benefits of a sponge as a spawning bed by the little dragon sculpin *Blepsias cirrhosis*. *Jap J Ichthyol* 38:179–184
- Munehara H, Goto A, Yabe M (2011) Diversity of Cottoid Fishes -Adaptation and Evolution-. Tokai Univ Press, Kanagawa
- Munehara H, Takano K, Koya Y (1989) Internal gametic association and external fertilization in the elkhorn sculpin, *Alcichthys alcicornis*. *Copeia* 1989:673–678
- Nakabo T, Kai Y (2013) Cottidae. In: Nakabo T (Ed.) *Fishes of Japan with pictorial keys to the species* third edition. Tokai University Press, Hadano, pp 1160–1188, 2061–2067
- Nelson JS, Grande TC, Wilson MVH (2016) *Fishes of the World*, 5th ed. John Wiley & Sons, Inc, New Jersey
- Nishida T, Inui R, Onikura N (2008) A note on the spawning bed of *Pseudoblennius percoides* (Scorpaeniformes, Cottidae) in shallow sea areas around coastal Fukutsu, northern Kyusyu Island, Japan. *Biogeography* 10:45–51
- Okamura O, Amaoka K Eds. (1997) *Seawater Fishes in Japan*. Yamatokeikoku-sha, Tokyo
- Paul VJ, Puglisi MP (2004) Chemical mediation of interactions among marine organisms. *Nat Prod Rep* 21:189–209

- Paul VJ, Puglisi MP, Ritson-Williams R (2006) Marine chemical ecology. *Nat Prod Rep* 23:153–180
- Palumbi SR (1996) What can molecular genetics contribute to marine biogeography? An urchin's tale. *J Exp Mar Biol Ecol* 203:75–92
- Peden AE, Corbett CA (1973) Commensalism between a liparid fish, *Careproctus* sp., and the lithodid box crab, *Lopholithodes foraminatus*. *Can J Zool* 51:555–556
- Poltev YN, Mukhametov IN (2009) Concerning the problem of carcinophilia of *Careproctus* species (Scorpaeniformes: Liparidae) in the North Kurils. *Russ J Mar Biol* 35:215–223
- Reichard M, Liu H, Smith C (2007) The co-evolutionary relationship between bitterling fishes and freshwater mussels: insights from interspecific comparisons. *Evol Ecol Res* 9:239–259
- Reichard M, Ondrackova M, Przybylski M, Liu H, Smith C (2006) The costs and benefits in an unusual symbiosis: experimental evidence that bitterling fish (*Rhodeus sericeus*) are parasites of unionid mussels in Europe. *J Evol Biol* 19:788–796
- Schmidt TR, Gold JR (1993) Complete sequence of the mitochondrial cytochrome *b* gene in the cherryfin shiner, *Lythrurus roseipinnis* (Teleostei: Cyprinidae). *Copeia* 1993:880–883
- Shinomiya A (1985) Studies on the reproductive physiology and ecology in three marine cottid fish. Dissertation, Hokkaido University
- Shinomiya A, Ikemoto M (1987) Spawning habits of the sculpin *Pseudoblennius percoides* in relation to sea squirt. In: Advance abstracts for the 20th annual meeting, The Ichthyological Society of Japan, p 26
- Shiogaki M, Dotsu Y (1974) The spawning of the sea sculpin, *Pseudoblennius cottoides*. *Bull Fac Fish Nagasaki Univ* 38:71–76
- Smith C, Reichard M, Jurajda P, Przybylski M (2004) The reproductive ecology of the European bitterling (*Rhodeus sericeus*). *J Zool* 262:107–124
- Smith C, Rippon K, Douglas A, Jurajda P (2001) A proximate cue for oviposition site choice in the bitterling (*Rhodeus sericeus*). *Freshw Biol* 46:903–911
- Somerton DA, Donaldson W (1998) Parasitism of the golden king crab, *Lithodes aequispinus*, by two species of snailfish, genus *Careproctus*. *Fish Bull* 96:871–884
- Spence R, Smith C (2013) Rose bitterling (*Rhodeus ocellatus*) embryos parasitize freshwater mussels by competing for nutrients and oxygen. *Acta Zool* 94:113–118
- Stoecker D (1980) Chemical defenses of ascidians against predators. *Ecology* 61:1327–1334
- Tamura K, Nei M (1993) Estimation of the number of nucleotide substitutions in the control region of mitochondrial-DNA in humans and chimpanzees. *Mol Biol Evol* 10:512–526
- Uchida K (1932) Fish laying eggs in the body of tunicates. *Kagaku* 2:56–57
- Uchida K (1979) *Chigyo Wo Motomete*. Iwanamisyoten, Tokyo
- Uryu T (2011) *Marine Fishes of Izu*. Kaiyusha, Tokyo
- Yabe M (1985) Comparative osteology and myology of the superfamily Cottoidea (Pisces: Scorpaeniformes), and its phylogenetic classification. *Mem Fac Fish Hokkaido Univ*

32:1–130

## Figure legends

**Fig. 1** Location of sampling sites. (a) Sado Island located 45 km north west of Niigata, Japan. We collected sculpins that oviposit on rock surface at Usujiri (41° 56' 11.85" N, 140° 56' 54.66" E), Hokkaido, Japan. (b) Four sampling sites on the coast of the island. Kitakoura: 38° 16' 26.90" N, 138° 30' 34.42" E; Mushizaki: 38° 14' 58.66" N, 138° 30' 24.37" E; Tassya: 38° 04' 28.64" N, 138° 14' 40.81" E and; Kotoura: 37° 48' 13.43" N, 138° 15' 08.26" E. (c) Kitakoura Port. Monthly collection of tunicates was conducted along the south breakwater of the port (indicated by an ellipse).

**Fig. 2** Examples of photographs of eggs deposited in sponges and tunicates. (a) Egg masses deposited by females of *Pseudoblennius marmoratus* in sponges. (b) Eyed eggs of *Vellitor centropomus* in a colonial tunicate. (c) Colonial tunicate; at least five egg masses were deposited in this colony. Colony width: c.a. 100 mm. (d) *Halocynthia ritteri*, the solitary tunicates most frequently used by sculpins. (e) Underwater photograph of *H. roretzi*. (f) Eyed eggs of *Ps. cottoides* found in *H. ritteri*. (g) Eyed eggs of *Pseudoblennius* sp. "Kirin-anahaze" deposited in *H. roretzi*. (f) Eyed *Furcina osimae* eggs in *Pyura sacciformis*. Scale bars indicated in (a), (b), (f), (g) and (h): 10 mm.

**Fig. 3** Size frequency distribution of solitary tunicates collected during study periods. (a) *Microcosmus hartmeyeri*, (b) *Herdmania japonica*, (c) *Halocynthia roretzi*, (d) *Halocynthia ritteri*, (e) *Polycarpa cryptocarpa kroboja*, (f) *Pyura sacciformis* and (g) *Cnemidocarpa irene*. Panels are arranged in decreasing order of the average long diameter (LD) of tunicates. *White, grey and black bars* indicate tunicates with sculpin egg masses, with Japanese tubesnout egg masses and without any egg masses, respectively.  $n_{sc}$ : number of tunicates with sculpin eggs,

$n_{tu}$ : number of tunicates with eggs of Japanese tubesnout (the details of which will be published in a separate paper) and  $n_{ne}$ : number of tunicates with no fish eggs.

**Fig. 4** (a) Ovipositor length (OL) increased with standard length (SL) of sculpins and differed among sculpins that utilised four different types of hosts. *Circles, triangles and squares* denote the species that used solitary tunicates, colonial tunicates and sponges, and rock surfaces as deposition sites, respectively. Regression lines for different types of hosts are from the reduced model of ANCOVA. Relative OL were statistically significantly different between all combinations of the four types (Tukey's HSD,  $p < 0.05$ ). (b)–(e) Examples of photographs of ovipositors (indicated by the arrows) in the sculpins of the four types. (b) *Pseudoblennius cottoides*; (c) *Vellitor centropomus*; (d) *Ps. marmoratus*; and (e) *Alcichthys alcicornis*. Scale bars: 10 mm.

**Fig. 5** Body size and ovipositor length of parasitic sculpins were closely related to the size of solitary tunicates selected by each sculpin species as spawning hosts. (a) Female standard length (SL) and (b) ovipositor length (OL) of six species of sculpins that used solitary tunicates as hosts. (c) Long diameter (LD) and (d) egg mass position of tunicates utilised as hosts. Means ( $\pm$  SD) with different letters are significantly different (Tukey's HSD,  $p < 0.05$ ). Sample sizes are indicated as numerals in bars. *ND*, no data. Bars are arranged in decreasing order of the average SL of sculpins from left to right.




Figure 1











Figure 2




Figure 3





Figure 4




Figure 5

**Table 1** Host (solitary tunicates) use by sculpins and Japanese tubesnout. The number of tunicates where single, double and triple clutches of fish were found is shown in parenthesis.

Host tunicate species	Total number of tunicates	Total number of tunicates with sculpin egg masses	Number of tunicates used by sculpins							Number of tunicates used by <i>Aulichthys japonicus</i>
			<i>Pseudoblennius cottoides</i>	<i>Ps. zonostigma</i>	<i>Pseudoblennius</i> sp. "Kirin-anahaze"	<i>Ps. percoides</i>	<i>Furcina osimae</i>	<i>F. ishikawae</i>	Unidentified	
<i>Halocynthia Ritteri</i>	678	75 (69, 5, 1)	46 (42, 3, 1)	11 (9, 2, 0)	8	6	0	1	3	10 (9, 1, 0)
<i>Polycarpa cryptocarpa kroboja</i>	307	0	0	0	0	0	0	0	0	0
<i>Cnemidocarpa irene</i>	141	0	0	0	0	0	0	0	0	0
<i>Pyura sacciformis</i>	42	6	1	0	0	2	3	0	0	6 (5, 0, 1)
<i>Microcosmus hartmeyeri</i>	24	0	0	0	0	0	0	0	0	0
<i>Halocynthia roretzi</i>	19	1	0	0	1	0	0	0	0	1
<i>Herdmania japonica</i>	1	0	0	0	0	0	0	0	0	0
Total	1212	82 (76, 5, 1)	47 (43, 3, 1)	11 (9, 2, 0)	9	8	3	1	3	17 (15, 1, 1)

## Electronic Supplementary Material

### Marine Biology

#### Host selection and ovipositor length in eight sympatric species of sculpins that deposit their eggs into tunicates or sponges

Satoshi Awata<sup>1,2</sup> · Haruka Sasaki<sup>2</sup> · Tomohito Goto<sup>2</sup> · Yasunori Koya<sup>3</sup> · Hirohiko Takeshima<sup>4,5</sup> · Aya Yamazaki<sup>6</sup> · Hiroyuki Munchara<sup>6</sup>

*<sup>1</sup>Laboratory of Animal Sociology, Graduate School of Science, Osaka City University, Osaka, Japan, <sup>2</sup>Sado Marine Biological Station, Faculty of Science, Niigata University, Sado, Japan, <sup>3</sup>Department of Biology, Faculty of Education, Gifu University, Gifu, Japan, <sup>4</sup>Department of Marine Biology, School of Marine Science and Technology, Tokai University, Shizuoka, Japan, <sup>5</sup>Research Institute for Humanity and Nature, Kyoto, Japan, <sup>6</sup>Usujiri Fisheries Station, Field Science Center for Northern Biosphere, Hokkaido University, Hakodate, Japan*

Correspondence

Satoshi Awata

Laboratory of Animal Sociology, Graduate School of Science, Osaka City University, 3-3-138 Sugimoto, Sumiyoshi, Osaka 558-8585, Japan.

Tel.: +81 6 6605 2607

e-mail: sa-awata@sci.osaka-cu.ac.jp



<i>Vellitor centropomus</i> -17	<i>Vellitor centropomus</i>	Kitakoura, Sado, Japan	March-13	same sequence as LC424816
<i>Vellitor centropomus</i> -18	<i>Vellitor centropomus</i>	Kitakoura, Sado, Japan	March-13	same sequence as LC424820
<i>Vellitor centropomus</i> -19	<i>Vellitor centropomus</i>	Tassya, Sado, Japan	October-13	same sequence as LC424820
<i>Vellitor centropomus</i> -20	<i>Vellitor centropomus</i>	Tassya, Sado, Japan	October-13	same sequence as LC424820
<i>Vellitor centropomus</i> -21	<i>Vellitor centropomus</i>	Tassya, Sado, Japan	October-13	same sequence as LC424820
<i>Vellitor centropomus</i> -22	<i>Vellitor centropomus</i>	Sado, Japan	November-13	same sequence as LC424820
<i>Vellitor centropomus</i> -23	<i>Vellitor centropomus</i>	Sado, Japan	November-13	LC424819
<i>Vellitor centropomus</i> -24	<i>Vellitor centropomus</i>	Sado, Japan	November-13	same sequence as LC424820
<i>Pseudoblennius zonostigma</i> -3	<i>Pseudoblennius zonostigma</i>	Kitakoura, Sado, Japan	December-12	LC424821
<i>Pseudoblennius zonostigma</i> -4	<i>Pseudoblennius zonostigma</i>	Kitakoura, Sado, Japan	December-12	LC424822
<i>Pseudoblennius zonostigma</i> -7	<i>Pseudoblennius zonostigma</i>	Tassya, Sado, Japan	November-14	same sequence as LC424822
<i>Pseudoblennius zonostigma</i> -8	<i>Pseudoblennius zonostigma</i>	Kotoura, Sado, Japan	April-15	same sequence as LC424822
<i>Pseudoblennius zonostigma</i> -9	<i>Pseudoblennius zonostigma</i>	Kotoura, Sado, Japan	April-15	same sequence as LC424821
<i>Pseudoblennius zonostigma</i> -12	<i>Pseudoblennius zonostigma</i>	Mushizaki, Sado, Japan	February-16	same sequence as LC424822
<i>Pseudoblennius zonostigma</i> -13	<i>Pseudoblennius zonostigma</i>	Mushizaki, Sado, Japan	September-16	same sequence as LC424821
<i>Pseudoblennius zonostigma</i> -15	<i>Pseudoblennius zonostigma</i>	Kotoura, Sado, Japan	October-16	same sequence as LC424822
<i>Pseudoblennius zonostigma</i> -16	<i>Pseudoblennius zonostigma</i>	Tassya, Sado, Japan	December-16	same sequence as LC424822
<i>Pseudoblennius zonostigma</i> -17	<i>Pseudoblennius zonostigma</i>	Mushizaki, Sado, Japan	December-16	same sequence as LC424822
<i>Pseudoblennius zonostigma</i> -18	<i>Pseudoblennius zonostigma</i>	Mushizaki, Sado, Japan	December-16	same sequence as LC424822
<i>Furcina osimae</i> -16	<i>Furcina osimae</i>	Tassya, Sado, Japan	July-13	LC424823
<i>Furcina osimae</i> -19	<i>Furcina osimae</i>	Tassya, Sado, Japan	August-13	LC424824
<i>Furcina osimae</i> -20	<i>Furcina osimae</i>	Tassya, Sado, Japan	August-13	same sequence as LC424823
<i>Furcina osimae</i> -27	<i>Furcina osimae</i>	Tassya, Sado, Japan	November-13	same sequence as LC424823
<i>Furcina osimae</i> -37	<i>Furcina osimae</i>	Sado, Japan	December-13	same sequence as LC424823
<i>Furcina osimae</i> -38	<i>Furcina osimae</i>	Sado, Japan	December-13	same sequence as LC424823
<i>Furcina osimae</i> -39	<i>Furcina osimae</i>	Sado, Japan	December-13	same sequence as LC424823
<i>Furcina osimae</i> -43	<i>Furcina osimae</i>	Sado, Japan	December-13	same sequence as LC424823
<i>Furcina osimae</i> -44	<i>Furcina osimae</i>	Sado, Japan	December-13	same sequence as LC424823
<i>Furcina osimae</i> -45	<i>Furcina osimae</i>	Sado, Japan	December-13	same sequence as LC424823
<i>Furcina osimae</i> -47	<i>Furcina osimae</i>	Sado, Japan	December-13	same sequence as LC424823
<i>Furcina osimae</i> -48	<i>Furcina osimae</i>	Mushizaki, Sado, Japan	January-14	same sequence as LC424824
<i>Furcina osimae</i> -48	<i>Furcina osimae</i>	Mushizaki, Sado, Japan	January-14	same sequence as LC424824
<i>Furcina osimae</i> -49	<i>Furcina osimae</i>	Tassya, Sado, Japan	December-13	same sequence as LC424823
<i>Furcina osimae</i> -50	<i>Furcina osimae</i>	Tassya, Sado, Japan	January-15	same sequence as LC424823
<i>Furcina osimae</i> -51	<i>Furcina osimae</i>	Tassya, Sado, Japan	January-15	same sequence as LC424824
<i>Furcina osimae</i> -52	<i>Furcina osimae</i>	Tassya, Sado, Japan	January-15	same sequence as LC424823
<i>Furcina osimae</i> -57	<i>Furcina osimae</i>	Tassya, Sado, Japan	November-15	same sequence as LC424823
<i>Furcina osimae</i> -58	<i>Furcina osimae</i>	Tassya, Sado, Japan	November-15	same sequence as LC424823
<i>Furcina osimae</i> -62	<i>Furcina osimae</i>	Kotoura, Sado, Japan	October-16	same sequence as LC424823
<i>Furcina ishikawae</i> -1	<i>Furcina ishikawae</i>	Kotoura, Sado, Japan	November-12	LC424825
<i>Furcina ishikawae</i> -2	<i>Furcina ishikawae</i>	Kitakoura, Sado, Japan	November-12	same sequence as LC424825
<i>Furcina ishikawae</i> -3	<i>Furcina ishikawae</i>	Kitakoura, Sado, Japan	November-12	same sequence as LC424825
<i>Furcina ishikawae</i> -5	<i>Furcina ishikawae</i>	Kitakoura, Sado, Japan	November-12	LC424826
<i>Furcina ishikawae</i> -6	<i>Furcina ishikawae</i>	Kitakoura, Sado, Japan	November-12	LC424827
<i>Furcina ishikawae</i> -7	<i>Furcina ishikawae</i>	Kitakoura, Sado, Japan	November-12	same sequence as LC424825
<i>Furcina ishikawae</i> -8	<i>Furcina ishikawae</i>	Kitakoura, Sado, Japan	November-12	same sequence as LC424825
<i>Furcina ishikawae</i> -9	<i>Furcina ishikawae</i>	Kitakoura, Sado, Japan	November-12	LC424828
<i>Furcina ishikawae</i> -10	<i>Furcina ishikawae</i>	Kitakoura, Sado, Japan	November-12	LC424829
<i>Furcina ishikawae</i> -11	<i>Furcina ishikawae</i>	Kitakoura, Sado, Japan	November-12	LC424830
<i>Furcina ishikawae</i> -12	<i>Furcina ishikawae</i>	Kitakoura, Sado, Japan	November-12	same sequence as LC424826
<i>Furcina ishikawae</i> -13	<i>Furcina ishikawae</i>	Kitakoura, Sado, Japan	November-12	same sequence as LC424826
<i>Furcina ishikawae</i> -14	<i>Furcina ishikawae</i>	Kitakoura, Sado, Japan	November-12	LC424831
<i>Furcina ishikawae</i> -15	<i>Furcina ishikawae</i>	Kitakoura, Sado, Japan	December-12	LC424832
<i>Furcina ishikawae</i> -16	<i>Furcina ishikawae</i>	Kitakoura, Sado, Japan	December-12	same sequence as LC424826
<i>Furcina ishikawae</i> -33	<i>Furcina ishikawae</i>	Kotoura, Sado, Japan	December-15	same sequence as LC424826
<i>Jordania zonope</i> -4	<i>Jordania zonope</i>	Ucluelet, BC, Canada	February-17	LC424833
<i>Jordania zonope</i> -6	<i>Jordania zonope</i>	Ucluelet, BC, Canada	February-17	same sequence as LC424833



**Table S2** Results of genetic identification of parental species of egg masses deposited in colonial tunicates and sponges. Sampling date, locality, depth, water temperature (WT), number of eggs in hosts, number of hatched larvae, the average egg diameter, egg (yolk) coloration and number of young analyzed for genetic species identification are shown. *ND*, no data.

No	Egg mass ID	Fish species	Host ID	Host animal	Host species	Date	Locality	Depth (m)	WT (°C)	Number of eggs	Number of hatched larva	Average egg	Egg (yolk) coloration	Number of analyzed young
												Diameter (mm)		
1	SPE-001	<i>Vellitor centropomus</i>	SP-0001	Colonial tunicate	Didemnidae sp.	02-Apr-12	Kotoura	7.0-9.0	9.2		15	1.4 (N = 10)	Orange	2
2	SPE-002	<i>Vellitor centropomus</i>	SP-0001	Colonial tunicate	Didemnidae sp.	02-Apr-12	Kotoura	7.0-9.0	9.2		124	1.4 (10)	Orange, yellow, clear	3
3	SPE-003	<i>Vellitor centropomus</i>	SP-0001	Colonial tunicate	Didemnidae sp.	02-Apr-12	Kotoura	7.0-9.0	9.2	138	57	1.4 (10)	Orange	3
4	SPE-004	<i>Vellitor centropomus</i>	SP-0002	Colonial tunicate	Didemnidae sp.	02-Apr-12	Kotoura	7.0-9.0	9.2		11	1.3 (10)	Yellow	2
5	SPE-006	<i>Vellitor centropomus</i>	SP-0007	Colonial tunicate	Didemnidae sp.	15-Apr-12	Kotoura	8.0-12.0	9.4	29	16	1.4 (10)	Yellow	3
6	SPE-008	<i>Vellitor centropomus</i>	SP-0014	Colonial tunicate	Didemnidae sp.	15-Apr-12	Kotoura	8.0-12.0	9.4	121	2	1.4 (10)	Yellow	2
7	SPE-011	<i>Vellitor centropomus</i>	SP-0014	Colonial tunicate	Didemnidae sp.	15-Apr-12	Kotoura	8.0-12.0	9.4	95	1	1.4 (10)	Orange	1
8	SPE-014	<i>Vellitor centropomus</i>	SP-0025	Colonial tunicate	Didemnidae sp.	02-May-12	Kotoura	6.0-9.0	13.6	54	1	ND	Yellow	1
9	SPE-015	<i>Vellitor centropomus</i>	SP-0027	Sponge	Demospongiae sp.	05-May-12	Kotoura	4.0-7.0	9.7	6	1	1.3 ( 6)	Yellow	2
10	SPE-016	<i>Vellitor centropomus</i>	SP-0031	Colonial tunicate	Didemnidae sp.	05-May-12	Kotoura	4.0-7.0	9.7	25	0	1.4 (10)	Yellow	1
11	SPE-017	<i>Vellitor centropomus</i>	SP-0031	Colonial tunicate	Didemnidae sp.	05-May-12	Kotoura	4.0-7.0	9.7	33	31	1.3 (10)	Yellow	2
12	SPE-018	<i>Vellitor centropomus</i>	SP-0032	Sponge	Demospongiae sp.	05-May-12	Kotoura	4.0-7.0	9.7	8	8	1.3 ( 5)	Yellow	2
13	SPE-020	<i>Vellitor centropomus</i>	SP-0034	Colonial tunicate	Didemnidae sp.	20-Mar-13	Kotoura	6.0-8.0	10.4	85	85	1.3 (10)	Yellow	3
14	SPE-021	<i>Vellitor centropomus</i>	SP-0035	Colonial tunicate	Didemnidae sp.	20-Mar-13	Kotoura	6.0-8.0	10.4	82	82	1.3 ( 8)	Yellow	3
15	SPE-022	<i>Vellitor centropomus</i>	SP-0036	Colonial tunicate	Didemnidae sp.	20-Mar-13	Kotoura	6.0-8.0	10.4	29	21	1.3 ( 9)	Yellow	3
16	SPE-023A	<i>Vellitor centropomus</i>	SP-0037	Colonial tunicate	Didemnidae sp.	20-Mar-13	Kotoura	6.0-8.0	10.4	40	1	1.3 (10)	Yellow	3
17	SPE-023B	<i>Vellitor centropomus</i>	SP-0037	Colonial tunicate	Didemnidae sp.	20-Mar-13	Kotoura	6.0-8.0	10.4	63	47	1.3 (10)	Yellow	3
18	SPE-024	<i>Vellitor centropomus</i>	SP-0038	Colonial tunicate	Didemnidae sp.	20-Mar-13	Kotoura	6.0-8.0	10.4	79	69	1.2 (10)	Yellow	3
19	SPE-026	<i>Vellitor centropomus</i>	SP-0040	Colonial tunicate	Didemnidae sp.	26-Apr-13	Kotoura	8.0-11.0	10.8	68	29	1.4 (10)	Yellow	2
20	SPE-027A	<i>Vellitor centropomus</i>	SP-0041	Colonial tunicate	Didemnidae sp.	26-Apr-13	Kotoura	8.0-11.0	10.8	66	66	1.4 (10)	Yellow	2
21	SPE-027B	<i>Vellitor centropomus</i>	SP-0041	Colonial tunicate	Didemnidae sp.	26-Apr-13	Kotoura	8.0-11.0	10.8	46	46	1.3 (10)	Yellow	2
22	SPE-029	<i>Vellitor centropomus</i>	SP-0043	Colonial tunicate	Didemnidae sp.	26-Apr-13	Kotoura	8.0-11.0	10.8	19	19	1.3 (10)	Yellow	2
23	SPE-030	<i>Vellitor centropomus</i>	SP-0044	Colonial tunicate	Didemnidae sp.	26-Apr-13	Kotoura	8.0-11.0	10.8	33	33	1.3 (10)	Yellow	3
24	SPE-031	<i>Vellitor centropomus</i>	SP-0045	Colonial tunicate	Didemnidae sp.	26-Apr-13	Kotoura	8.0-11.0	10.8	49	42	1.4 (10)	Yellow	2
25	SPE-032A	<i>Vellitor centropomus</i>	SP-0046	Colonial tunicate	Didemnidae sp.	26-Apr-13	Kotoura	8.0-11.0	10.8	31	23	1.3 (10)	Yellow	3
26	SPE-032B	<i>Vellitor centropomus</i>	SP-0046	Colonial tunicate	Didemnidae sp.	26-Apr-13	Kotoura	8.0-11.0	10.8	53	12	1.3 (10)	Yellow	2
27	SPE-034	<i>Vellitor centropomus</i>	SP-0048	Sponge	Demospongiae sp.	06-Mar-14	Kitakoura	2.0	8.8	21	5	ND	Yellow	2
28	SPE-035	<i>Vellitor centropomus</i>	SP-0049	Colonial tunicate	Didemnidae sp.	13-Apr-14	Kitakoura	2.0	9.9	2	2	1.5 ( 2)	Yellow	1
29	SPE-037	<i>Vellitor centropomus</i>	SP-0051	Sponge	Demospongiae sp.	11-Mar-16	Kotoura	6.0	10.7	7	2	1.4 ( 7)	Yellowish orange	2
30	SPE-039	<i>Vellitor centropomus</i>	SP-0053	Colonial tunicate	Didemnidae sp.	11-Mar-16	Kotoura	6.0	10.7	2	2	1.5 ( 2)	Yellowish orange	2
31	SPE-019	<i>Pseudoblennius marmoratus</i>	SP-0033	Sponge	Demospongiae sp.	05-May-12	Kotoura	4.0-7.0	9.7	10	10	1.6 (10)	Orange	3
32	SPE-025	<i>Pseudoblennius marmoratus</i>	SP-0039	Sponge	Demospongiae sp.	24-Mar-13	Kitakoura	9.0	9.8	33	33	1.6 (10)	Orange	3
33	SPE-033	<i>Pseudoblennius marmoratus</i>	SP-0047	Sponge	Demospongiae sp.	11-Feb-14	Kitakoura	1.0	10.2	16	6	1.5 ( 7)	Orange	1
34	SPE-038	<i>Pseudoblennius marmoratus</i>	SP-0052	Sponge	Demospongiae sp.	11-Mar-16	Kotoura	2.0	10.7	14	1	1.6 ( 7)	Orange	1
35	SPE-005	Not identified	SP-0002	Colonial tunicate	Didemnidae sp.	02-Apr-12	Kotoura	7.0-9.0	9.2		0	1.4 (10)	Yellow	0
36	SPE-007	Not identified	SP-0008	Colonial tunicate	Didemnidae sp.	15-Apr-12	Kotoura	8.0-12.0	9.4	29	0	1.4 (10)	Orange	0
37	SPE-009	Not identified	SP-0014	Colonial tunicate	Didemnidae sp.	15-Apr-12	Kotoura	8.0-12.0	9.4	89	0	1.4 (10)	Yellow	0
38	SPE-010	Not identified	SP-0014	Colonial tunicate	Didemnidae sp.	15-Apr-12	Kotoura	8.0-12.0	9.4	196	0	1.4 (10)	Yellow	0
39	SPE-012	Not identified	SP-0014	Colonial tunicate	Didemnidae sp.	15-Apr-12	Kotoura	8.0-12.0	9.4	44	0	1.5 (10)	Yellow	0
40	SPE-013	Not identified	SP-0024	Colonial tunicate	Didemnidae sp.	02-May-12	Kotoura	6.0-9.0	13.6	109	0	ND	Yellow	0
41	SPE-028	Not identified	SP-0042	Colonial tunicate	Didemnidae sp.	26-Apr-13	Kotoura	8.0-11.0	10.8			Lost		0
42	SPE-036	Not identified	SP-0050	Sponge	Demospongiae sp.	11-Mar-16	Kotoura	6.0	10.7	29	0	1.4 (10)	Yellowish orange	0

**Table S3** Results of genetic identification of parental species of egg masses deposited in solitary tunicates. Sampling date, locality, depth, water temperature (WT), number of eggs in hosts, number of hatched larvae, the average egg diameter, egg (yolk) coloration and number of young analyzed for genetic species identification are shown. *ND*, no data.

No	Egg mass ID	Fish species	Host ID	Host animal	Host species	Date	Locality	Depth (m)	WT (°C)	Number of eggs	Number of hatched larva	Average egg		Number of analyzed young
												Diameter (mm)	Egg (yolk) coloration	
1	STE-004	<i>Pseudoblennius cottoides</i>	ST-0021	Solitary tunicate	<i>Halocynthia ritteri</i>	22-Dec-12	Kitakoura	8.0-18.0	13.9	200	195	1.8 (10)	Yellowish orange	3
2	STE-005	<i>Pseudoblennius cottoides</i>	ST-0050	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-12	Kitakoura	3.0-7.0	13.3	90	86	1.9 (10)	Orange	1
3	STE-006	<i>Pseudoblennius cottoides</i>	ST-0078	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-12	Kitakoura	3.0-7.0	13.3	93	83	1.9 (10)	Yellow	2
4	STE-007	<i>Pseudoblennius cottoides</i>	ST-0078	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-12	Kitakoura	3.0-7.0	13.3	185	176	1.9 (10)	Orange	2
5	STE-008	<i>Pseudoblennius cottoides</i>	ST-0078	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-12	Kitakoura	3.0-7.0	13.3	150	143	1.9 (10)	Yellow	3
6	STE-015	<i>Pseudoblennius cottoides</i>	ST-0156	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.9	13.3	210	208	1.9 (10)	Orange	1
7	STE-018	<i>Pseudoblennius cottoides</i>	ST-0180	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.9	13.3	66	68	1.8 (10)	Yellow	1
8	STE-021	<i>Pseudoblennius cottoides</i>	ST-0191	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.9	13.3	318	257	1.9 (10)	Yellow	2
9	STE-022	<i>Pseudoblennius cottoides</i>	ST-0198	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.9	13.3	161	124	1.9 (10)	Yellow	2
10	STE-025	<i>Pseudoblennius cottoides</i>	ST-0262	Solitary tunicate	<i>Halocynthia ritteri</i>	26-Dec-13	Mushizaki	8.0-17.8	14.3	74	56	1.8 (10)	Yellow	1
11	STE-026	<i>Pseudoblennius cottoides</i>	ST-0270	Solitary tunicate	<i>Halocynthia ritteri</i>	26-Dec-13	Mushizaki	8.0-17.8	14.3	97	91	1.8 (10)	Yellow	3
12	STE-028	<i>Pseudoblennius cottoides</i>	ST-0301	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-4.0	13.3	92	42	1.9 (10)	Yellowish orange	1
13	STE-032	<i>Pseudoblennius cottoides</i>	ST-0375	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.0	14.1		247	1.9 (10)	Yellow	2
14	STE-034	<i>Pseudoblennius cottoides</i>	ST-0381	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.0	14.1		224	1.8 (10)	Yellowish orange	1
15	STE-035	<i>Pseudoblennius cottoides</i>	ST-0391	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.0	14.1	255	115	1.8 (10)	Yellowish orange	2
16	STE-036	<i>Pseudoblennius cottoides</i>	ST-0397	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.0	14.1	256	35	1.8 (10)	Yellow	2
17	STE-038	<i>Pseudoblennius cottoides</i>	ST-0431	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.0	14.1	164		1.9 (10)	Yellow	2
18	STE-040	<i>Pseudoblennius cottoides</i>	ST-0485	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.0	14.1	122	109	1.8 (10)	Orange	2
19	STE-042	<i>Pseudoblennius cottoides</i>	ST-0545b	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-3.9	10.2	440	434	1.8 (10)	Yellow	2
20	STE-043	<i>Pseudoblennius cottoides</i>	ST-0566	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-3.9	10.2	400	344	1.9 (10)	Yellow	2
21	STE-050	<i>Pseudoblennius cottoides</i>	ST-0631	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-4.0	10.2	240	213	1.9 (10)	Orange	2
22	STE-051A	<i>Pseudoblennius cottoides</i>	ST-0635	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-4.0	10.2	179	179	1.8 ( 6)	Yellow	2
23	STE-051B	<i>Pseudoblennius cottoides</i>	ST-0635	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-4.0	10.2	250	178	1.9 (10)	Orange	1
24	STE-052	<i>Pseudoblennius cottoides</i>	ST-0636	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-4.0	10.2	170	162	1.8 (10)	Yellowish orange	2
25	STE-053	<i>Pseudoblennius cottoides</i>	ST-0641	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-4.0	10.2	170	202	1.9 (10)	Orange	1
26	STE-054	<i>Pseudoblennius cottoides</i>	ST-0644	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-4.0	10.2	361	354	2.0 (10)	Orange	1
27	STE-055	<i>Pseudoblennius cottoides</i>	ST-0649	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-4.0	10.2	196	145	1.9 (10)	Orange	1
28	STE-056	<i>Pseudoblennius cottoides</i>	ST-0650	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-4.0	10.2	146	146	1.9 (10)	Yellow	2
29	STE-057	<i>Pseudoblennius cottoides</i>	ST-0654	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-4.0	10.2	85	85	1.8 ( 3)	Yellow	2
30	STE-058	<i>Pseudoblennius cottoides</i>	ST-0655	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-4.0	10.2	172	112	1.9 (10)	Yellowish orange	2
31	STE-059	<i>Pseudoblennius cottoides</i>	ST-0658	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-4.0	10.2	269	267	1.9 (10)	Yellowish orange	2
32	STE-063	<i>Pseudoblennius cottoides</i>	ST-0744	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-3.9	8.8	323	278	1.8 (10)	Orange	1
33	STE-064	<i>Pseudoblennius cottoides</i>	ST-0751	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-3.9	8.8	390	353	1.9 (10)	Yellowish orange	1
34	STE-065	<i>Pseudoblennius cottoides</i>	ST-0754	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-3.9	8.8	291	291	1.9 (10)	Yellow	2
35	STE-066	<i>Pseudoblennius cottoides</i>	ST-0755	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-3.9	8.8	292	158	1.9 (10)	Reddish orange	2
36	STE-067A	<i>Pseudoblennius cottoides</i>	ST-0757	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-3.9	8.8	360	7	1.9 (10)	Yellowish orange	1
37	STE-067B	<i>Pseudoblennius cottoides</i>	ST-0757	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-3.9	8.8	300	300	2.0 (10)	Yellow	1
38	STE-071	<i>Pseudoblennius cottoides</i>	ST-0824	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-4.0	8.8	311	297	1.9 (10)	Orange	1
39	STE-072	<i>Pseudoblennius cottoides</i>	ST-0856	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-4.0	8.8	371	316	2.0 (10)	Orange	2
40	STE-073	<i>Pseudoblennius cottoides</i>	ST-0857	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-4.0	8.8	300	286	1.8 (10)	Orange	1
41	STE-074	<i>Pseudoblennius cottoides</i>	ST-0869	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-4.0	8.8	380	339	1.8 (10)	Orange	1
42	STE-075	<i>Pseudoblennius cottoides</i>	ST-0873	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-4.0	8.8	260	184	1.8 (10)	Yellowish orange	1
43	STE-076	<i>Pseudoblennius cottoides</i>	ST-0884	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-4.0	8.8	400	384	1.9 (10)	Yellowish orange	1
44	STE-077	<i>Pseudoblennius cottoides</i>	ST-0887	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-4.0	8.8	157	138	1.9 (10)	Orange	1
45	STE-080	<i>Pseudoblennius cottoides</i>	ST-0918	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	293	379	1.9 (10)	Orange	2
46	STE-081	<i>Pseudoblennius cottoides</i>	ST-0922	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	312	314	2.0 (10)	Yellowish orange	2
47	STE-082	<i>Pseudoblennius cottoides</i>	ST-0923	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	344	313	1.9 (10)	Yellowish orange	1
48	STE-083A	<i>Pseudoblennius cottoides</i>	ST-0924	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	430	425	1.8 (10)	Yellowish orange	3

49	STE-083B	<i>Pseudoblennius cottoides</i>	ST-0924	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	310	303	1.8 (10)	Orange	1
50	STE-086	<i>Pseudoblennius cottoides</i>	ST-0940	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	400	316	1.9 (10)	Orange	1
51	STE-088	<i>Pseudoblennius cottoides</i>	ST-0944	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	300	240	1.9 (10)	Yellowish orange	2
52	STE-090	<i>Pseudoblennius cottoides</i>	ST-0986	Solitary tunicate	<i>Pyura saciformis</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	206	176	2.0 (10)	Orange	1
53	STE-003	<i>Pseudoblennius zonostigma</i>	ST-0021	Solitary tunicate	<i>Halocynthia ritteri</i>	22-Dec-12	Kitakoura	8.0-18.0	13.9	338	203	1.8 (10)	Orange	2
54	STE-009A	<i>Pseudoblennius zonostigma</i>	ST-0079	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-12	Kitakoura	3.0-7.0	13.3	37	27	2.0 (10)	Orange	3
55	STE-009B	<i>Pseudoblennius zonostigma</i>	ST-0079	Solitary tunicate	<i>Halocynthia ritteri</i>	26-Dec-12	Kitakoura	3.0-7.0	13.3	7	6	1.9 ( 6)	Orange	2
56	STE-011	<i>Pseudoblennius zonostigma</i>	ST-0120	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.9	13.3	290	222	1.8 (10)	Yellow	1
57	STE-014	<i>Pseudoblennius zonostigma</i>	ST-0149	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.9	13.3	163	135	1.7 (10)	Yellowish orange	2
58	STE-016A	<i>Pseudoblennius zonostigma</i>	ST-0172	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.9	13.3	57		1.7 (10)	Orange	1
59	STE-016B	<i>Pseudoblennius zonostigma</i>	ST-0172	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.9	13.3	83	81	1.8 (10)	Yellow	2
60	STE-017	<i>Pseudoblennius zonostigma</i>	ST-0178	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.9	13.3	106	84	1.8 (10)	Orange	1
61	STE-019	<i>Pseudoblennius zonostigma</i>	ST-0185	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.9	13.3	136	137	1.7 (10)	Orange	1
62	STE-020	<i>Pseudoblennius zonostigma</i>	ST-0190	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.9	13.3	52	28	1.8 (10)	Orange	1
63	STE-027	<i>Pseudoblennius zonostigma</i>	ST-0273	Solitary tunicate	<i>Halocynthia ritteri</i>	26-Dec-13	Mushizaki	8.0-17.8	14.3	229	156	1.8 (10)	Yellowish orange	1
64	STE-030	<i>Pseudoblennius zonostigma</i>	ST-0331	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.0	14.1	139	158	1.8 (10)	Orange	1
65	STE-049	<i>Pseudoblennius zonostigma</i>	ST-0623	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-4.0	10.2	140	128	1.8 ( 7)	Yellowish orange	2
66	STE-002	<i>Pseudoblennius</i> sp. "Kirin-anahaze"	ST-0000B	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Dec-11	Tassha	3.0	15.5	450	443	1.9 (10)	Yellow	3
67	STE-010	<i>Pseudoblennius</i> sp. "Kirin-anahaze"		Solitary tunicate	<i>Halocynthia ritteri</i>						525	ND		3
68	STE-013	<i>Pseudoblennius</i> sp. "Kirin-anahaze"	ST-0134	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.9	13.3	702	690	1.9 (10)	Yellow	3
69	STE-024	<i>Pseudoblennius</i> sp. "Kirin-anahaze"	ST-0259	Solitary tunicate	<i>Halocynthia roretzi</i>	26-Dec-13	Mushizaki	8.0-17.8	14.3	655	214	2.0 (10)	Yellow	3
70	STE-029	<i>Pseudoblennius</i> sp. "Kirin-anahaze"	ST-0317	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-4.0	13.3	395	144	2.0 (10)	Yellow	2
71	STE-033	<i>Pseudoblennius</i> sp. "Kirin-anahaze"	ST-0379	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.0	14.1	1200	642	2.0 (10)	Yellow	2
72	STE-037	<i>Pseudoblennius</i> sp. "Kirin-anahaze"	ST-0399	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.0	14.1	870	15	2.0 (10)	Yellow	2
73	STE-044	<i>Pseudoblennius</i> sp. "Kirin-anahaze"	ST-0567	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-3.9	10.2	1000	913	2.0 (10)	Yellow	2
74	STE-046	<i>Pseudoblennius</i> sp. "Kirin-anahaze"	ST-0576	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-3.9	10.2	680	624	2.0 (10)	Yellow	2
75	STE-023	<i>Pseudoblennius percoides</i>	ST-0222	Solitary tunicate	<i>Pyura saciformis</i>	25-Dec-13	Kitakoura	0.5-3.9	13.3	147	125	1.9 (10)	Yellowish green	2
76	STE-031	<i>Pseudoblennius percoides</i>	ST-0369	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.0	14.1	37	23	1.9 (10)	Yellow	1
77	STE-045	<i>Pseudoblennius percoides</i>	ST-0574	Solitary tunicate	<i>Halocynthia roretzi</i>	11-Feb-14	Kitakoura	0.5-3.9	10.2	121	121	1.8 (10)	Yellowish orange	2
78	STE-047	<i>Pseudoblennius percoides</i>	ST-0588	Solitary tunicate	<i>Halocynthia ritteri</i>	11-Feb-14	Kitakoura	0.5-3.9	10.2	161	106	1.9 (10)	Orange	2
79	STE-060	<i>Pseudoblennius percoides</i>	ST-0728	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-3.9	8.8	89		1.9 (10)	Yellow	2
80	STE-062	<i>Pseudoblennius percoides</i>	ST-0735	Solitary tunicate	<i>Pyura saciformis</i>	06-Mar-14	Kitakoura	0.5-3.9	8.8		107	1.9 (10)	Yellowish green	2
81	STE-068	<i>Pseudoblennius percoides</i>	ST-0773	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-3.9	8.8	140	140	2.0 (10)	Yellowish green	2
82	STE-078	<i>Pseudoblennius percoides</i>	ST-0910	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-4.0	8.8	80	80	2.0 (10)	Yellowish green	1
83	STE-048	<i>Furcina osimae</i>	ST-0612	Solitary tunicate	<i>Pyura saciformis</i>	11-Feb-14	Kitakoura	0.5-3.9	10.2	73	72	1.6 (10)	Reddish orange	2
84	STE-069	<i>Furcina osimae</i>	ST-0786	Solitary tunicate	<i>Pyura saciformis</i>	06-Mar-14	Kitakoura	0.5-3.9	8.8	75	75	1.6 (10)	Red	2
85	STE-070	<i>Furcina osimae</i>	ST-0794	Solitary tunicate	<i>Pyura saciformis</i>	06-Mar-14	Kitakoura	0.5-3.9	8.8	83	83	1.7 (10)	Red	2
86	STE-001	<i>Furcina ishikawae</i>	ST-0000A	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Dec-11	Tassha	2.0	15.5	35	31	ND	Reddish orange	2
87	STE-012	Not identified	ST-0130	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.9	13.3	26	0	Dead	White (dead)	0
88	STE-039	Not identified	ST-0435	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.0	14.1	250	0	Dead	White (dead)	0
89	STE-041	Not identified	ST-0491	Solitary tunicate	<i>Halocynthia ritteri</i>	25-Dec-13	Kitakoura	0.5-3.0	14.1	300	0	Dead	White (dead)	0
90	STE-061	<i>Aulichthys japonicas</i>	ST-0731	Solitary tunicate	<i>Halocynthia ritteri</i>	06-Mar-14	Kitakoura	0.5-3.9	8.8	58	48	2.6 (10)	Orange	-
91	STE-079	<i>Aulichthys japonicas</i>	ST-0863	Solitary tunicate	<i>Halocynthia roretzi</i>	06-Mar-14	Kitakoura	0.5-4.0	8.8		208	ND	Orange	-
92	STE-084	<i>Aulichthys japonicas</i>	ST-0938	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	64	69	2.5 (10)	Orange	-
93	STE-085	<i>Aulichthys japonicas</i>	ST-0939	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	110	81	2.6 (10)	Orange	-
94	STE-087	<i>Aulichthys japonicas</i>	ST-0943	Solitary tunicate	<i>Pyura saciformis</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	95	133	2.6 (10)	Orange	-
95	STE-089A	<i>Aulichthys japonicas</i>	ST-0948	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	38	46	2.6 (10)	Reddish orange	-
96	STE-089B	<i>Aulichthys japonicas</i>	ST-0948	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	59	56	2.6 (10)	Orange	-
97	STE-091	<i>Aulichthys japonicas</i>	ST-0987	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	61	49	2.7 (10)	Orange	-
98	STE-092	<i>Aulichthys japonicas</i>	ST-0991	Solitary tunicate	<i>Pyura saciformis</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	94	106	2.7 (10)	Reddish orange	-
99	STE-093	<i>Aulichthys japonicas</i>	ST-0992	Solitary tunicate	<i>Pyura saciformis</i>	13-Apr-14	Kitakoura	0.5-3.9	9.9	72	73	2.6 (10)	Orange	-
100	STE-094	<i>Aulichthys japonicas</i>	ST-1021	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-4.0	9.9	71	75	2.6 (10)	Orange	-
101	STE-095	<i>Aulichthys japonicas</i>	ST-1043	Solitary tunicate	<i>Pyura saciformis</i>	13-Apr-14	Kitakoura	0.5-4.0	9.9	70	0	2.6 (10)	Orange	-
102	STE-096	<i>Aulichthys japonicas</i>	ST-1051	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-4.0	9.9	72	0	2.7 (10)	Orange	-
103	STE-097A	<i>Aulichthys japonicas</i>	ST-1086	Solitary tunicate	<i>Pyura saciformis</i>	13-Apr-14	Kitakoura	0.5-4.0	9.9	59	63	2.8 (10)	Orange	-
104	STE-097B	<i>Aulichthys japonicas</i>	ST-1086	Solitary tunicate	<i>Pyura saciformis</i>	13-Apr-14	Kitakoura	0.5-4.0	9.9	55	27	2.8 (10)	Reddish orange	-
105	STE-097C	<i>Aulichthys japonicas</i>	ST-1086	Solitary tunicate	<i>Pyura saciformis</i>	13-Apr-14	Kitakoura	0.5-4.0	9.9	68	54	2.8 (10)	Orange	-
106	STE-098	<i>Aulichthys japonicas</i>	ST-1087	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-4.0	9.9	62	0	2.8 (10)	Orange	-
107	STE-099	<i>Aulichthys japonicas</i>	ST-1091	Solitary tunicate	<i>Halocynthia ritteri</i>	13-Apr-14	Kitakoura	0.5-4.0	9.9	59	33	2.7 (10)	Orange	-
108	STE-100	<i>Aulichthys japonicas</i>	ST-1126	Solitary tunicate	<i>Halocynthia ritteri</i>	25-May-14	Kitakoura	0.5-3.9	14.4	60	0	2.3 ( 6)	Reddish orange	-
109	STE-101	<i>Aulichthys japonicas</i>	ST-1144	Solitary tunicate	<i>Pyura saciformis</i>	25-May-14	Kitakoura	0.5-3.9	14.4	59	0	2.3 ( 7)	Orange	-

## Supporting information

**Figure S1** Maximum likelihood phylogeny of nine species of potentially ostracophilous sculpins, with *Jordania zonope* as an outgroup, as reconstructed by MEGA ver. 7.0.14. using 800 bp *cytb* sequences, with the settings of the Tamura Nei model and 1000 bootstrap replicates. Values at nodes are bootstrap percentages.


**Figure S2** Maximum likelihood phylogeny of the potential parental species and 205 eggs from 113 egg masses as reconstructed by MEGA ver. 7.0.14. using 800 bp *cytb* sequences, with the settings of the Tamura Nei model and 1000 bootstrap replicates. Values at nodes are bootstrap percentages.

**Figure S3** Maximum likelihood phylogeny of the potential parental species and 18 eggs from 14 egg masses as reconstructed by MEGA ver. 7.0.14., using 628 bp *cytb* sequences, with the settings of Tamura Nei model and 1000 bootstrap replicates. Values at nodes are bootstrap percentages.


**Movie S1** Spawning behaviours of female *Pseudoblennius cottoides* in the aquarium. Six *Halocynthia ritteri* on wire-netting were placed in a 75 × 45 × 45 cm aquarium. Eggs were deposited in the atrium of a tunicate through the atrial siphon using extremely long ovipositors. Eggs were probably released into the atrium when the female opened her mouth. The female stayed still on the tunicate after spawning. She ejected her ovipositor 1 min 22 sec and left the tunicate 2 min 34 sec after the insert, respectively (not shown). This movie was filmed by H. S. on Feb. 7, 2015. Fish and tunicates were collected on the coast of Sado Island in the Sea of Japan.


**Movie S2** Failure of spawning by female *Pseudoblennius cottoides* in the aquarium. The atrial siphon of the tunicate closed before the sculpin female inserted her ovipositor. Settings were the same as in Movie S1. This movie was filmed by H. S. on Feb. 1, 2015.

**Movie S3** Spawning behaviours of female *Pseudoblennius marmoratus* in the field. Eggs were deposited in the spongocoel of a sponge using short ovipositors. This movie was filmed by Tomonobu Uryu on Dec. 28, 2009 at Izu Oceanic Park, Izu Peninsula, Pacific coast of Japan. Water temperature was about 16 °C and water depth was about 5 m.




**Figure S1** Maximum likelihood phylogeny of nine species of potentially ostracophilous sculpins, with *Jordania zonope* as an outgroup, as reconstructed by MEGA ver. 7.0.14. using 800 bp *cytb* sequences, with the settings of the Tamura Nei model and 1000 bootstrap replicates. Values at nodes are bootstrap percentages.





**Figure S2** Maximum likelihood phylogeny of the potential parental species and 205 eggs from 113 egg masses as reconstructed by MEGA ver. 7.0.14. using 800 bp *cytb* sequences, with the settings of the Tamura Nei model and 1000 bootstrap replicates. Values at nodes are bootstrap percentages.





**Figure S3** Maximum likelihood phylogeny of the potential parental species and 18 eggs from 14 egg masses as reconstructed by MEGA ver. 7.0.14., using 628 bp *cytb* sequences, with the settings of Tamura Nei model and 1000 bootstrap replicates. Values at nodes are bootstrap percentages.



**Movie S1** Spawning behaviours of female *Pseudoblennius cottoides* in the aquarium. Six *Halocynthia ritteri* on wire-netting were placed in a  $75 \times 45 \times 45$  cm aquarium. Eggs were deposited in the atrium of a tunicate through the atrial siphon using extremely long ovipositors. Eggs were probably released into the atrium when the female opened her mouth. The female stayed still on the tunicate after spawning. She ejected her ovipositor 1 min 22 sec and left the tunicate 2 min 34 sec after the insert, respectively (not shown). This movie was filmed by H. S. on Feb. 7, 2015. Fish and tunicates were collected on the coast of Sado Island in the Sea of Japan.



**Movie S2** Failure of spawning by female *Pseudoblennius cottoides* in the aquarium. The atrial siphon of the tunicate closed before the sculpin female inserted her ovipositor. Settings were the same as in Movie S1. This movie was filmed by H. S. on Feb. 1, 2015.



**Movie S3** Spawning behaviours of female *Pseudoblennius marmoratus* in the field. Eggs were deposited in the spongocoel of a sponge using short ovipositors. This movie was filmed by Tomonobu Uryu on Dec. 28, 2009 at Izu Oceanic Park, Izu Peninsula, Pacific coast of Japan. Water temperature was about 16 °C and water depth was about 5 m.